

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
29 December 2004 (29.12.2004)

PCT

(10) International Publication Number  
**WO 2004/113522 A1**

(51) International Patent Classification<sup>7</sup>: **C12N 9/56**,  
9/50, 9/64, 9/00, A61K 38/48

(21) International Application Number:  
PCT/EP2004/051173

(22) International Filing Date: 18 June 2004 (18.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
03013819.2 18 June 2003 (18.06.2003) EP  
03025851.1 10 November 2003 (10.11.2003) EP  
03025871.9 11 November 2003 (11.11.2003) EP  
04003058.7 11 February 2004 (11.02.2004) EP

(71) Applicant (for all designated States except US): **DIREVO BIOTECH AG** [DE/DE]; Nattermannallee 1, 50667 Köln (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HAUPTS, Ulrich** [DE/DE]; c/o Direvo Biotech AG, 50667 Köln (DE). **KOLTERMANN, Andre** [DE/DE]; c/o Direvo Biotech AG, 50667 Köln (DE). **SCHEIDIG, Andreas** [DE/DE]; c/o Direvo Biotech AG, 50667 Köln (DE). **VOETSMEIER, Christian** [DE/DE]; c/o Direvo Biotech AG, 50667 Köln (DE). **KETTLING, Ulrich** [DE/DE]; c/o Direvo Biotech AG, 50667 Köln (DE).

(74) Agents: **HELBING, Jörg** et al.; Von Kreisler Selting Werner, Postfach 10 22 41, 50462 Köln (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NEW BIOLOGICAL ENTITIES AND THE PHARMACEUTICAL OR DIAGNOSTIC USE THEREOF

(57) Abstract: The present invention provides method for the treatment of a disease by applying a medicament comprising a protease with a defined specificity is capable to hydrolyze specific peptide bonds within a target substrate related to such disease. The proteases with such a defined specificity can further be used for related therapeutic or diagnostic purposes.

WO 2004/113522 A1

## New Biological Entities and the Pharmaceutical or Diagnostic Use Thereof

The present invention provides methods for the treatment of a disease by applying a medicament comprising a protease with a defined specificity is capable to hydrolyze specific peptide bonds within a target substrate related to such disease. The proteases with such a defined specificity can further be used for related therapeutic or diagnostic purposes.

### Background

Academic and industrial research continuously searches for functional proteins to be used as therapeutic, research, diagnostic, nutritional, personal care or industrial agents. Today, such functional proteins can be classified mainly into two categories: natural proteins and engineered proteins. Natural proteins, on the one hand, are discovered from nature, e.g. by screening natural isolates or by sequencing genomes from diverse species. Engineered proteins, on the other hand, are typically based on known proteins and are altered in order to acquire modified functionalities. The present invention discloses engineered proteins with novel functions as compared to the starting components. Such proteins are called NBEs (New Biologic Entities). The NBEs disclosed in the present invention are engineered enzymes with novel substrate specificities or fusion proteins of such engineered enzymes with other functional components.

Specificity is an essential element of enzyme function. A cell consists of thousands of different, highly reactive catalysts. Yet the cell is able to maintain a coordinated metabolism and a highly organized three-dimensional structure. This is due in part to the specificity of enzymes, i.e. the selective conversion of their respective substrates. Specificity is a qualitative and a quantitative property: the specificity of a particular enzyme can vary widely, ranging from just one particular type of target molecules to all molecular types with certain chemical substructures. In nature, the specificity of an organism's enzymes has been evolved to the particular needs of the organism. Arbitrary specificities with high value for therapeutic, research, diagnostic, nutritional or industrial applications are unlikely to be found in any organism's enzymatic repertoire due to the large

space of possible specificities. The only realistic way of obtaining such specificities is their generation de novo.

When comparing enzymes with binders, a paradigm of specificity is given by antibodies recognizing individual epitopes as small distinct structures within large molecules. The naturally occurring vast range of antibody specificities is attributed to the diversity generated by the immune system combined with natural selection. Several mechanisms contribute to the vast repertoire of antibody specificity and occur at different stages of immune response generation and antibody maturation (Janeway, C et al. (1999) Immunobiology, Elsevier Science Ltd., Garland Publishing, New York). Specifically, antibodies contain complementarity determining regions (CDRs) which interact with the antigen in a highly specific manner and allow discrimination even between very similar epitopes. The light as well as the heavy chain of the antibody each contribute three CDRs to the binding domain. Nature uses recombination of various gene segments combined with further mutagenesis in the generation of CDRs. As a result, the sequences of the six CDR loops are highly variable in composition and length and this forms the basis for the diversity of binding specificities in antibodies. A similar principle for the generation of a diversity of catalytic specificities is not known from nature.

Catalysis, i.e. the increase of the rate of a specific chemical reaction, is besides binding the most important protein function. Catalytic proteins, i.e. enzymes, are classified according to the chemical reaction they catalyze.

Transferases are enzymes transferring a group, for example, the methyl group or a glycosyl group, from one compound (generally regarded as donor) to another compound (generally regarded as acceptor). For example, glycosyltransferases (EC 2.4) transfer glycosyl residues from a donor to an acceptor molecule. Some of the glycosyltransferases also catalyze hydrolysis, which can be regarded as transfer of a glycosyl group from the donor to water. The subclass is further subdivided into hexosyltransferases (EC 2.4.1), pentosyltransferases (EC 2.4.2) and those transferring other glycosyl groups (EC 2.4.99, Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (NC-IUBMB)).

Oxidoreductases catalyze oxido-reductions. The substrate that is oxidized is regarded as hydrogen or electron donor. Oxidoreductases are classified as dehydrogenases, oxidases, mono- and dioxygenases. Dehydrogenases transfer hydrogen from a hydrogen donor to a hydrogen acceptor molecule. Oxidases react with molecular oxygen as hydrogen acceptor and produce oxidized products as well as either hydrogen peroxide or water. Monooxygenases transfer one oxygen atom from molecular oxygen to the substrate and one is reduced to water. In contrast, dioxygenases catalyze the insert of both oxygen atoms from molecular oxygen into the substrate.

Lyases catalyze elimination reactions and thereby generate double bonds or, in the reverse direction, catalyze the additions at double bonds. Isomerases catalyze intramolecular rearrangements. Ligases catalyze the formation of chemical bonds at the expense of ATP consumption.

Finally, hydrolases are enzymes that catalyze the hydrolysis of chemical bonds like C-O or C-N. The E.C. classification for these enzymes generally classifies them by the nature of the bond hydrolysed and by the nature of the substrate. Hydrolases such as lipases and proteases play an important role in nature as well in technical applications of biocatalysts. Proteases hydrolyse a peptide bond within the context of an oligo- or polypeptide. Depending on the catalytic mechanism proteases are grouped into aspartic, serin, cysteine, metallo- and threonine proteases (Handbook of proteolytic enzymes. (1998) Eds: Barret, A; Rawling, N.; Woessner, J.; Academic Press, London). This classification is based on the amino acid side chains that are responsible for catalysis and which are typically presented in the active site in very similar orientation to each other. The scissile bond of the substrate is brought into register with the catalytic residues due to specific interactions between the amino acid side chains of the substrate and complementary regions of the protease (Perona, J. & Craik, C (1995) *Protein Science*, 4, 337-360). The residues on the N- and C-terminal side of the scissile bond are usually called  $P_1$ ,  $P_2$ ,  $P_3$  etc and  $P_1'$ ,  $P_2'$ ,  $P_3'$  and the binding pockets complementary to the substrate  $S_1$ ,  $S_2$ ,  $S_3$  and  $S_1'$ ,  $S_2'$ ,  $S_3'$ , respectively (nomenclature according to Schlechter & Berger, Biochem. Biophys. Res. Commun. 27 (1967) 157-162). The selectivity of proteases can vary widely from



being virtually nonselective – e.g. the Subtilisins – over a strict preference at the  $P_1$  position – e.g. Trypsin selectively cutting on the C-terminal side of arginine or lysine residues – to highly specific proteases – e.g. human tissue-type plasminogen activator (t-PA) cleaving at the C-terminal side of the arginine in the sequence CPGRVVG (Ding, L et al. (1995) *Proc. Natl. Acad. Sci. USA* 92, 7627-7631; Coombs, G et al. (1996) *J. Biol. Chem.* 271, 4461-4467).

The specificity of proteases, i.e. their ability to recognize and hydrolyze preferentially certain peptide substrates, can be expressed qualitatively and quantitatively. Qualitative specificity refers to the kind of amino acid residues that are accepted by a protease at certain positions of the peptide substrate. For example, trypsin and t-PA are related with respect to their qualitative specificity, since both of them require at the  $P_1$  position an arginine or a similar residue. On the other hand, quantitative specificity refers to the relative number of peptide substrates that are accepted as substrates by the protease, or more precisely, to the relative  $k_{cat}/K_M$  ratios of the protease for the different peptides that are accepted by the protease. Proteases that accept only a small portion of all possible peptides have a high specificity, whereas the specificity of proteases that, as an extreme, cleave any peptide substrate would theoretically be zero.

Comparison of the primary, secondary as well as the tertiary structure of proteases (Fersht, A., *Enzyme Structure and Mechanism*, W. H. Freeman and Company, New York, 1995) allows identification of classes showing a high degree of conservation (Rawlings, N.D. & Barrett, A.J. (1997) In: *Proteolysis in Cell Functions* Eds. Hopsu-Havu, V.K.; Järvinen, M.; Kirschke, H, pp. 13-21, IOS Press, Amsterdam). A widely accepted scheme for protease classification has been proposed by Rawlings & Barrett (*Handbook of proteolytic enzymes*. (1998) Eds: Barret, A; Rawling, N.; Woessner, J.; Academic Press, London). For example, the serine proteases family can be subdivided into structural classes with chymotrypsin (class S1), subtilisin (class S8) and carboxypeptidase (class SC) folds, each of which includes nonspecific as well as specific proteases (Rawlings, N.D. & Barrett, A.J. (1994) *Methods Enzymol.* 244, 19-61). This applies to other protease families analogously. An additional distinction can be made according to the relative location of the cleaved bond in the substrate. Carboxy- and

aminopeptidases cleave amino acids from the C- and N-terminus, respectively, while endopeptidases cut anywhere along the oligopeptide.

Many applications would be conceivable if enzymes with a basically unlimited spectrum of specificities were available. However, the use of such enzymes with high, low or any defined specificity is currently limited to those which can be isolated from natural sources. The field of application for these enzymes varies from therapeutic, research, diagnostic, nutritional to personal care and industrial purposes.

Enzyme additives in detergents have come to constitute nearly a third of the whole industrial enzyme market. Detergent enzymes include proteinases for removing organic stains, lipases for removing greasy stains, amylases for removing residues of starchy foods and cellulases for restoring of smooth surface of the fiber. The best-known detergent enzyme is probably the nonspecific proteinase subtilisin, isolated from various *Bacillus* species.

Starch enzymes, such as amylases, occupy the majority of those used in food processing. While starch enzymes include products that are important for textile desizing, alcohol fermentation, paper and pulp processing, and laundry detergent additives, the largest application is for the production of high fructose corn syrup. The production of corn syrup from starch by means of industrial enzymes was a successful alternative to acid hydrolysis.

Apart from starch processing, enzymes are used for an increasing range of applications in food. Enzymes in food can improve texture, appearance and nutritional value or may generate desirable flavours and aromas. Currently used food enzymes in bakery are amylase, amyloglycosidases, pentosanases for breakdown of pentosan and reduced gluten production or glucose oxidases to increase the stability of dough. Common enzymes for dairy are rennet (protease) as coagulant in cheese production, lactase for hydrolysis of lactose, protease for hydrolysis of whey proteins or catalase for the removal of hydrogen peroxides. Enzymes used in brewing process are the above named amylases, but also cellulases or proteases to clarify the beer from suspended proteins. In wines and fruit juices, cloudiness is more commonly caused by starch and pectins so that

amylases and pectinases increase yield and clarification. Papain and other proteinases are used for meat tenderizing.

Enzymes have also been developed to aid animals in the digestion of feed. In the western hemisphere, corn is a major source of food for cattle, swine, and poultry. In order to improve the bioavailability of phosphate from corn, phytase is commonly added (Wyss, M. et al., Biochemical characterization of fungal phytases (myo-inositol hexakisphosphate phosphohydrolases); Catalytic properties. Applied & Environmental Microbiology 65, 367-373 (1999)). Moreover, phytate hydrolysis has been shown to bring about improvements in digestibility of protein and absorption of minerals such as calcium (Bedford, M.R. & Schulze, H., Exogenous Enzymes for Pigs and Poultry [Review]. Nutrition Research Reviews 11, 91-114 (1998)). Another major feed enzyme is xylanase. This enzyme is particularly useful as a supplement for feeding stuff comprising more than about 10% of wheat barley or rye, because of their relatively high soluble fiber content. Xylanases cause two important actions: reduction of viscosity of the intestinal contents by hydrolyzing the gel-like high molecular weight arabinoxylans in feed (Murphy, T et al., Effect of range of new xylanases on in vitro viscosity and on performance of broiler diets. British Poultry Science 44, S16-S18 (2003)) and break down of polymers in cell walls which improve the bioavailability of protein and starch.

Biotech research and development laboratories routinely use special enzymes in small quantities along with many other reagents. These enzymes create a significant market for various enzymes. Enzymes like alkaline phosphatase, horseradish peroxidase and luciferase are only some examples. Thermostable DNA polymerases like Taq polymerase or restriction endonucleases revolutionized laboratory work.

The use of enzymes in the diagnosis of disease is another important benefit derived from the intensive research in biochemistry. Within the recent past few years that interest in diagnostic enzymology has increased and there are still large areas of medical research in which the diagnostic potential of enzyme reactions has not been explored at all. Common enzymes used for clinical diagnosis are acid phosphatase, alanine aminotransferase, alkaline phosphatase,

amylase, angiotensin converting enzymes, aspartate aminotransferase, cholinesterase, creatinine kinase, gamma glutamyltransferase, lactate dehydrogenase or rennin.

Therapeutic enzymes are a particular class of drugs, categorized by the FDA as biologicals, with a lot of advantages compared to other, especially non-biological pharmaceuticals. Examples for successful therapeutic enzymes are human clotting factors like factor VIII and factor IX for human treatment. In addition, digestive enzymes are used for various deficiencies in human digestive processes. Other examples are t-PA and streptokinase for the treatment of cardiovascular disease, beta-glucocerebrosidase for the treatment of Type I Gaucher disease, L-asparaginase for the treatment of acute lymphoblastic leukemia and DNase for the treatment of cystic fibrosis. An important issue in the application of proteins as therapeutics is their potential immunogenicity. To reduce this risk, one would prefer enzymes of human origin, which narrows down the set of available enzymes. The provision of designed enzymes, preferably of human origin, with novel, tailor-made specificities would allow the specific modification of target substrates at will, while minimizing the risk of immunogenicity. A further advantage of highly specific enzymes as therapeutics would be their lower risk of side effects. Due to the limited possibility of specific interactions between a small molecule and a protein, binding to non-target proteins and therefore side effects are quite common and often cause termination of an otherwise promising lead compound. Specific enzymes, on the other hand, provide many more contact sites and mechanisms for substrate discrimination and therefore enable a higher specificity and thereby less side activities.

Proteases represent an important class of therapeutic agents (*Drugs of today*, 33, 641-648 (1997)). However, currently the therapeutic protease is usually a substitute for insufficient activity of the body's own proteases. For example, factor VII can be administered in certain cases of coagulation deficiencies of bleeders or during surgery (Heuer L.; Blumenberg D. (2002) *Anaesthetist* 51:388). Tissue-type plasminogen activator (t-PA) is applied in acute cardiac infarction, initializing the dissolution of fibrin clots through specific cleavage and activation of plasminogen (Verstraete, M. et al. (1995) *Drugs*, 50, 29-41). So far

a protease with tailor-made specificity is generated to provide a therapeutic agent that specifically activates or inactivates a disease related target protein.

Monoclonal antibodies represent another important biological class of substances with therapeutic capabilities. One of the main antibody targets are tumor necrosis factors (TNFs) which belong to the family of cytokines. TNFs play a major role in the inflammation process. As homotrimers they could bind to receptors of nearly every cell. They activate a multiplicity of cellular genes, multiple signal transduction mechanisms, kinases and transcription factors. The most important TNFs are TNF-alpha and TNF-beta. TNF-alpha is produced by macrophages, monocytes and other cells. TNF-alpha is an inflammation mediator. Therefore, research of the last decade has been focused on TNF-alpha inhibitors like monoclonal antibodies as possible therapeutics for different therapeutic indications like Rheumatoid Arthritis, Crohn's disease or Psoriasis (Hamilton et al. (2000) *Expert Opin Pharmacother*, 1 (5): 1041-1052). One of the major disadvantages of monoclonal antibodies are their high costs, so that new biological alternatives are of great importance.

There are a lot of examples for engineered enzymes in literature. Fulani et al. (Fulani F. et al. (2003) *Protein Engineering* 16, 515-519) describe a rhodanase (thiosulfat:cyanide sulfurtransferase) from *Azotobacter vinelandii* which has a catalytic domain structurally related to catalytic subunit of Cdc25 phosphatase enzymes. The difference in catalytic mechanism depends on the different size of the active site. Both rhodanase and phosphatase are highly specific on different substrates (sulfate vs. phosphate). The catalytic mechanism of the rhodanase could be shifted towards serine/threonine phosphatase by single-residue insertion. Therefore, Fulani et al. give a single example for the change of a catalytic mechanism by structural comparison and sequence alignment of naturally known enzymes from different enzyme classes but lack an indication of how to generate a user-definable substrate specificity while keeping the same catalytic mechanism.

The thioredoxin reductase described by Briggs et al. (WO 02/090300 A2) has an altered cofactor specificity which preferably binds NADPH compared to NADH. Thus, both enzymes, the starting point as well as the resulting engineered

enzyme are highly specific towards different substrates. The methods to achieve such an altered substrate specificity are either computational processing methods or sequence alignments of related proteins to define variable and conserved residues. They all have in common that they are based on the comparison of structures and sequences of proteins with known specificities followed by the transfer of the same to another backbone.

There are other examples of specificity-engineered enzymes and, in particular, of proteases which have been published in the literature. None of these examples, however, provides a means for generating novel specificities compared to the specificity of the starting material used within the described methods. The methods range from structure-directed single point mutations (Kurth, T. et al. (1998) *Biochemistry* 37, 11434-11440; Ballinger, M et al. (1996) *Biochemistry*, 35:13579-13585), exchange of surface loops between two specific proteases (Horrevoets et al. (1993) *J. Biol. Chem.* 268, 779-782), to random mutagenesis either regio-selectively or across the whole gene combined with in-vitro or in-vivo selection (Sices, H. & Kristie, T. (1998) *Proc. Natl. Acad. Sci. USA*, 95, 2828-2833).

The rational design of protease specificity is limited to very few examples. This approach is severely limited by the insufficient understanding of the complexities that govern folding and dynamics as well as structure-function relationships in proteins (Corey, M.J. & Corey, E. (1996) *Proc. Natl. Acad. Sci. USA*, 93:11428-11434). It is therefore difficult to alter the primary amino acid sequence of a protease in order to change its activity or specificity in a predictive way. In a successful example, Kurth et al. engineered trypsin to show a preference for a dibasic motive (Kurth, T. et al. (1998) *Biochemistry*, 37:11434-11440). In another example, Hedstrom et al. converted the S<sub>1</sub> substrate specificity of trypsin to that of chymotrypsin (Hedstrom, L. et al. (1992) *Science*, 255:1249-1253). This is an example where a known property was transferred from one backbone to another.

Ballinger et al. (WO 96/27671) describe subtilisin variants with combination mutations (N62D/G166D, and optionally Y104D) having a shift of substrate specificity towards peptide or polypeptide substrates with basic amino acids at

the P1, P2 and P4 positions of the substrate. Suitable substrates of the variant subtilisin were revealed by sorting a library of phage particles (substrate phage) containing five contiguous randomized residues. These subtilisin variants are useful for cleaving fusion proteins with basic substrate linkers and processing hormones or other proteins (in vitro or in vivo) that contain basic cleavage sites. The problems associated with rational redesign of enzymes can partially be overcome by directed evolution (as disclosed in PCT/EP03/04864). These studies can be classified by their expression and selection systems. Genetic selection means to produce inside an organism an enzyme, e.g. a protease, which is able to cleave a precursor protein which in turn results in an alteration of the growth behavior of the producing organism. From a population of organisms with different proteases those can be selected which have an altered growth behavior. This principle was for example reported by Davis et al. (US 5258289, WO 96/21009). The production of a phage system is dependent on the cleavage of a phage protein which only can be activated in the presence of a proteolytic enzyme which is able to cleave the phage protein. Other approaches use a reporter system which allows a selection by screening instead of a genetic selection, but also cannot overcome the intrinsic insufficiency of the intracellular characterization of enzymes.

Systems to generate enzymes with altered sequence specificities with self-secreting enzymes are also reported. Duff et al. (WO 98/11237) describe an expression system for a self-secreting protease. An essential element of the experimental design is that the catalytic reaction acts on the protease itself by an autoproteolytic processing of the membrane-bound precursor molecule to release the matured protease from the cellular membrane into the extracellular environment. Therefore, a fusion protein must be constructed where the target peptide sequence replaces the natural cleavage site for autoproteolysis. Limitations of such a system are that positively identified proteases will have the ability to cleave a certain amino acid sequence but they also may cleave many other peptide sequences. Therefore, high substrate specificity cannot be achieved. Additionally, such a system is not able to control that selected proteases cleave at a specific position in a defined amino acid sequence and it does not allow a precise characterization of the kinetic constants of the selected proteases ( $k_{\text{cat}}$ ,  $K_M$ ).

A method has been described that aims at the generation of new catalytic activities and specificities within the  $\alpha/\beta$ -barrel proteins (WO 01/42432; Fersht et al, Methods of producing novel enzymes; Altamirano et al. (2000) *Nature* 403, 617-622). The  $\alpha/\beta$ -barrel proteins comprise a large superfamily of proteins accounting for a large fraction of all known enzymes. The structure of the proteins is made from a  $\beta$ -barrel surrounded by  $\alpha$ -helices. The loops connecting  $\beta$ -strands and helices comprise the so-called lid-structure including the active site residues. The method is based on the classification of  $\alpha/\beta$ -barrel proteins into two classes based on the catalytic lid structure. An extensive comparison of  $\alpha/\beta$ -barrel protein structures led the authors to the conclusion that the substrate binding and specificity is primarily defined by the barrel structure while the specificity of the chemical reaction resides within the loops. It is suggested that barrels and lid structures from different enzymes can be combined to generate new enzymatic activities and to provide a starting point to fine tune the properties by targeted or randomized mutagenesis and selection. The method does not provide for the generation of user-defined specificity.

In summary, it is clear that there are many possible applications in the fields of therapeutics, research and diagnostics, industrial enzymes, food and feed processing, cosmetics and other areas that would become possible by the availability of enzymes with a novel substrate specificity. However, only a limited number of specific enzymes has been identified from natural sources so far. Methods of rational design to modify, alter, convert or transfer sequence specificity as well as random approaches described above did not enable the generation of a novel and user-definable specificity that was not present in the employed starting material.

Therefore, none of the currently available methods can provide enzymes with a novel and user-defined sequence specificity. In contrast, the current invention provides such enzymes as well as methods for generating them.

#### Summary of the Invention

The objective of the present invention is to provide a method for the treatment of a disease by applying a medicament comprising a protease. Further the



present invention provides engineered proteins with novel functions that do not exist in the components used for the engineering of such proteins. In particular, the invention provides enzymes with user-definable specificities. User-definable specificity means that enzymes are provided with specificities that do not exist in the components used for the engineering of such enzymes. The specificities can be chosen by the user so that one or more intended target substrates are preferentially recognised and converted by the enzymes. Furthermore, the invention provides enzymes that possess essentially identical sequences to human proteins but have different specificities. In a particular embodiment, the invention provides proteases with user-definable specificities.

Furthermore, the present invention is directed to engineered enzymes which are fused to one or more further functional components. These further components can be proteinacious components which preferably have binding properties and are of the group consisting of substrate binding domains, antibodies, receptors or fragments thereof. Furthermore, these further components can be further functional components, preferably being selected from the group consisting of polyethylenglycols, carbohydrates, lipids, fatty acids, nucleic acids, metals, metal chelates, and fragments or derivatives thereof. The resulting fusion proteins are understood as enzymes with user-definable specificities within the present invention.

Besides, the invention is directed to the application of such enzymes with novel, user-definable specificities for therapeutic, research, diagnostic, nutritional, personal care or industrial purposes. Moreover, the invention is directed to a method for generating engineered enzymes with user-definable specificities. In particular, the invention is directed to generate enzymes that possess essentially identical sequences to human enzymes but have different specificities.

This problem has been solved by the embodiments of the invention specified in the description below and in the claims. The present invention is thus directed to (1) the use of a protease with defined specificity for a target substrate for preparing a medicament for the treatment of a specific disease related to said target substrate,

(2) an engineered enzyme with defined specificity characterized by the combination of the following components,:

(a) a protein scaffold which catalyzes at least one chemical reaction on at least one substrate, and

(b) one or more specificity determining regions (SDRs) located at sites in the protein scaffold that enable the resulting engineered protein to discriminate between at least one target substrate and one or more different substrates, and wherein the SDRs are essentially synthetic peptide sequences;

(3) the use of an engineered enzyme as defined in (2) above for therapeutic, research, diagnostic, nutritional, personal care or industrial purposes, preferably for the use as defined in (1) above;

(4) a method for generating engineered enzymes as defined in (2) above having specificities towards target substrates, such specificities not being present in the individual starting components, comprising at least the following steps:

(a) providing a protein scaffold which catalyzes at least one chemical reaction on at least one substrate,

(b) generating a library of engineered enzymes by combining the protein scaffold from step (a) with fully or partially random peptide sequences at sites in the protein scaffold that enable the resulting engineered enzyme to discriminate between at least one target substrate and one or more different substrates, and

(c) selecting out of the library of engineered enzymes generated in step (b) one or more enzymes that have specificities towards at least one target substrate;

(5) a fusion protein which is comprised of at least one engineered enzyme as defined in (2) above and at least one further component, preferably the at least one further component having binding properties and more preferably being selected from the group consisting of antibodies, binding domains, receptors, and fragments thereof;

(6) a composition or pharmaceutical composition comprising one or more engineered enzymes as defined in (2) above or a fusion protein as defined in (5) above, said pharmaceutical composition may optionally comprise an acceptable carrier, excipient and/or auxiliary agent;

(7) a DNA encoding the engineered enzyme as defined in (2) above;

(8) a vector comprising the DNA as defined in (7) above;

- (9) a host cell or transgenic organism being transformed/transfected with a vector as defined in (8) above and/or containing the DNA as defined in (7) above; and
- (10) a method for producing the engineered enzyme of (2) above comprising culturing a cell or organism as defined in (8) above and isolating the enzyme from the culture broth.

#### Brief description of the Figures

The following figures are provided in order to explain further the present invention in supplement to the detailed description:

Figure 1 illustrates the three-dimensional structure of human trypsin I with the active site residues shown in "ball-and-stick" representation and with the marked regions indicating potential SDR insertion sites.

Figure 2 shows the alignment of the primary amino acid sequence of three members of the serine protease class S1 family: human trypsin I, human alpha-thrombin and human enteropeptidase (see also SEQ ID NOs: 1, 5 and 6).

Figure 3 illustrates the three-dimensional structure of subtilisin with the active site residues being shown in "ball-and-stick" representation and with the numbered regions indicating potential SDR insertion sites.

Figure 4 shows the alignment of the primary amino acid sequences of four members of the serine protease class S8 family: subtilisin E, furin, PC1 and PC5 (see also SEQ ID NOs: 7-10).

Figure 5 illustrates the three-dimensional structure of pepsin with the active site residues being shown in "ball-and-stick" representation and with the numbered regions indicating potential SDR insertion sites.

Figure 6 shows the alignment of the primary amino acid sequences of three members of the A1 aspartic acid protease family: pepsin,  $\beta$ -secretase and cathepsin D (see also SEQ ID NOs: 11-13).

Figure 7: Illustrates the three-dimensional structure of caspase 7 with the active site residues being shown in “ball-and-stick” representation and with the numbered regions indicating potential SDR insertion sites.

Figure 8: shows the primary amino acid sequence of caspase 7 as a member of the cysteine protease class C14 family (see also SEQ ID NO: 14).

Figure 9 depicts schematically the third aspect of the invention.

Figure 10 shows a Western blot analysis of a culture supernatant of cells expressing variants of human trypsin I with SDR1 and SDR2, compared to negative controls.

Figure 11 shows the time course of the proteolytic cleavage of a target substrate by human trypsin I.

Figure 12 shows the relative activities of three variants of inventive engineered proteolytic enzymes in comparison with human trypsin I on two different peptide substrates.

Figure 13 shows the relative specificities of human trypsin I and variants of inventive engineered proteolytic enzymes with one or two SDRs, respectively.

Figure 14: shows the relative specificities of human trypsin I and of variants of inventive engineered proteolytic enzymes being specific for human TNF-alpha with this scaffold on peptides with a target sequence of human TNF-alpha.

Figure 15: shows the reduction of cytotoxicity induced by TNF-alpha when incubating the TNF-alpha with concentrated supernatant from cultures expressing the inventive engineered proteolytic enzymes being specific for human TNF-alpha.

Figure 16: shows the reduction of cytotoxicity induced by TNF-alpha when incubating the TNF-alpha with purified inventive engineered proteolytic enzyme being specific for human TNF-alpha.

Figure 17: compares the activity of inventive engineered proteolytic enzymes being specific for human TNF-alpha with the activity of human trypsin I on two protein substrates: (a) human TNF-alpha; (b) mixture of human serum proteins.

Figure 18: shows the specific activity of an inventive engineered proteolytic enzyme with specificity for human VEGF.

### Definitions

In the framework of the present invention the following terms and definitions are used.

The term "protease" means any protein molecule that is capable of hydrolysing peptide bonds. This includes naturally-occurring or artificial proteolytic enzymes, as well as variants thereof obtained by site-directed or random mutagenesis or any other protein engineering method, any active fragment of a proteolytic enzyme, or any molecular complex or fusion protein comprising one of the aforementioned proteins. A "chimera of proteases" means a fusion protein of two or more fragments derived from different parent proteases.

The term "substrate" means any molecule that can be converted catalytically by an enzyme. The term "peptide substrate" means any peptide, oligopeptide, or protein molecule of any amino acid composition, sequence or length, that contains a peptide bond that can be hydrolyzed catalytically by a protease. The peptide bond that is hydrolyzed is referred to as the "cleavage site". Numbering of positions in the substrate is done according to the system introduced by Schlechter & Berger (Biochem. Biophys. Res. Commun. 27 (1967) 157-162). Amino acid residues adjacent N-terminal to the cleavage site are numbered  $P_1$ ,  $P_2$ ,  $P_3$ , etc., whereas residues adjacent C-terminal to the cleavage site are numbered  $P_1'$ ,  $P_2'$ ,  $P_3'$ , etc.

The term "target substrate" describes a user-defined substrate which is specifically recognized and converted by an enzyme according to the invention. The term "target peptide substrate" describes a user-defined peptide substrate.

The term "target specificity" describes the qualitative and quantitative specificity of an enzyme that is capable of recognizing and converting a target substrate. Catalytic properties of enzymes are expressed using the kinetic parameters " $K_M$ " or "Michaelis Menten constant", " $k_{cat}$ " or "catalytic rate constant", and " $k_{cat} / K_M$ " or "catalytic efficiency", according to the definitions of Michaelis and Menten (Fersht, A., Enzyme Structure and Mechanism, W. H. Freeman and Company, New York, 1995). The term "catalytic activity" describes quantitatively the conversion of a given substrate under defined reaction conditions.

The term "specificity" means the ability of an enzyme to recognize and convert preferentially certain substrates. Specificity can be expressed qualitatively and quantitatively. "Qualitative specificity" refers to the chemical nature of the substrate residues that are recognized by an enzyme. "Quantitative specificity" refers to the number of substrates that are accepted as substrates. Quantitative specificity can be expressed by the term  $s$ , which is defined as the negative logarithm of the number of all accepted substrates divided by the number of all possible substrates. Proteases, for example, that accept preferentially a small portion of all possible peptide substrates have a "high specificity". Proteases that accept almost any peptide substrate have a "low specificity". Definitions are made in accordance to WO 03/095670 which is therefore incorporated by reference. Proteases with very low specificity are also referred to as "unspecific proteases". The term "defined specificity" refers to a certain type of specificity, i.e. to a certain target substrate or a set of certain target substrates that are preferentially converted versus other substrates.

The term "engineered" in combination with the term "enzyme" describes an enzyme that is comprised of different components and that has features not being conferred by the individual components alone.

The term "protein scaffold" or "scaffold protein" refers to a variety of primary, secondary and tertiary polypeptide structures.

The term "peptide sequence" indicates any peptide sequence used for insertion or substitution into or combination with a protein scaffold. Peptide sequences are usually obtained by expression from DNA sequences which can be synthesized

according to well-established techniques or can be obtained from natural sources. Insertion, substitution or combination of peptide sequences with the protein scaffold are generated by insertion, substitution or combination of oligonucleotides into or with a polynucleotide encoding the protein scaffold. The term "synthetic" in combination with the term "peptide sequence" refers to peptide sequences that are not present in the protein scaffold in which the peptide sequences are inserted or substituted or with which they are combined.

The term "components" in combination with the term "engineered enzyme" refers to peptide or polypeptide sequences that are combined in the engineering of such enzymes. Such components may among others comprise one or more protein scaffolds and one or more synthetic peptide sequences. The term "library of engineered enzymes" describes a mixture of engineered enzymes, whereby every single engineered enzyme is encoded by a different polynucleotide sequence. The term "gene library" indicates a library of polynucleotides that encodes the library of engineered enzymes. The term "SDR" or "Specificity determining region" refers to a synthetic peptide sequence that provides the defined specificity when combined with the protein scaffold at sites that enable the resulting enzymes to discriminate between the target substrate and one or more other substrates. Such sites are termed "SDR sites".

The terms "tertiary structure similar to the structure of" and "similar tertiary structure" in combination with the terms "enzyme" or "protein" refer to proteins in which the type, sequence, connectivity and relative orientation of the typical secondary structural elements of a protein, e.g. alpha-helices, beta-sheets, beta-turns and loops, are similar and the proteins are therefore grouped into the same structural or topological class or fold. This includes proteins that have altered, additional or deleted structural elements of any type but otherwise unchanged topology. Examples of such structural classes are the TNF superfamily, the S1 fold or the S8 fold within the serine proteases, the GPCRs, or the  $\alpha/\beta$ -barrel fold.

The term "positions that correspond structurally" indicates amino acids in proteins of similar tertiary structure that correspond structurally to each other, i.e. they are usually located within the same structural or topological element of the structure. Within the structural element they possess the same relative

positions with respect to beginning and end of the structural element. If, e.g. the topological comparison of two proteins reveals two structurally corresponding sequences of different length, then amino acids within, e.g. 20% and 40% of the respective region lengths, correspond to each other structurally.

The term "library of engineered enzymes" of the present invention refers to a multiplicity of enzymes or enzyme variants, which may exist as a mixture or in isolated form.

Amino acids residues are abbreviated according to the following Table 1 either in one- or in three-letter code.

Table 1: Amino acid abbreviations

Abbreviations		Amino acid
A	Ala	Alanine
C	Cys	Cysteine
D	Asp	Aspartic acid
E	Glu	Glutamic acid
F	Phe	Phenylalanine
G	Gly	Glycine
H	His	Histidine
I	Ile	Isoleucine
K	Lys	Lysine
L	Leu	Leucine
M	Met	Methionine
N	Asn	Asparagine
P	Pro	Proline
Q	Gln	Glutamine
R	Arg	Arginine
S	Ser	Serine
T	Thr	Threonine
V	Val	Valine
W	Trp	Tryptophane
Y	Tyr	Tyrosine



#### Detailed description of the invention

The present invention provides engineered proteins with novel functions. In particular, the invention provides enzymes with user-definable specificities. In a particular embodiment, the invention provides proteases with user-definable specificities. Besides, the invention provides applications of such enzymes with novel, user-definable specificities for therapeutic, research, diagnostic, nutritional, personal care or industrial purposes. Moreover, the invention provides a method for generating enzymes with specificities that are not present in the components used for the engineering of such enzymes. In particular, the invention is directed to the generation of enzymes that have sequences that are essentially identical to mammalian, especially human enzymes but have different specificities. Moreover, the invention provides libraries of specific engineered enzymes with corresponding specificities encoded genetically, a method for the generation of libraries of specific engineered enzymes with corresponding specificities encoded genetically, and the application of such libraries for technical, diagnostic, nutritional, personal care or research purposes.

A first aspect of the invention is directed to the application of engineered enzymes with specificities for therapeutic, research, diagnostic, nutritional, personal care or industrial purposes. The application comprises at least the following steps:

- (a) identification of a target peptide substrate whose hydrolysis has a positive effect in connection with the intended purpose, such as curing a disease, diagnosing a disease, processing of ingredients for human or animal nutrition, or other technical processes;
- (b) provision of an engineered enzyme, the enzyme being specific for the target peptide identified in step (a); and
- (c) use of the enzyme as provided in step (b) for the intended purpose.

In a first variant of this aspect of the invention, the engineered enzyme is used as a therapeutic means to inactivate a disease-related target substrate. This application comprises at least the following steps:

- (a) identification of a target substrate whose function is connected to a disease and whose inactivation has a positive effect in connection with the disease, and determination of a target site within the target substrate

- characterized by the fact that modification at the target site leads to the inactivation of the target substrate;
- (b) provision of an engineered enzyme, the enzyme being specific for the target site identified in step (a); and
  - (c) use of the enzyme for the inactivation of the target substrate inside or outside the human body.

Preferably, the scaffold is a protease and the modification is hydrolysis of a target site in a protein target. Preferably, the hydrolysis leads to the activation or inactivation of the peptide or protein target. Potential peptide or protein targets include soluble proteins, in particular cytokines, such as proteins of the TNF-superfamily, interleukines, interferons, chemokines and growth factors; hormones; toxins; enzymes, such as oxidoreductases, transferases, hydrolases, lyases, isomerases and ligases; structural proteins, such as collagen; and immunoglobulins; or membrane associated proteins, in particular single pass transmembrane proteins; multipass transmembrane proteins, such as G-protein coupled receptors, ion channels and transporters; lipid-anchored membrane proteins and GPI-anchored membrane proteins.

In a first embodiment of this variant the engineered enzyme is a protease and is capable of hydrolysing human tumor necrosis factor-alpha (hTNF- $\alpha$ ). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, rheumatoid arthritis, inflammatory bowel diseases, psoriasis, Crohn's disease, Ulcerative colitis, diabetes type II, classical Hodgkin's Lymphoma (cHL), Grave's disease, Hashimoto's thyroiditis, Sjogren's Syndrome, systemic lupus erythematosus, multiple sclerosis, Systemic inflammatory response syndrome (SIRS) which leads to distant organ damage and multiple organ dysfunction syndrome (MODS), eosinophilia, neurodegenerative disease, stroke, closed head injury, encephalitis, CNS disorders, asthma, rheumatoid arthritis, sepsis, vasodilation, intravascular coagulation and multiple organ failure, as well as other diseases connected with hTNF- $\alpha$ . Preferably, said enzyme or said fusion protein is capable of specifically inactivating hTNF- $\alpha$  (SEQ ID NO:96). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 31/32, 32/33, 44/45, 45/46, 87/88, 128/129, 130/131, 140/141 and/or 141/142 (most

preferred between positions 31/32, 32/33 and/or 45/46) in hTNF- $\alpha$ , or a peptide bond in proximity to these positions in hTNF- $\alpha$ , or peptide bonds in protein targets related to hTNF- $\alpha$  between positions having structural homology or sequence homology to these positions. In this embodiment it is most preferred that the protease has the a sequence shown in SEQ ID NO:74, SEQ ID NO:75 and is capable of hydrolysing hTNF- $\alpha$  at positions 31/32 and/or 32/33.

In a second embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Tumor necrosis factor ligand superfamily member 5 (hCD40-L). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, systemic lupus erythematosus and classical Hodgkin's Lymphoma (cHL), as well as other diseases connected with hCD40-L. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCD40-L (SEQ ID NO:143). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 133/134, 145/146, 165/166, 200/201, 201/202, 207/208 and/or 216/217 (most preferred between positions 133/134, 165/166, 201/202 and/or 216/217) in hCD40-L, or a peptide bond in proximity to these positions in hCD40-L, or peptide bonds in protein targets related to hCD40-L at positions having structural homology or sequence homology to these positions.

In a third embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Macrophage migration inhibitory factor (hMIF). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, inflammatory diseases, as well as other diseases connected with hMIF. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hMIF (SEQ ID NO:109). More preferably said engineered or said fusion protein is capable of hydrolysing the peptide bonds between positions 16/17, 44/45, 66/67, 73/74, 77/78, 88/89, 92/93 and/or 100/101 (most preferred between positions 16/17 and/or 92/93) in hMIF, or a peptide bond in proximity to these positions in hMIF, or peptide bonds in protein targets related to hMIF at positions having structural homology or sequence homology to these positions.

In a fourth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin-1 beta precursor (hIL-1 beta). The enzymes or

the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, diabetes, brain inflammation in cancer, arthritis, autoimmune and inflammatory diseases, as well as other diseases connected with hIL-1 beta. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-1 beta (SEQ ID NO:112). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 24/25, 35/36, 46/47, 54/55, 74/75, 75/76, 76/77, 77/78, 86/87, 88/89, 93/94, 94/95, 97/98 and/or 150/151 (most preferred between positions 35/36, 75/76, 76/77, 88/89, 93/94, 94/95 and/or 150/151) in hIL-1 beta, or a peptide bond in proximity to these positions in hIL-1 beta, or peptide bonds in protein targets related to hIL-1 beta at positions having structural homology or sequence homology to these positions.

In a fifth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 2 (hIL-2). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, T-cell leukemia and hairy cell leukemia, Crohn's disease, Ulcerative colitis, Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, systemic lupus erythematosus, multiple sclerosis, asthma and chronic obstructive pulmonary and classical Hodgkin's Lymphoma (cHL), as well as other diseases connected with hIL-2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-2 (SEQ ID NO:99). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 20/21, 32/33, 38/39, 43/44, 45/46, 48/49, 49/50, 54/55, 64/65, 76/77, 83/84, 84/85, 107/108, 109/110 and/or 120/121 (most preferred between positions 109/110) in hIL-2, or a peptide bond in proximity to these positions in hIL-2, or peptide bonds in protein targets related to hIL-2 at positions having structural homology or sequence homology to these positions.

In a sixth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 3 (hIL-3). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL) and eosinophilia, as well as other diseases connected with hIL-3. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-3 (SEQ ID NO:148). More preferably said enzyme or said fusion protein is capable of hydrolysing the

peptide bonds between positions 21/22, 28/29, 36/37, 44/45, 46/47, 51/52, 63/64, 66/67, 79/80, 94/95, 101/102, 108/109 and/or 109/110 (most preferred between positions 21/22, 28/29, 46/47, 63/64, 66/67, 79/80 and/or 101/102) in hIL-3, or a peptide bond in proximity to these positions in hIL-3, or peptide bonds in protein targets related to hIL-3 at positions having structural homology or sequence homology to these positions.

In a seventh embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 4 (hIL-4). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL), Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, Asthma, chronic obstructive pulmonary disease and allergic inflammatory reactions, as well as other diseases connected with hIL-4. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-4 (SEQ ID NO:118). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 4/5, 12/13, 31/32, 37/38, 61/62, 62/63, 64/65, 91/92, 102/103, 121/122 and/or 126/127 (most preferred between positions 4/5, 61/62, 62/63, 64/65 and/or 121/122) in hIL-4, or a peptide bond in proximity to these positions in hIL-4, or peptide bonds in protein targets related to hIL-4 at positions having structural homology or sequence homology to these positions.

In a eighth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin-5 (hIL-5). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL), asthma, chronic obstructive pulmonary disease, eosinophilia, allergic inflammatory diseases, as well as other diseases connected with hIL-5. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-5 (SEQ ID NO:133). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 12/13, 32/33, 67/68, 76/77, 77/78, 80/81, 83/84, 84/85, 85/86, 90/91, 91/92, 92/93 and/or 98/99 (most preferred between positions 90/91, 91/92, 92/93 and/or 98/99) in hIL-5, or a peptide bond in proximity to these positions in hIL-5, or peptide bonds in protein targets related to hIL-5 at positions having structural homology or sequence homology to these positions.

In a ninth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin-6 (hIL-6). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL), breast cancer, renal cell carcinoma, multiple myeloma, lymphoma, leukemia, Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, systemic lupus erythematosus, Systemic inflammatory response syndrome (SIRS) which leads to distant organ damage and multiple organ dysfunction syndrome (MODS), chronic obstructive pulmonary disease (COPD), Castleman's diseases, inflammatory bowel diseases, Crohn's disease, as well as other diseases connected with hIL-6. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-6 (SEQ ID NO:134). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 32/33, 35/36, 55/56, 71/72, 129/130, 130/131, 132/133, 135/136, 141/142, 161/162, 180/181 and/or 183/184 (most preferred between positions 135/136 and/or 141/142) in hIL-6, or a peptide bond in proximity to these positions in hIL-6, or peptide bonds in protein targets related to hIL-6 at positions having structural homology or sequence homology to these positions.

In a tenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 8 (hIL-8). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease, Ulcerative colitis, classical Hodgkin's Lymphoma (cHL), Systemic inflammatory response syndrome (SIRS) which leads to distant organ damage and multiple organ dysfunction syndrome (MODS), chronic obstructive pulmonary disease (COPD), endometriosis, psoriasis and atherosclerotic lesions, as well as other diseases connected with hIL-8. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-8 (SEQ ID NO:100). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 1/12, 15/16, 45/46, 47/48, 52/53, 54/55, 60/61, 64/65 and/or 67/68 (most preferred between positions 45/46) in hIL-8, or a peptide bond in proximity to these positions in hIL-8, or peptide bonds in protein targets related to hIL-8 at positions having structural homology or sequence homology to these positions.

In a eleventh embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin-10 (hIL-10). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL) and diseases related to the suppression of cytotoxic T-cells, as well as other diseases connected with hIL-10. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-10 (SEQ ID NO:135). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 24/25, 25/26, 27/28, 28/29, 40/41, 44/45, 49/50, 57/58, 59/60, 84/85, 86/87, 106/107, 107/108, 110/111, 130/131, 134/135, 137/138, 138/139 and/or 144/145 (most preferred between positions 24/25, 27/28, 44/45, 49/50, 86/87, 137/138 and/or 144/145) in hIL-10, or a peptide bond in proximity to these positions in hIL-10, or peptide bonds in protein targets related to hIL-10 at positions having structural homology or sequence homology to these positions.

In a twelfth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 12 beta chain (hIL-12 $\beta$ ). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease and classical Hodgkin's Lymphoma (cHL), as well as other diseases connected with hIL-12 $\beta$ . Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-12 $\beta$  (SEQ ID NO:97). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 14/15, 18/19, 29/30, 34/35, 87/88, 99/100, 102/103, 104/105, 161/162, 174/175, 222/223, 225/226, 228/229, 238/239, 268/269 and/or 293/294 (most preferred between positions 18/19, 34/35, 87/88 and/or 161/162) in hIL-12 $\beta$ , or a peptide bond in proximity to these positions in hIL-12 $\beta$ , or peptide bonds in protein targets related to hIL-12 $\beta$  at positions having structural homology or sequence homology to these positions.

In a thirteenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 13 (hIL-13). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of cancer, such as, but not limited to, classical Hodgkin's Lymphoma (cHL), eosinophilia, asthma, chronic obstructive pulmonary disease, fibrosis, psoriasis,

atopic dermatitis and Ulcerative colitis, as well as other diseases connected with hIL-13. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-13 (SEQ ID NO:119). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 25/26, 62/63, 65/66, 86/87, 87/88, 98/99, 108/109 and/or 111/112 (most preferred between positions 87/88) in hIL-13, or a peptide bond in proximity to these positions in hIL-13, or peptide bonds in protein targets related to hIL-13 at positions having structural homology or sequence homology to these positions.

In a fourteenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interleukin 18 (hIL-18). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease, inflammation liver injuries, pulmonary tuberculosis, plural tuberculosis and rheumatoid arthritis, as well as other diseases connected with hIL-18. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIL-18 (SEQ ID NO:98). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 17/18, 32/33, 37/38, 39/40, 40/41, 53/54, 58/59, 79/80, 90/91, 93/94, 98/99, 110/111, 120/121, 123/124, 131/132, 132/133, 142/143, 147/148 and/or 157/158 (most preferred between positions 37/38, 132/133, 142/143 and/or 157/158) in hIL-18, or a peptide bond in proximity to these positions in hIL-18, or peptide bonds in protein targets related to hIL-18 at positions having structural homology or sequence homology to these positions.

In a fifteenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interferon-gamma (hIFN-gamma). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL), Crohn's disease and type I diabetes, as well as other diseases connected with hIFN-gamma. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIFN-gamma (SEQ ID NO:137). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 2/3, 6/7, 13/14, 21/22, 24/25, 34/35, 36/37, 37/38, 62/63, 68/69, 83/84, 86/87, 90/91, 102/103, 107/108 and/or 108/109 (most preferred



between positions 13/14, 24/25, 37/38, 62/63, 68/69, 102/103 and/or 107/108) in hIFN-gamma, or a peptide bond in proximity to these positions in hIFN-gamma, or peptide bonds in protein targets related to hIFN-gamma at positions having structural homology or sequence homology to these positions.

In a sixteenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human small inducible cytokine A2 (hCCL2). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease and Ulcerative colitis, as well as other diseases connected with hCCL2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCCL2 (SEQ ID NO:102). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 3/4, 13/14, 18/19, 19/20, 24/25, 29/30, 38/39, 54/55, 56/57, 58/59, 62/63, 65/66 and/or 68/69 (most preferred between positions 19/20, 29/30, 38/39, 54/55 and/or 62/63) in hCCL2, or a peptide bond in proximity to these positions in hCCL2, or peptide bonds in protein targets related to hCCL2 at positions having structural homology or sequence homology to these positions.

In a seventeenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Eotaxin (hCCL11). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease and Ulcerative colitis, classical Hodgkin's Lymphoma (cHL), chronic pathophysiologic dysfunction, characterized by an influx mainly of Th2 cells, and eosinophilia, as well as other diseases connected with hCCL11. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCCL11 (SEQ ID NO:101). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 11/12, 16/17, 17/18, 22/23, 27/28, 33/34, 44/45, 47/48, 48/49, 52/53, 54/55, 56/57, 60/61, 66/67 and/or 73/74 (most preferred between positions 48/49 and/or 66/67) in hCCL11, or a peptide bond in proximity to these positions in hCCL11, or peptide bonds in protein targets related to hCCL11 at positions having structural homology or sequence homology to these positions.

In an eighteenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Vascular endothelial growth factor (hVEGF). The

enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, all solid tumors and metastatic solid tumors, inflammatory breast cancer, as well as other diseases connected with hVEGF. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hVEGF (SEQ ID NO:103). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 16/17, 19/20, 23/24, 34/35, 41/42, 56/57, 62/63, 63/64, 64/65, 65/66, 82/83, and/or 84/85 (most preferred between positions 23/24, 41/42, 63/64, 82/83 and/or 84/85) in hVEGF, or a peptide bond in proximity to these positions in hVEGF, or peptide bonds in protein targets related to hVEGF at positions having structural homology or sequence homology to these positions.

In an nineteenth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Transforming growth factor beta 1 (hTGF- $\beta$ 1). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, a variety of cancers, including breast cancer, colorectal cancer and classical Hodgkin's Lymphoma (cHL), fibrosis, suppression of cell-mediated immunity, glaucoma, diffuse systemic sclerosis as well as other diseases connected with hTGF- $\beta$ 1. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hTGF- $\beta$ 1 (SEQ ID NO:104). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 23/24, 25/26, 26/27, 27/28, 37/38, 55/56 and/or 94/95 (most preferred between positions 25/26, 55/56 and/or 94/95) in hTGF- $\beta$ 1, or a peptide bond in proximity to these positions in hTGF- $\beta$ 1, or peptide bonds in protein targets related to hTGF- $\beta$ 1 at positions having structural homology or sequence homology to these positions.

In a twentieth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Somatotropin (human Growth hormone; hGH). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, acromegaly, diabetes and diabetic kidney disease including renal hypertrophy and glomerular enlargement and cardiovascular disorders, as well as other diseases connected with hGH. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hGH (SEQ ID NO:121). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 8/9,

16/17, 19/20, 26/27, 33/34, 38/39, 41/42, 70/71, 77/78, 94/95, 103/104, 112/113, 115/116, 116/117, 130/131, 147/148, 154/155 and/or 178/179 (most preferred between positions 112/113, 147/148 and/or 154/155) in hGH, or a peptide bond in proximity to these positions in hGH, or peptide bonds in protein targets related to hGH at positions having structural homology or sequence homology to these positions.

In a twenty-first embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Insulin-like growth factor II (hIGF-II). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, diabetes and diabetic kidney disease, as well as other diseases connected with hIGF-II. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIGF-II (SEQ ID NO:122). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 15/16, 23/24, 24/25, 34/35, 37/38, 38/39, 48/49 and/or 49/50 (most preferred between positions 23/24) in hIGF-II, or a peptide bond in proximity to these positions in hIGF-II, or peptide bonds in protein targets related to hIGF-II at positions having structural homology or sequence homology to these positions.

In a twenty-second embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Hepatocyte growth factor (hHGF). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, angiogenic disorders and hepatocellular carcinoma, as well as other diseases connected with hHGF. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hHGF (SEQ ID NO:120). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 54/55, 60/61, 62/63, 63/64, 68/69, 76/77, 112/113, 123/124, 134/135, 168/169, 198/199 and/or 202/203 (most preferred between positions 63/64, 68/69, 76/77, 168/169 and/or 202/203) in hHGF, or a peptide bond in proximity to these positions in hHGF, or peptide bonds in protein targets related to hHGF at positions having structural homology or sequence homology to these positions.

In a twenty-third embodiment of this variant the enzyme is a protease and is capable of hydrolysing human hInsulin (hInsulin). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of

diseases, such as, but not limited to, insulin overdosage, as well as other diseases connected with hInsulin. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hInsulin B chain (SEQ ID NO:105) and/or hInsulin A chain (SEQ ID NO:106). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 16/17 and/or 22/23 in hInsulin B and/or between position 14/15 in Insulin A, or a peptide bond in proximity to these positions in hInsulin A or B, or peptide bonds in protein targets related to hInsulin A or B at positions having structural homology or sequence homology to these positions.

In a twenty-fourth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human hGhrelin (hGhrelin). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, obesity, as well as other diseases connected with hGhrelin. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hGhrelin (SEQ ID NO:107). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 1/2, 2/3, 3/4 and/or 4/5 in hGhrelin, or a peptide bond in proximity to these positions in hGhrelin, or peptide bonds in protein targets related to hGhrelin at positions having structural homology or sequence homology to these positions.

In a twenty-fifth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human angiotensinogen (angiotensin). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, essential hypertension, as well as other diseases connected with angiotensin. Preferably, said enzyme or said fusion protein is capable of specifically inactivating angiotensin (SEQ ID NO:108). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 1/2, 3/4 and/or 7/8 (most preferred between positions 3/4) in angiotensin, or a peptide bond in proximity to these positions in angiotensin, or peptide bonds in protein targets related to angiotensin at positions having structural homology or sequence homology to these positions.

In a twenty-sixth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human leptin precursor (leptin). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of

diseases, such as, but not limited to, obesity, as well as other diseases connected with leptin. Preferably, said enzyme or said fusion protein is capable of specifically inactivating leptin (SEQ ID NO:127). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 8/9, 9/10, 15/16, 23/24, 40/41, 53/54, 71/72, 85/86, 94/95, 108/109 and/or 141/142 (most preferred between positions 9/10, 40/41, 71/72, 94/95 and/or 108/109) in leptin, or a peptide bond in proximity to these positions in leptin, or peptide bonds in protein targets related to leptin at positions having structural homology or sequence homology to these positions.

In a twenty-seventh embodiment of this variant the enzyme is a protease and is capable of hydrolysing Protective antigen (PA-83). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, anthrax infection, as well as other diseases connected with PA-83. Preferably, said enzyme or said fusion protein is capable of specifically inactivating PA-83 (SEQ ID NO:123). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 72/73, 73/74, 92/93, 93/94, 131/132, 149/150, 178/179, 213/214, 214/215, 387/388, 425/426, 426/427, 427/428, 453/454, 520/521, 608/609, 617/618, 671/672, 679/680, 680/681, 683/684 and/or 684/685 (most preferred between positions 72/73, 73/74, 93/94, 149/150, 387/388, 425/426, 427/428 and/or 683/684) in PA-83, or a peptide bond in proximity to these positions in PA-83, or peptide bonds in protein targets related to PA-83 at positions having structural homology or sequence homology to these positions.

In a twenty-eighth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human plasminogen (plasminogen). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, thrombosis, as well as other diseases connected with plasminogen. Preferably, said enzyme or said fusion protein is capable of specifically inactivating plasminogen (SEQ ID NO:140). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bond between position 580/581 in plasminogen, or a peptide bond in proximity to this position in plasminogen, or peptide bonds in protein targets related to plasminogen at positions having structural homology or sequence homology to these positions.

In a twenty-ninth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Prothrombin (thrombin). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, bleeding, as well as other diseases connected with thrombin. Preferably, said enzyme or said fusion protein is capable of specifically inactivating thrombin (SEQ ID NO:149). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 198/199, 327/328, 363/364 (most preferred between positions 327/328 and/or 363/364) in thrombin, or a peptide bond in proximity to these positions in thrombin, or peptide bonds in protein targets related to thrombin at positions having structural homology or sequence homology to these positions

In a thirty embodiment of this variant the enzyme is a protease and is capable of hydrolysing human beta-secretase. The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Alzheimer, as well as other diseases connected with human beta-secretase precursor. Preferably, said enzyme or said fusion protein is capable of specifically inactivating human beta-secretase precursor (SEQ ID NO:139). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 61/62, 64/65, 159/160, 238/239, 239/240, 246/247, 256/257, 330/331 and/or 365/366 (most preferred between positions 61/62, 246/247 and/or 365/366) in human beta-secretase precursor, or a peptide bond in proximity to these positions in human beta-secretase precursor, or peptide bonds in protein targets related to human beta-secretase precursor at positions having structural homology or sequence homology to these positions.

In a thirty-first embodiment of this variant the enzyme is a protease and is capable of hydrolysing human matrix metalloproteinase-2 (hMMP-2). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, a variety of cancers including bladder cancer, breast tumor cancer, gastric cancer and lung cancer, as well as other diseases connected with hMMP-2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hMMP-2 (SEQ ID NO:131). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 62/63, 68/69, 75/76, 76/77, 79/80, 88/89,

110/111, 112/113, 115/116, 120/121, 164/165, 254/255, 267/268, 296/297, 324/325, 325/326, 382/383, 383/384, 470/471, 500/501, 550/551, 564/565, 595/596, 597/598, 608/609, 646/647, 649/650 and/or 650/651 (most preferred between positions 68/69, 115/116, 120/121, 164/165, 325/326, 383/384, 470/471, 500/501, 595/596, 608/609 and/or 650/651) in hMMP-2, or a peptide bond in proximity to these positions in hMMP-2, or peptide bonds in protein targets related to hMMP-2 at positions having structural homology or sequence homology to these positions.

In a thirty-second embodiment of this variant the enzyme is a protease and is capable of hydrolysing human matrix metalloproteinase-9 (hMMP-9). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, a variety of cancers including bladder cancer, breast tumor cancer, gastric cancer and lung cancer, as well as other diseases connected with hMMP-9. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hMMP-9 (SEQ ID NO:132). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 41/42, 42/43, 106/107, 113/114, 134/135, 160/161, 162/163, 163/164, 222/223, 226/227, 265/266, 266/267, 267/268, 284/285, 309/310, 321/322, 322/323, 324/325, 356/357, 380/381, 433/434 and/or 440/441 (most preferred between positions 160/161, 163/164, 226/227, 284/285, 321/322, 322/323 and/or 433/434) in hMMP-9, or a peptide bond in proximity to these positions in hMMP-9, or peptide bonds in protein targets related to hMMP-9 at positions having structural homology or sequence homology to these positions.

In a thirty-third embodiment of this variant the enzyme is a protease and is capable of hydrolysing HIV membrane glycoprotein (GP120). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, AIDS or HIV infection, as well as other diseases connected with GP120 or HIV infection. Preferably, said enzyme or said fusion protein is capable of specifically inactivating GP120 (SEQ ID NO:124). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 97/98, 99/100, 107/108, 113/114, 117/118, 227/228, 231/233, 279/280, 335/336, 337/338, 368/369, 412/413, 419/420, 429/430, 444/445, 457/458, 474/475, 476/477, 477/478, 485/486 and/or

490/491 (most preferred between positions 99/100, 368/369, 412/413, 419/420, 444/445 and/or 490/491) in GP120, or a peptide bond in proximity to these positions in GP120, or peptide bonds in protein targets related to GP120 at positions having structural homology or sequence homology to these positions.

In a thirty-fourth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Cytotoxic T-lymphocyte protein 4 (hCTLA-4). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, breast cancer, as well as other diseases connected with hCTLA-4. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCTLA-4 (SEQ ID NO:144). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 14/15, 28/29, 33/34, 38/39, 41/42, 62/63, 72/73, 85/86, 95/96, 100/101, 105/106, 119/120, 125/126 and/or 127/128 (most preferred between positions 14/15, 28/29, 38/39, 41/42, 62/63 and/or 85/86) in hCTLA-4, or a peptide bond in proximity to these positions in hCTLA-4, or peptide bonds in protein targets related to hCTLA-4 at positions having structural homology or sequence homology to these positions.

In a thirty-fifth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Integrin alpha-2 (hVLA-2). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, renal tumors, uveal melanomas and gastrointestinal tumors, as well as other diseases connected with hVLA-2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hVLA-2 (SEQ ID NO:147). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 160/161, 174/175, 201/202, 219/220, 231/232, 232/233, 233/234, 243/244, 259/260, 264/265, 268/269, 288/289, 292/293, 294/295, 298/299, 301/302, 310/311 and/or 317/318 (most preferred between positions 160/161, 174/175, 201/202, 219/220, 243/244, 264/265, 292/293 and/or 294/295) in hVLA-2, or a peptide bond in proximity to these positions in hVLA-2, or peptide bonds in protein targets related to hVLA-2 at positions having structural homology or sequence homology to these positions.

In a thirty-sixth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Vascular endothelial growth factor receptor 1



(hVEGFR 1). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, solid tumors and metastatic solid tumors, astrocytic brain tumors, pancreatic cancer, metastatic renal cancer, metastatic solid tumors, as well as other diseases connected with hVEGFR 1. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hVEGFR 1 (SEQ ID NO:114). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 189/190, 190/191, 224/225 and/or 331/332 (most preferred between positions 189/190 and/or 331/332) in hVEGFR 1, or a peptide bond in proximity to these positions in hVEGFR 1, or peptide bonds in protein targets related to hVEGFR 1 at positions having structural homology or sequence homology to these positions.

In a thirty-seventh embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Vascular endothelial growth factor receptor 2 (hVEGFR 2). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, solid tumors and metastatic solid tumors, pancreatic cancer, metastatic renal cancer, metastatic CRC, as well as other diseases connected with hVEGFR 2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hVEGFR 2 (SEQ ID NO:115). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 214/215, and/or 323/324 (most preferred between position 214/215) in hVEGFR 2, or a peptide bond in proximity to these positions in hVEGFR 2, or peptide bonds in protein targets related to hVEGFR 2 at positions having structural homology or sequence homology to these positions.

In a thirty-eighth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Epidermal growth factor receptor (hEGFr). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, bladder cancer, breast cancer, cervical cancer, colorectal cancer, endometrial cancer, oesophageal cancer, head and neck cancer, gastric cancer, non-small-cell lung carcinoma and ovarian cancer, as well as other diseases connected with hEGFr. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hEGFr (SEQ ID NO:116). More preferably said enzyme or said fusion protein is capable of

hydrolysing the peptide bonds between positions 20/21, 29/30, 48/49, 74/75, 165/166, 202/203, 220/221, 246/247, 251/252, 269/270, 270/271, 304/305, 305/306, 357/358, 430/431, 443/444, 454/455, 455/456, 463/464, 465/466, 476/477, 507/508 and/or 509/510 in hEGFr, or a peptide bond in proximity to these positions in hEGFr, or peptide bonds in protein targets related to hEGFr at positions having structural homology or sequence homology to these positions.

In a thirty-ninth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Epithelial cell adhesion molecule (hEp-CAM). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, colorectal cancer, as well as other diseases connected with hEp-CAM. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hEp-CAM (SEQ ID NO:125). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 14/15, 19/20, 25/26, 30/31, 33/34, 55/56 and/or 70/71 (most preferred between positions 14/15, 30/31 and/or 70/71) in hEp-CAM, or a peptide bond in proximity to these positions in hEp-CAM, or peptide bonds in protein targets related to hEp-CAM at positions having structural homology or sequence homology to these positions.

In a forty embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Insulin-like growth factor I receptor (hIGF-1r). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, a variety of cancers including breast cancer, as well as other diseases connected with hIGF-1r. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIGF-1r (SEQ ID NO:126). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 59/60, 115/116, 146/147, 171/172, 191/192, 290/291, 306/307, 307/308, 335/336, 336/337, 455/456 and/or 470/471 (most preferred between positions 306/307, 307/308, 335/336 and/or 470/471) in hIGF-1r, or a peptide bond in proximity to these positions in hIGF-1r, or peptide bonds in protein targets related to hIGF-1r at positions having structural homology or sequence homology to these positions.

In a forty-first embodiment of this variant the enzyme is a protease and is capable of hydrolysing human T-cell surface antigen CD2 precursor (hCD2). The enzymes or the fusion protein can thus be used for preparing medicaments for

the treatment of diseases, such as, but not limited to, psoriasis, as well as other diseases connected with hCD2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCD2 (SEQ ID NO:128). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 42/43, 43/44, 48/49, 49/50, 51/52, 54/55, 63/64, 69/70, 89/90 and/or 91/92 (most preferred between positions 43/44, 51/52 and/or 89/90) in hCD2, or a peptide bond in proximity to these positions in hCD2, or peptide bonds in protein targets related to hCD2 at positions having structural homology or sequence homology to these positions.

In a forty-second embodiment of this variant the enzyme is a protease and is capable of hydrolysing human T-cell surface glycoprotein CD4 (hCD4). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, psoriasis, transplant rejection, graft-versus-host colitis, autoimmune disorders and rheumatoid arthritis, as well as other diseases connected with hCD4. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCD4 (SEQ ID NO:129). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 166/167, 167/168, 206/207, 219/220, 224/225, 226/227, 251/252, 252/253, 322/323, 329/330 and/or 334/335 (most preferred between positions 206/207, 219/220, 251/252 and/or 252/253) in hCD4, or a peptide bond in proximity to these positions in hCD4, or peptide bonds in protein targets related to hCD4 at positions having structural homology or sequence homology to these positions.

In a forty-third embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Integrin alpha-L (hCD11a). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, psoriasis as well as other diseases connected with hCD11a. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCD11a (SEQ ID NO:130). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 145/146, 152/153, 156/157, 159/160, 160/161, 177/178, 178/179, 189/190, 190/191, 191/192, 193/194, 197/198, 200/201, 221/222, 229/230, 249/250, 253/254, 268/269, 290/291, 297/298, 304/305 and/or 305/306 (most preferred between positions 145/146, 159/160, 160/161,

189/190, 229/230, 249/250, 268/269, 297/298, 304/305 and/or 305/306) in hCD11a, or a peptide bond in proximity to these positions in hCD11a, or peptide bonds in protein targets related to hCD11a at positions having structural homology or sequence homology to these positions.

In a forty-fourth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Interferon-gamma receptor alpha chain (hIFN-gamma-R1). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, classical Hodgkin's Lymphoma (cHL) and type I diabetes, as well as other diseases connected with hIFN-gamma-R1. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hIFN-gamma-R1 (SEQ ID NO:136). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 49/50, 52/53, 62/63, 106/107, 122/123, 174/175, 215/216 and/or 222/223 (most preferred between positions 49/50, 122/123, 174/175 and/or 215/216) in hIFN-gamma-R1, or a peptide bond in proximity to these positions in hIFN-gamma-R1, or peptide bonds in protein targets related to hIFN-gamma-R1 at positions having structural homology or sequence homology to these positions.

In a forty-fifth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Platelet membrane glycoprotein IIb/IIIa (hGPIIb/IIIa). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, unstable angina, carotid stenting, ischemic stroke, peripheral vascular diseases, angiogenesis-related diseases and disseminating tumors, as well as other diseases connected with hGPIIb/IIIa. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hGPIIb/IIIa (SEQ ID NO:141). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 67/68, 91/92, 129/130, 143/144, 144/145, 181/182, 208/209, 209/210, 216/217, 239/240, 261/262, 410/411, 532/533, 556/557, 557/558, 597/598, 650/651 and/or 689/690 (most preferred between positions 67/68, 261/262, 410/411, 650/651 and/or 689/690) in hGPIIb/IIIa, or a peptide bond in proximity to these positions in hGPIIb/IIIa, or peptide bonds in protein targets related to hGPIIb/IIIa at positions having structural homology or sequence homology to these positions.

In a forty-sixth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Intercellular adhesion molecule-1 (hICAM-1). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease, as well as other diseases connected with hICAM-1. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hICAM-1 (SEQ ID NO:142). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 40/41, 88/89, 97/98, 102/103, 128/129, 131/132, 132/133, 149/150, 150/151, 160/161 and/or 166/167 (most preferred between positions 88/89, 102/103, 150/151, 160/161 and/or 166/167) in hICAM-1, or a peptide bond in proximity to these positions in hICAM-1, or peptide bonds in protein targets related to hICAM-1 at positions having structural homology or sequence homology to these positions.

In a forty-seventh embodiment of this variant the enzyme is a protease and is capable of hydrolysing human TGF-beta receptor type II (hTGF-beta RII). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, diffuse systemic sclerosis, as well as other diseases connected with hTGF-beta RII. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hTGF-beta RII (SEQ ID NO:145). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 32/33, 34/35, 35/36, 66/67, 67/68, 69/70, 82/83, 103/104, 104/105, 105/106, 118/119, 122/123 and/or 130/131 (most preferred between positions 32/33, 34/35, 66/67, 69/70, 104/105, 122/123 and/or 130/131) in hTGF-beta RII, or a peptide bond in proximity to these positions in hTGF-beta RII, or peptide bonds in protein targets related to hTGF-beta RII at positions having structural homology or sequence homology to these positions.

In a forty-eighth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Membrane cofactor protein (hMCP). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, renal tumors, uveal melanomas and gastrointestinal tumors, as well as other diseases connected with hMCP. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hMCP (SEQ ID NO:146). More preferably said enzyme or said fusion

protein is capable of hydrolysing the peptide bonds between positions 15/16, 17/18, 25/26, 31/32, 32/33, 35/36, 48/49, 67/68, 69/70, 110/111, 119/120 and/or 125/126 (most preferred between positions 15/16, 32/33, 48/49, 119/120 and/or 125/126) 130/131) in hMCP, or a peptide bond in proximity to these positions in hMCP, or peptide bonds in protein targets related to hMCP at positions having structural homology or sequence homology to these positions.

In a forty-ninth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Protease activated receptor 1 (hPAR1). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, thrombosis, as well as other diseases connected with hPAR1. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hPAR1 (SEQ ID NO:110). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 46/47, 51/52 and/or 52/53 in PAR1, or a peptide bond in proximity to these positions in hPAR1, or peptide bonds in protein targets related to hPAR1 at positions having structural homology or sequence homology to these positions.

In a fifth embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Protease activated receptor 2 (hPAR2). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, Crohn's disease, Ulcerative colitis and Inflammatory bowel disease, asthma, inflammation associated pain and arthritis, as well as other diseases connected with hPAR2. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hPAR2 (SEQ ID NO:111). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 41/42, 51/52 and/or 59/60 in hPAR2, or a peptide bond in proximity to these positions in hPAR2, or peptide bonds in protein targets related to hPAR2 at positions having structural homology or sequence homology to these positions.

In a fifty-first embodiment of this variant the enzyme is a protease and is capable of hydrolysing human Protease activated receptor 4 (hPAR4). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, thrombosis, as well as other diseases connected with hPAR4. Preferably, said enzyme or said fusion

protein is capable of specifically inactivating hPAR4 (SEQ ID NO:113). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 68/69, 74/75 and/or 78/79 in hPAR4, or a peptide bond in proximity to these positions in hPAR4, or peptide bonds in protein targets related to hPAR4 at positions having structural homology or sequence homology to these positions.

In a fifty-second embodiment of this variant the enzyme is a protease and is capable of hydrolysing human 5-hydroxytryptamine 1A receptor (h5-HT-1A). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, irritable bowel syndrome, as well as other diseases connected with h5-HT-1A. Preferably, said enzyme or said fusion protein is capable of specifically inactivating h5-HT-1A (SEQ ID NO:117). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 101/102, 102/103, 181/182 and/or 370/371 in h5-HT-1A a peptide bond in proximity to these positions in h5-HT-1A, or peptide bonds in protein targets related to h5-HT-1A at positions having structural homology or sequence homology to these positions.

In a fifty-third embodiment of this variant the enzyme is a protease and is capable of hydrolysing human carcinoembryonic antigen (hCEA). The enzymes or the fusion protein can thus be used for preparing medicaments for the treatment of diseases, such as, but not limited to, colon cancer, as well as other diseases connected with hCEA. Preferably, said enzyme or said fusion protein is capable of specifically inactivating hCEA (SEQ ID NO:138). More preferably said enzyme or said fusion protein is capable of hydrolysing the peptide bonds between positions 17/18, 69/70, 71/72, 74/75, 77/78, 98/99, 116/117, 126/127 and/or 128/129 in hCEA, or a peptide bond in proximity to these positions in hCEA, or peptide bonds in protein targets related to hCEA at positions having structural homology or sequence homology to these positions.

It is obvious to someone skilled in the art that also polymorphisms of all target sequences referred to are included. The expression "proximity to these positions" in all embodiments above refer to positions of peptide bonds that are between 10 and 5 Ångström and/or 5 amino acids, preferably 3 amino acids, next to the positions of the peptide bonds

Preferably, in this variant the scaffold of the engineered enzyme provided in step (c) is of human origin in order to avoid or reduce immunogenicity or allergenic effects associated with the application of the enzyme in the human body.

Alternatively, immunoenicity and allergenicity can be reduced by deimmunization of the engineered enzyme.

In further embodiment of this variant, the target substrate is a pro-drug which is activated by the engineered enzyme. In a particular embodiment of this variant, the engineered enzyme has proteolytic activity and the target substrate is a protein target which is proteolytically activated. Examples of such pro-drugs are pro-proteins such as the inactivated forms of coagulations factors. In another particular variant, the engineered enzyme is an oxidoreductase and the target substrate is a chemical that can be activated by oxidation.

In a second variant of this aspect of the invention, the engineered enzyme is used for diagnostic puposes. In a particular embodiment of this variant, the engineered enzyme is target-specific protease. Such diagnostic purposes comprise but are not limited to applications with the aim of diagnosing diseases, testing genetic predispositions or monitoring disease progression during therapy. In a particular embodiment, the diagnosis is based on the testing for the presence or absence of a disease-specific marker protein or a disease-specific variant of a human protein in test samples such as human tissue samples, blood samples or other samples taken from patients. The testing employs a protease with specificity for a particular, disease-related target protein. The testing is done by analysing the proteolytic degradation of such protein in the test sample. In a preferred embodiment the aim of the diagnostic test is to detect and/or quantify a disease-specific variant of a native human protein. Such a diagnostic test employs a protease that is specific for the disease-related protein variant, i.e. it has significantly higher proteolytic activity on the disease-related protein variant compared to the native human protein. The disease-related protein variant is therefore detected and/or quantified by detecting and/or quantifying the activity of the target-specific protease. Such detection and/or quantification is done by directly measuring the degradation products of the target protein or indirectly by measuring the influence of the target protein on the activity of the



target-specific protease by a competition assay. In another preferred embodiment the aim of the diagnostic test is to detect and/or quantify a protein that is specific for an infection by an infectious agent such as a virus or a bacterium. Such a diagnostic test employs a protease that is specific for a protein specifically expressed upon infection by the infectious agent, i.e. it has significantly higher proteolytic activity on a particular infection-indicating protein compared to any other native human protein. The infection-indicating protein is therefore detected and/or quantified by detecting and/or quantifying the proteolytic activity of the target-specific protease. Such detection and/or quantification is done by directly measuring the degradation products of the infection-indicating protein or indirectly by measuring the influence of the infection-indicating protein on the activity of the target-specific protease by a competition assay.

In a third variant of this aspect of the invention, the engineered enzyme is used as a technical means in order to catalyze an industrially or nutritionally relevant reaction with defined specificity. In a particular embodiment of this variant the engineered enzyme has proteolytic activity, the catalyzed reaction is a proteolytic processing, and the engineered enzyme specifically hydrolyses one or more industrially or nutritionally relevant protein substrates. In a preferred embodiment of this variant the engineered enzyme hydrolyses one or more industrially or nutritionally relevant protein substrates at specific sites, thereby leading to industrially or nutritionally desired product properties such as texture, taste or precipitation characteristics. In a further particular embodiment of this variant, the engineered enzyme catalyzes the hydrolysis of glycosidic bonds (glycosidase or glycosylases activity). Then, preferably, the catalyzed reaction is a polysaccharide processing, and the engineered enzyme specifically hydrolyses one or more industrially, technically or nutritionally relevant polysaccharide substrates. In a further particular embodiment of this variant, the engineered enzyme catalyzes the hydrolysis of triglyceride esters or lipids (lipase activity). Then, preferably, the catalyzed reaction is a lipid processing step, and the engineered enzyme specifically hydrolyses one or more industrially, technically or nutritionally relevant lipid substrates. In a further particular variant of this embodiment, the engineered enzyme catalyzes the oxidation or reduction of substrates (oxidoreductase activity). Then, preferably, the engineered enzyme

specifically oxidizes or reduces one or more industrially, technically or nutritionally relevant chemical substrates.

A second aspect of the invention discloses engineered enzymes with defined specificities. These engineered enzymes are characterized by the following components:

- (a) a protein scaffold capable of catalyzing at least one chemical reaction on a substrate, and
- (b) one or more specificity determining regions (SDRs) located at sites in the protein scaffold that enable the resulting engineered protein to discriminate between at least one target substrate and one or more different substrates, wherein the SDRs are essentially synthetic peptide sequences.

Preferably, such defined specificity of the engineered enzymes is not conferred by the protein scaffold.

In principle, the protein scaffold can have a variety of primary, secondary and tertiary structures. The primary structure, i.e. the amino acid sequence, can be an engineered sequence or can be derived from any viral, prokaryotic or eukaryotic origin. For human therapeutic use, however, the protein scaffold is preferably of mammalian origin, and more preferably, of human origin. Furthermore, the protein scaffold is capable to catalyze one or more chemical reactions and has preferably only a low specificity.

Preferably, derivatives of the protein scaffold are used that have modified amino acid sequences that confer improved characteristics for the applicability as protein scaffolds. Such improved characteristics comprise, but are not limited to, stability; expression or secretion yield; folding, in particular after combination of the protein scaffold with SDRs; increased or decreased sensitivity to regulators such as activators or inhibitors; immunogenicity; catalytic rate;  $K_M$  or substrate affinity.

The engineered enzymes reveal their quantitative specificity from the synthetic peptide sequences that are combined with the protein scaffold. Therefore, the engineered peptide sequences are acting as Specificity Determining Regions or

SDRs. The number, the length and the positions of such SDRs can vary over a wide range. The number of SDRs within the scaffold is at least one, preferably more than one, more preferably between two and eleven, most preferably between two and six. The SDRs have a length between one and 50 amino acid residues, preferably a length between one and 15 amino acid residues, more preferably a length between one and six amino acid residues. Alternatively, the SDRs have a length between two and 20 amino acid residues, preferably a length between two and ten amino acid residues, more preferably a length between three and eight amino acid residues.

The inventive engineered enzymes can further be described as antibody-like protein molecules comprising constant and variable regions, but having a non-immunoglobulin backbone and having an active site (catalytic activity) in the constant region, whereby the substrate specificity of the active site is modulated by the variable region. Preferably, as in the immunoglobulin structure, the variable regions are loops of variable length and composition that interact with a target molecule.

In a particular variant of the invention, the engineered enzymes have hydrolase activity. In a preferred variant, the engineered enzymes have proteolytic activity. Particularly preferred protein scaffolds for this variant are unspecific proteases or are parts from unspecific proteases or are otherwise derived from unspecific proteases. The expressions "derived from" or "a derivative thereof" in this respect and in the following variants and embodiments refer to derivatives of proteins that are mutated at one or more amino acid positions and/or have a homology of at least 70%, preferably 90%, more preferably 95% and most preferably 99% to the original protein, and/or that are proteolytically processed, and/or that have an altered glycosylation pattern, and/or that are covalently linked to non-protein substances, and/or that are fused with further protein domains, and/or that have C-terminal and/or N-terminal truncations, and/or that have specific insertions, substitutions and/or deletions. Alternatively, "derived from" may refer to derivatives that are combinations or chimeras of two or more fragments from two or more proteins, each of which optionally comprises any or all of the aforementioned modifications. The tertiary structure of the protein scaffold can be of any type. Preferably, however, the tertiary structure belongs to

one of the following structural classes: class S1 (chymotrypsin fold of the serine proteases family), class S8 (subtilisin fold of the serine proteases family), class SC (carboxypeptidase fold of the serine proteases family), class A1 (pepsin A fold of the aspartic proteases), or class C14 (caspase-1 fold of the cysteine proteases). Examples of proteases that can serve as the protein scaffold of engineered proteolytic enzymes for the use as human therapeutics are or are derived from human trypsin, human thrombin, human chymotrypsin, human pepsin, human endothiapepsin, human caspases 1 to 14, and/or human furin.

The defined specificity of the engineered proteolytic enzymes is a measure of their ability to discriminate between at least one target peptide or protein substrates and one or more further peptide or protein substrates. Preferably, the defined specificity refers to the ability to discriminate peptide or protein substrates that differ in other positions than the P1 site, more preferably, the defined specificity refers to the ability to discriminate peptide or protein substrates that differ in other positions than the P1 site and the P1' site. Most preferably, the engineered proteolytic enzymes distinguish target peptide or protein substrates at as many sites as is necessary to preferentially hydrolyse the target substrate versus other proteins. As an example, a therapeutically useful engineered proteolytic enzyme applied intravenously in the human body should be sufficiently specific to discriminate between the target substrate and any other protein in the human serum. Preferably, such an engineered proteolytic enzyme recognizes and discriminates peptide substrates at three or more amino acid positions, more preferably at four or more positions, and even more preferably at five or more amino acid positions. These positions may either be adjacent or non-adjacent.

In a first embodiment, the protein scaffold has a tertiary structure or fold equal or similar to the tertiary structure or fold of the S1 structural subclass of serine proteases, i. e. the chymotrypsin fold, and/or has at least 70% identity on the amino acid level to a protein of the S1 structural subclass of serine proteases. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 18-25, 38-48, 54-63, 73-86, 122-130, 148-156, 165-171 and 194-204 in human trypsin I, and more preferably at one

or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 20-23, 41-45, 57-60, 76-83, 125-128, 150-153, 167-169 and 197-201 (numbering of amino acids according to SEQ ID NO:1). The number of SDRs to be combined with this type of protein scaffold is preferably between 1 and 10, and more preferably between 2 and 4. Preferably, the protein scaffold is equal to or is a derivative or homologue of one or more of the following proteins: chymotrypsin, granzyme, kallikrein, trypsin, mesotrypsin, neutrophil elastase, pancreatic elastase, enteropeptidase, cathepsin, thrombin, ancrod, coagulation factor IXa, coagulation factor VIIa, coagulation factor Xa, activated protein C, urokinase, tissue-type plasminogen activator, plasmin, Desmodus-type plasminogen activator. More preferably, the protein scaffold is trypsin or thrombin or is a derivative or homologue from trypsin or thrombin. For the use as a human therapeutic, the trypsin or thrombin scaffold is most preferably of human origin in order to minimize the risk of an immune response or an allergenic reaction.

Preferably, derivatives with improved characteristics derived from human trypsin I or from proteins with similar tertiary structure are used. Preferred examples of such derivatives are derived from human trypsin I (SEQ ID NO:1) and comprise one or more of the following amino acid substitutions E56G; R78W; Y131F; A146T; C183R.

It is preferred that at least one of two SDRs are inserted into human trypsin I, or a derivative thereof, between residues 42 and 43 (SDR 1) and between 123 and 124 (SDR 2), respectively (numbering of amino acids according to SEQ ID NO:1). In addition the SDR 1 has a preferred length of 6 and the SDR 2 has a preferred length of 5 amino acids, respectively. In a preferred variant of this embodiment, the SDR 1 and SDR 2 sequences comprise one of the amino acid sequences listed in table 2. Such engineered proteolytic enzymes have specificity for the target substrate B as exemplified in example IV.

In a further embodiment the protein scaffold belongs to the S8 structural subclass of serine proteases and/or has a tertiary structure similar to subtilisin E from *Bacillus subtilis* and/or has at least 70% identity on the amino acid level to a protein of the S8 structural subclass of serine proteases. Preferably, the scaffold belongs to the subtilisin family or the human pro-protein convertases. It

is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 6-17, 25-29, 47-55, 59-69, 101-111, 117-125, 129-137, 139-154, 158-169, 185-195 and 204-225 in subtilisin E from *Bacillus subtilis*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 59-69, 101-111, 129-137, 158-169 and 204-225 (numbering of amino acids according to SEQ ID NO:7). It is preferred that the protein scaffold is equal to or is a derivative or homologue of one or more of the following proteins: subtilisin Carlsberg; *B. subtilis* subtilisin E; subtilisin BPN'; *B. licheniformis* subtilisin; *B. lentus* subtilisin; *Bacillus alcalophilus* alkaline protease; proteinase K; kexin; human pro-protein convertase; human furin. In a preferred variant, subtilisin BPN' or one of the proteins SPC 1 to 7 is used as the protein scaffold.

In a further embodiment the protein scaffold belongs to the family of aspartic proteases and/or has a tertiary structure similar to human pepsin. Preferably, the scaffold belongs to the A1 class of proteases and/or has at least 70% identity on the amino acid level to a protein of the A1 class of proteases. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 6-18, 49-55, 74-83, 91-97, 112-120, 126-137, 159-164, 184-194, 242-247, 262-267 and 277-300 in human pepsin, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 10-15, 75-80, 114-118, 130-134, 186-191 and 280-296 (numbering of amino acids according to SEQ ID NO:11). It is preferred that the protein scaffold is equal to or is a derivative or homologue of one or more of the following proteins: pepsin, chymosin, renin, cathepsin, yapsin. Preferably, pepsin or endothiopepsin or a derivative or homologue thereof is used as the protein scaffold.

In a further embodiment the protein scaffold belongs to the cysteine protease family and/or has a tertiary structure similar to human caspase 7. Preferably the scaffold belongs to the C14 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C14 class of cysteine proteases. It is preferred that SDRs are inserted into the protein scaffold at one

or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 78-91, 144-160, 186-198, 226-243 and 271-291 in human caspase 7, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 80-86, 149-157, 190-194 and 233-238 (numbering of amino acids according to SEQ ID NO:14). It is preferred that the protein scaffold is equal to or is a derivative or homologue of one of the caspases 1 to 9.

In a further embodiment the protein scaffold belongs to the S11 class of serine proteases or has at least 70% identity on the amino acid level to a protein of the S11 class of serine proteases and/or has a tertiary structure similar to D-alanyl-D-alanine transpeptidase from *Streptomyces* species K15. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 67-79, 137-150, 191-206, 212-222 and 241-251 in D-alanyl-D-alanine transpeptidase from *Streptomyces* species K15, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 70-75, 141-147, 195-202 and 216-220 (numbering of amino acids according to SEQ ID NO:15). It is preferred that the D-alanyl-D-alanine transpeptidase from *Streptomyces* species K15 or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the S21 class of serine proteases or has at least 70% identity on the amino acid level to a protein of the S21 class of serine proteases and/or has a tertiary structure similar to assemblin from human cytomegalovirus. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 25-33, 64-69, 134-155, 162-169 and 217-244 in assemblin from human cytomegalovirus, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 27-31, 164-168 and 222-239 (numbering of amino acids according to SEQ ID NO:16). It is preferred that the assemblin from human cytomegalovirus or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the S26 class of serine proteases or has at least 70% identity on the amino acid level to a protein of the S26 class of serine proteases and/or has a tertiary structure similar to the signal peptidase from *Escherichia coli*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 8-14, 57-68, 125-134, 239-254, 200-211 and 228-239 in signal peptidase from *Escherichia coli*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 9-13, 60-67, 127-132 and 203-209 (numbering of amino acids according to SEQ ID NO:17). It is preferred that the signal peptidase from *Escherichia coli* or a derivative or homologue thereof is used as the scaffold.

In an further embodiment the protein scaffold belongs to the S33 class of serine proteases or has at least 70% identity on the amino acid level to a protein of the S33 class of serine proteases and/or has a tertiary structure similar to the prolyl aminopeptidase from *Serratia marcescens*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 47-54, 152-160, 203-212 and 297-302 in prolyl aminopeptidase from *Serratia marcescens*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 50-53, 154-158 and 206-210 (numbering of amino acids according to SEQ ID NO:18). It is preferred that the prolyl aminopeptidase from *Serratia marcescens* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the S51 class of serine proteases or has at least 70% identity on the amino acid level to a protein of the S51 class of serine proteases and/or has a tertiary structure similar to aspartyl dipeptidase from *Escherichia coli*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 8-16, 38-46, 85-92, 132-140, 159-170 and 205-211 in aspartyl dipeptidase from *Escherichia coli*, and more preferably at one or more positions from the group of



positions that correspond structurally or by amino acid sequence homology to the regions 10-14, 87-90, 134-138 and 160-165 (numbering of amino acids according to SEQ ID NO:19). It is preferred that the aspartyl dipeptidase from *Escherichia coli* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the A2 class of aspartic proteases or has at least 70% identity on the amino acid level to a protein of the A2 class of aspartic proteases and/or has a tertiary structure similar to the protease from human immunodeficiency virus. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 5-12, 17-23, 27-30, 33-38 and 77-83 in protease from human immunodeficiency virus, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 7-10, 18-21, 34-37 and 79-82 (numbering of amino acids according to SEQ ID NO:20). It is preferred that the protease from human immunodeficiency virus, preferably HIV-1 protease, or a derivative or homologue thereof is used as the scaffold.

In an further embodiment the protein scaffold belongs to the A26 class of aspartic proteases or has at least 70% identity on the amino acid level to a protein of the A26 class of aspartic proteases and/or has a tertiary structure similar to the omptin from *Escherichia coli*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 28-40, 86-98, 150-168, 213-219 and 267-278 in omptin from *Escherichia coli*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 33-38, 161-168 and 273-277 (numbering of amino acids according to SEQ ID NO:21). It is preferred that the omptin from *Escherichia coli* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C1 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C1 class of cysteine proteases and/or has a tertiary structure similar to the

papain from *Carica papaya*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 17-24, 61-68, 88-95, 135-142, 153-158 and 176-184 in papain from *Carica papaya*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 63-66, 136-139 and 177-181 (numbering of amino acids according to SEQ ID NO:22). It is preferred that the papain from *Carica papaya* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C2 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C2 class of cysteine proteases and/or has a tertiary structure similar to human calpain-2. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 90-103, 160-172, 193-199, 243-260, 286-294 and 316-322 in human calpain-2, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 92-101, 245-250 and 287-291 (numbering of amino acids according to SEQ ID NO:23). It is preferred that the human calpain-2 or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C4 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C4 class of cysteine proteases and/or has a tertiary structure similar to Nla protease from tobacco etch virus. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 23-31, 112-120, 144-150, 168-176 and 205-218 in Nla protease from tobacco etch virus, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 145-149, 169-174 and 212-218 (numbering of amino acids according to SEQ ID NO:24). It is preferred that the Nla protease from tobacco etch virus (TEV protease) or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C10 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C10 class of cysteine proteases and/or has a tertiary structure similar to the streptopain from *Streptococcus pyogenes*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 81-90, 133-140, 150-164, 191-199, 219-229, 246-256, 306-312 and 330-337 in streptopain from *Streptococcus pyogenes*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 82-87, 134-138, 250-254 and 331-335 (numbering of amino acids according to SEQ ID NO:25). It is preferred that the streptopain from *Streptococcus pyogenes* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C19 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C19 class of cysteine proteases and/or has a tertiary structure similar to human ubiquitin specific protease 7. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 3-15, 63-70, 80-86, 248-256, 272-283 and 292-304 in human ubiquitin specific protease 7, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 10-15, 251-255, 277-281 and 298-304 (numbering of amino acids according to SEQ ID NO:26). It is preferred that the human ubiquitin specific protease 7 or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C47 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C47 class of cysteine proteases and/or has a tertiary structure similar to the staphopain from *Staphylococcus aureus*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 15-23, 57-66, 108-119, 142-149 and 157-164 in staphopain from *Staphylococcus aureus*, and more preferably at one or more positions from the group of positions

that correspond structurally or by amino acid sequence homology to the regions 17-22, 111-117, 143-147 and 159-163 (numbering of amino acids according to SEQ ID NO:27). It is preferred that the staphopain from *Staphylococcus aureus* or a derivative or homologue thereof is used as the scaffold.

In an further embodiment the protein scaffold belongs to the C48 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C48 class of cysteine proteases and/or has a tertiary structure similar to the Ulp1 endopeptidase from *Saccharomyces cerevisiae*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 40-51, 108-115, 132-141, 173-179 and 597-605 in Ulp1 endopeptidase from *Saccharomyces cerevisiae*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 43-49, 110-113, 133-137 and 175-178 (numbering of amino acids according to SEQ ID NO:28). It is preferred that the Ulp1 endopeptidase from *Saccharomyces cerevisiae* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the C56 class of cysteine proteases or has at least 70% identity on the amino acid level to a protein of the C56 class of cysteine proteases and/or has a tertiary structure similar to the Pfpl endopeptidase from *Pyrococcus horikoshii*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 8-16, 40-47, 66-73, 118-125 and 147-153 in Pfpl endopeptidase from *Pyrococcus horikoshii*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 9-14, 68-71, 120-123 and 148-151 (numbering of amino acids according to SEQ ID NO:29). It is preferred that the Pfpl endopeptidase from *Pyrococcus horikoshii* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the M4 class of metallo proteases or has at least 70% identity on the amino acid level to a protein of the M4 class of metallo proteases and/or has a tertiary structure similar to

thermolysin from *Bacillus thermoproteolyticus*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 106-118, 125-130, 152-160, 197-204, 210-213 and 221-229 in thermolysin from *Bacillus thermoproteolyticus*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 108-115, 126-129, 199-203 and 223-227 (numbering of amino acids according to SEQ ID NO:30). It is preferred that the thermolysin from *Bacillus thermoproteolyticus* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the M10 class of metallo proteases or has at least 70% identity on the amino acid level to a protein of the M10 class of metallo proteases and/or has a tertiary structure similar to human collagenase. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 2-7, 68-79, 85-90, 107-111 and 135-141 in human collagenase, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 3-6, 71-78 and 136-140 (numbering of amino acids according to SEQ ID NO:31). It is preferred that human collagenase or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzymes have glycosidase activity. A particularly suited protein scaffold for this variant is a glycosylase or is derived from a glycosylase. Preferably, the tertiary structure belongs to one of the following structural classes: class GH13, GH7, GH12, GH11, GH10, GH28, GH26, and GH18 (beta/alpha)<sub>8</sub> barrel.

In a first embodiment the protein scaffold belongs to the GH13 class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH13 class of glycosylases and/or has a tertiary structure similar to human pancreatic alpha-amylase. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 50-60, 100-110,

148-167, 235-244, 302-310 and 346-359 in human pancreatic alpha-amylase, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 51-58, 148-155 and 303-309 (numbering of amino acids according to SEQ ID NO:32). It is preferred that human pancreatic alpha-amylase or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GH7 class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH7 class of glycosylases and/or has a tertiary structure similar to cellulase from *Trichoderma reesei*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 47-56, 93-104, 173-182, 215-223, 229-236 and 322-334 in cellulase from *Trichoderma reesei*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 175-180, 218-222 and 324-332 (numbering of amino acids according to SEQ ID NO:33). It is preferred that cellulase from *Trichoderma reesei* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GH12 class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH12 class of glycosylases and/or has a tertiary structure similar to cellulase from *Aspergillus niger*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 18-28, 55-60, 106-113, 126-132 and 149-159 in cellulase from *Aspergillus niger*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 20-26, 56-59, 108-112 and 151-156 (numbering of amino acids according to SEQ ID NO:34). It is preferred that cellulase from *Aspergillus niger* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GH11 class of glycosylases or has at least 70% identity on the amino acid level to a protein of

the GH11 class of glycosylases and/or has a tertiary structure similar to xylanase from *Aspergillus niger*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 7-14, 33-39, 88-97, 114-126 and 158-167 in xylanase from *Aspergillus niger*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 20-26, 56-59, 108-112 and 151-156 (numbering of amino acids according to SEQ ID NO:35). It is preferred that xylanase from *Aspergillus niger* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GH10 class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH10 class of glycosylases and/or has a tertiary structure similar to xylanase from *Streptomyces lividans*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 21-29, 42-50, 84-92, 130-136, 206-217 and 269-278 in xylanase from *Streptomyces lividans*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 43-49, 86-90, 208-213 and 271-276 (numbering of amino acids according to SEQ ID NO:36). It is preferred that xylanase from *Streptomyces lividans* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GH28 class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH28 class of glycosylases and/or has a tertiary structure similar to pectinase from *Aspergillus niger*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 82-88, 118-126, 171-178, 228-236, 256-264 and 289-299 in pectinase from *Aspergillus niger*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 116-124, 174-178 and 291-296 (numbering of amino acids according

to SEQ ID NO:37). It is preferred that pectinase from *Aspergillus niger* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GH26 class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH26 class of glycosylases and/or has a tertiary structure similar to mannanase from *Pseudomonas cellulosa*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 75-83, 113-125, 174-182, 217-224, 247-254, 324-332 and 325-340 in mannanase from *Pseudomonas cellulosa*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 115-123, 176-180, 286-291 and 328-337 (numbering of amino acids according to SEQ ID NO:38). It is preferred that mannanase from *Pseudomonas cellulosa* or a derivative or homologue thereof is used as the scaffold.

In an further embodiment the protein scaffold belongs to the GH18 (beta/alpha)<sub>8</sub> barrel class of glycosylases or has at least 70% identity on the amino acid level to a protein of the GH18 class of glycosylases and/or has a tertiary structure similar to chitinase from *Bacillus circulans*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 21-29, 57-65, 130-136, 176-183, 221-229, 249-257 and 327-337 in chitinase from *Bacillus circulans*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 59-63, 178-181, 250-254 and 330-336 (numbering of amino acids according to SEQ ID NO:39). It is preferred that chitinase from *Bacillus circulans* or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzymes have esterhydrolase activity. Preferably, the protein scaffold for this variant have lipase, phosphatase, phytase, or phosphodiesterase activity.



In a first embodiment the protein scaffold belongs to the GX class of esterases or has at least 70% identity on the amino acid level to a protein of the GX class of esterases and/or has a tertiary structure similar to the structure of the lipase B from *Candida antarctica*. Preferably, the scaffold has lipase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 139-148, 188-195, 216-224, 256-266, 272-287 in lipase B from *Candida antarctica*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 141-146, 218-222, 259-263 and 275-283 (numbering of amino acids according to SEQ ID NO:40). It is preferred that lipase B from *Candida antarctica* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GX class of esterases or has at least 70% identity on the amino acid level to a protein of the GX class of esterases and/or has a tertiary structure similar to the pancreatic lipase from guinea pig. Preferably, the scaffold has lipase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 78-90, 91-100, 112-120, 179-186, 207-218, 238-247 and 248-260 in pancreatic lipase from guinea pig, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 80-87, 114-118, 209-215 and 239-246 (numbering of amino acids according to SEQ ID NO:41). It is preferred that pancreatic lipase from guinea pig or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold has a tertiary structure similar to the structure of the alkaline phosphatase from *Escherichia coli* or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the alkaline phosphatase from *Escherichia coli*. Preferably, the scaffold has phosphatase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the

regions 110-122, 187-142, 170-175, 186-193, 280-287 and 425-435 in alkaline phosphatase from *Escherichia coli*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 171-174, 187-191, 282-286 and 426-433 (numbering of amino acids according to SEQ ID NO:42). It is preferred that alkaline phosphatase from *Escherichia coli* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold has a tertiary structure similar to the structure of the bovine pancreatic desoxyribonuclease I or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the bovine pancreatic desoxyribonuclease I. Preferably, the scaffold has phosphodiesterase activity. More preferably, a nuclease, and most preferably, an unspecific endonuclease or a derivative thereof is used as the scaffold. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 14-21, 41-47, 72-77, 97-111, 135-143, 171-178, 202-209 and 242-251 in bovine pancreatic desoxyribonuclease I, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 16-19, 42-46, 136-141 and 172-176 (numbering of amino acids according to SEQ ID NO:43). It is preferred that bovine pancreatic desoxyribonuclease I or human desoxyribonuclease I or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzyme has transferase activity. A particularly suited protein scaffold for this variant is a glycosyl-, a phospho- or a methyltransferase, or is a derivative thereof. Particularly preferred protein scaffolds for this variant are glycosyltransferases or are derived from glycosyltransferases. The tertiary structure of the protein scaffold can be of any type. Preferably, however, the tertiary structure belongs to one of the following structural classes: GH13 and GT1.

In a first embodiment the protein scaffold belongs to the GH13 class of transferases or has at least 70% identity on the amino acid level to a protein of

the GH13 class of transferases and/or has a tertiary structure similar to the structure of the cyclomaltodextrin glucanotransferase from *Bacillus circulans*. Preferably, the scaffold has transferase activity, and more preferably a glycosyltransferase is used as the scaffold. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 38-48, 85-94, 142-154, 178-186, 259-266, 331-340 and 367-377 in cyclomaltodextrin glucanotransferase from *Bacillus circulans*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 87-92, 180-185, 261-264 and 269-275 (numbering of amino acids according to SEQ ID NO:44). It is preferred that cyclomaltodextrin glucanotransferase from *Bacillus circulans* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold belongs to the GT1 class of transferases or has at least 70% identity on the amino acid level to a protein of the GT1 class of transferases and/or has a tertiary structure similar to the structure of the glycosyltransferase from *Amycolatopsis orientalis* A82846. Preferably the scaffold has transferase activity, and more preferably glycosyltransferase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 58-74, 130-138, 185-193, 228-236 and 314-323 in glycosyltransferase from *Amycolatopsis orientalis* A82846, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 61-71, 230-234 and 316-321 (numbering of amino acids according to SEQ ID NO:45). It is preferred that the glycosyltransferase from *Amycolatopsis orientalis* A82846 or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzymes have oxidoreductase activity. A particularly suited protein scaffold for this variant is a monooxygenase, a dioxygenase or an alcohol dehydrogenase, or a derivative thereof. The tertiary structure of the protein scaffold can be of any type.

In a first embodiment the protein scaffold has a tertiary structure similar to the structure of the 2,3-diphydroxybiphenyl dioxygenase from *Pseudomonas* sp. or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the 2,3-diphydroxybiphenyl dioxygenase from *Pseudomonas* sp. Preferably, the scaffold has dioxygenase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 172-185, 198-206, 231-237, 250-259 and 282-287 in 2,3-diphydroxybiphenyl dioxygenase from *Pseudomonas* sp., and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 175-182, 200-204, 252-257 and 284-287 (numbering of amino acids according to SEQ ID NO:46). It is preferred that the 2,3-diphydroxybiphenyl dioxygenase from *Pseudomonas* sp or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold has a tertiary structure similar to the structure of the catechol dioxygenase from *Acinetobacter* sp. or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the catechol dioxygenase from *Acinetobacter* sp.. Preferably, the scaffold has dioxygenase activity, and more preferably catechol dioxygenase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 66-72, 105-112, 156-171 and 198-207 in catechol dioxygenase from *Acinetobacter* sp., and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 107-110, 161-171 and 201-205 (numbering of amino acids according to SEQ ID NO:47). It is preferred that the catechol dioxygenase from *Acinetobacter* sp or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold has a tertiary structure similar to the structure of the camphor-5-monooxygenase from *Pseudomonas putida* or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the camphor-5-monooxygenase from

*Pseudomonas putida*. Preferably, the scaffold has monooxygenase activity, and more preferably camphor monooxygenase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 26-31, 57-63, 84-98, 182-191, 242-256, 292-299 and 392-399 in camphor-5-monooxygenase from *Pseudomonas putida*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 85-96, 183-188, 244-253, 293-298 and 393-398 (numbering of amino acids according to SEQ ID NO:48). It is preferred that the camphor-5-monooxygenase from *Pseudomonas putida* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold has a tertiary structure similar to the structure of the alcohol dehydrogenase from *Equus caballus* or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the alcohol dehydrogenase from *Equus caballus*. Preferably, the scaffold has alcohol dehydrogenase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 49-63, 111-112, 294-301 and 361-369 in alcohol dehydrogenase from *Equus caballus*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 51-61 and 295-299 (numbering of amino acids according to SEQ ID NO:49). It is preferred that the alcohol dehydrogenase from *Equus caballus* or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzymes have lyase activity. A particularly suited protein scaffold for this variant is a oxoacid lyase or is a derivative thereof. Particularly preferred protein scaffolds for this variant are aldolases or synthases, or are derived thereof. The tertiary structure of the protein scaffold can be of any type, but a (beta/alpha)<sub>8</sub> barrel structure is preferred.

In a first embodiment the protein scaffold has a tertiary structure similar to the structure of the N-acetyl-d-neuramic acid aldolase from *Escherichia coli* or has at

least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the N-acetyl-d-neuramic acid aldolase from *Escherichia coli*. Preferably, the scaffold has aldolase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 45-55, 78-87, 105-113, 137-146, 164-171, 187-193, 205-210, 244-255 and 269-276 in N-acetyl-d-neuramic acid aldolase from *Escherichia coli*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 45-52, 138-144, 189-192, 247-253 and 271-275 (numbering of amino acids according to SEQ ID NO:50). It is preferred that the N-acetyl-d-neuramic acid aldolase from *Escherichia coli* or a derivative or homologue thereof is used as the scaffold.

In a further embodiment the protein scaffold has a tertiary structure similar to the structure of the tryptophan synthase from *Salmonella typhimurium* or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the tryptophan synthase from *Salmonella typhimurium*. Preferably, the scaffold has synthase activity. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 56-63, 127-134, 154-161, 175-193, 209-216 and 230-240 in tryptophan synthase from *Salmonella typhimurium*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 57-62, 155-160, 178-190 and 210-215 (numbering of amino acids according to SEQ ID NO:51). It is preferred that the tryptophan synthase from *Salmonella typhimurium* or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzymes have isomerase activity. A particularly suited protein scaffold for this variant is a converting aldose or a converting ketose, or is a derivative thereof.

In a first embodiment, the protein scaffold has a tertiary structure similar to the structure of the xylose isomerase from *Actinoplanes missouriensis* or has at least

70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the xylose isomerase from *Actinoplanes missouriensis*. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 18-31, 92-103, 136-147, 178-188 and 250-257 in xylose isomerase from *Actinoplanes missouriensis*, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 20-27, 92-99 and 180-186 (numbering of amino acids according to SEQ ID NO:52). It is preferred that the xylose isomerase from *Actinoplanes missouriensis* or a derivative or homologue thereof is used as the scaffold.

It is further preferred that the engineered enzymes have ligase activity. A particularly suited protein scaffold for this variant is a DNA ligase, or is a derivative thereof.

In a first embodiment, the protein scaffold has a tertiary structure similar to the structure of the DNA ligase from Bacteriophage T7 or has at least 70% identity on the amino acid level to a protein that has a tertiary structure similar to the structure of the DNA-ligase from Bacteriophage T7. It is preferred that SDRs are inserted into the protein scaffold at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 52-60, 94-108, 119-131, 241-248, 255-263 and 302-318 in DNA ligase from Bacteriophage T7, and more preferably at one or more positions from the group of positions that correspond structurally or by amino acid sequence homology to the regions 96-106, 121-129, 256-262 and 304-316 (numbering of amino acids according to SEQ ID NO:53). It is preferred that the DNA ligase from Bacteriophage T7 or a derivative or homologue thereof is used as the scaffold.

A third aspect of the invention is directed to a method for generating engineered enzymes with specificities that are qualitatively and/or quantitatively novel in combination with the protein scaffold. The inventive method comprises at least the following steps:

- (a) providing a protein scaffold capable to catalyze at least one chemical reaction on at least one target substrate,

- (b) generating a library of engineered enzymes or isolated engineered enzymes by combining the protein scaffold from step (a) with one or more fully or partially random peptide sequences at sites in the protein scaffold that enable the resulting engineered enzyme to discriminate between at least one target substrate and one or more different substrates and
- (c) selecting out of the library of engineered enzymes generated in step (b) one or more enzymes that have defined specificities towards at least one target substrate.

In a first variant of this aspect of the invention, the inventive method comprises at least the following steps:

- (a) providing a protein scaffold capable to catalyze at least one chemical reaction on at least one target substrate,
- (b) generating a library of engineered enzymes or isolated engineered enzymes by inserting into the protein scaffold from step (a) one or more fully or partially random peptide sequences at sites in the protein scaffold that enable the resulting engineered enzyme to discriminate between at least one target substrate and one or more different substrates and
- (c) selecting out of the library of engineered enzymes generated in step (b) one or more enzymes that have defined specificities towards at least one target substrate.

Preferably, the positions at which the one or more fully or partially random peptide sequences are combined with or inserted into the protein scaffold are identified prior to the combination or insertion.

The number of insertions or other combinations of fully or partially random peptide sequences as well as their length may vary over a wide range. The number is at least one, preferably more than one, more preferably between two and eleven, most preferably between two and six. The length of such fully or partially random peptide sequences is usually less than 50 amino acid residues. Preferably, the length is between one and 15 amino acid residues, more preferably between one and six amino acid residues. Alternatively, the length is between two and 20 amino acid residues, preferably between two and ten amino acid residues, more preferably between three and eight amino acid residues.



Preferably such insertions or other combinations are performed on the DNA level, using polynucleotides encoding such protein scaffolds and polynucleotides or oligonucleotides encoding such fully or partially random peptide sequences.

Optionally, steps (a) to (c) are repeated cyclically, whereby enzymes selected in step (c) serve as the protein scaffold in step (a) of a further cycle, and randomized peptide sequences are either inserted or, alternatively, substituted for peptide sequences that have been inserted in former cycles. Thereby, the number of inserted peptide sequences is either constant or increases over the cycles. The cycles are repeated until one or more enzymes with the intended specificities are generated.

Moreover, during or after one or more rounds of steps (a) to (c), the scaffold may be mutated at one or more positions in order to make the scaffold more acceptable for the combination with SDR sequences, and/or to increase catalytic activity at a specific pH and temperature, and/or to change the glycosylation pattern, and/or to decrease sensitivity towards enzyme inhibitors, and/or to change enzyme stability.

In a second variant of this aspect of the invention, the inventive method comprises at least the following steps:

- (a) providing a first protein scaffold fragment,
- (b) connecting said protein scaffold fragment via a peptide linkage with a first SDR, and optionally
- (c) connecting the product of step (b) via a peptide linkage with a further SDR peptide or with a further protein scaffold fragment, and optionally
- (d) repeating step (c) for as many cycles as necessary in order to generate a sufficiently specific enzyme, and
- (e) selecting out of the population generated in steps (a) – (d) one or more enzymes that have the desired specificities toward the one or more target substrates.

Protein scaffold fragment means a part of the sequence of a protein scaffold. A protein scaffold is comprised of at least two protein scaffold fragments.

In a third variant of this aspect of the invention, the protein scaffold, the SDRs and the engineered enzyme are encoded by a DNA sequence and an expression system is used in order to produce the protein. In an alternative variant, the protein scaffold, the SDRs and/or the engineered enzyme are chemically synthesized from peptide building blocks.

In a fourth variant of this aspect of the invention, the inventive method comprises at least the following steps:

- (a) providing a polynucleotide encoding a protein scaffold capable of catalyzing one or more chemical reactions on one or more target substrates;
- (b) combining one or more fully or partially random oligonucleotide sequence with the polynucleotide encoding the protein scaffold, the fully or partially random oligonucleotide sequences being located at sites in the polynucleotide that enable the encoded engineered enzyme to discriminate between the one or more target substrates and one or more other substrates; and
- (c) selecting out of the population generated in step (b) one or more polynucleotides that encode enzymes that have the defined specificities toward the one or more target substrates.

Any enzyme can serve as the protein scaffold in step (a). It can be a naturally occurring enzyme, a variant or a truncated derivate therefore, or an engineered enzyme. For human therapeutic use, the protein scaffold is preferably a mammalian enzyme, and more preferably a human enzyme. In that aspect, the invention is directed to a method for the generation of essentially mammalian, especially of essentially human enzymes with specificities that are different from specificities of any enzyme encoded in mammalian genomes or in the human genome, respectively.

According to the invention, the protein scaffold provided in step (a) of this aspect requires to be capable of catalyzing one or more chemical reactions on a target substrate. Therefore, a protein scaffold is selected from the group of potential protein scaffolds by its activity on the target substrate.

In a preferred variant of this aspect of the invention, a protein scaffold with hydrolase activity is used. Preferably, a protein scaffold with proteolytic activity

is used, and more preferably, a protease with very low specificity having basic activity on the target substrate is used as the protein scaffold. Examples of proteases from different structural classes with low substrate specificity are Papain, Trypsin, Chymotrypsin, Subtilisin, SET (trypsin-like serine protease from *Streptomyces erythraeus*), Elastase, Cathepsin G or Chymase. Before being employed as the protein scaffold, the amino acid sequence of the protease may be modified in order to change protein properties other than specificity, e.g. catalytic activity, stability, inhibitor sensitivity, or expression yield, essentially as described in WO 92/18645, or in order to change specificity, essentially as described in EP 02020576.3 and PCT/EP03/04864.

Another option for a feasible protein scaffold are lipases. Hepatic lipase, lipoprotein lipase and pancreatic lipase belong to the "lipoprotein lipase superfamily", which in turn is an example of the GX-class of lipases (M. Fischer, J. Pleiss (2003), Nucl. Acid. Res., 31, 319-321). The substrate specificity of lipases can be characterized by their relative activity towards triglycerol esters of fatty acids and phospholipids, bearing a charged head group. Alternatively, other hydrolases such as esterases, glycosylases, amidases, or nitrilases may be used as scaffolds.

Transferases are also feasible protein scaffolds. Glycosyltransferases are involved in many biological synthesis involving a variety of donors and acceptors. Alternatively, the protein scaffold may have ligase, lyase, oxidoreductase, or isomerase activity.

In a first embodiment, the one or more fully or partially random peptide sequences are inserted at specific sites in the protein scaffold. These insertion sites are characterized by the fact that the inserted peptide sequences can act as discriminators between different substrates, i.e. as Specificity Determining Regions or SDRs. Such insertion sites can be identified by several approaches. Preferably, insertion sites are identified by analysis of the three-dimensional structure of the protein scaffolds, by comparative analysis of the primary sequences of the protein scaffold with other enzymes having different quantitative specificities, or experimentally by techniques such as alanine scanning, random mutagenesis, or random deletion, or by any combination thereof.

A first approach to identify insertion sites for SDRs bases on the three-dimensional structure of the protein scaffold as it can be obtained by x-ray crystallography or by nuclear magnetic resonance studies. Structural alignment of the protein scaffold in comparison with other enzymes of the same structural class but having different quantitative specificities reveals regions of high structural similarity and regions with low structural similarity. Such an analysis can for example be done using public software such as Swiss PDB viewer (Guex, N. and Peitsch, M.C. (1997) *Electrophoresis* 18, 2714-2723). Regions of low structural similarity are preferred SDR insertion sites.

In a second approach to identify insertion sites for SDRs, three-dimensional structures of the scaffold protein in complex with competitive inhibitors or substrate analogs are analysed. It is assumed that the binding site of a competitive inhibitor significantly overlaps with the binding site of the substrate. In that case, atoms of the protein that are within a certain distance of atoms of the inhibitor are likely to be in a similar distance to the substrate as well. Choosing a short distance, e.g.  $< 5 \text{ \AA}$ , will result in an ensemble of protein atoms that are in close contact with the substrate. These residues would constitute the first shell contacts and are therefore preferred insertion sites for SDRs. Once first shell contacts have been identified, second shell contacts can be found by repeating the distance analysis starting from first shell atoms. In yet another alternative of the invention the distance analysis described above is performed starting from the active site residues.

In third approach to identify insertion sites for SDRs, the primary sequence of the scaffold protein is aligned with other enzymes of the same structural class but having different quantitative specificities using an alignment algorithm. Examples of such alignment algorithms are published (Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990) *J. Mol. Biol.* 215:403-410; "Statistical methods in Bioinformatics: an introduction" by Ewens, W. & Grant, G.R. 2001, Springer, New York). Such an alignment may reveal conserved and non-conserved regions with varying sequence homology, and, in particular, additional sequence elements in one or more enzymes compared to the scaffold protein. Conserved regions of are more likely to contribute to phenotypes shared

among the different proteins, e.g. stabilizing the three-dimensional fold. Non-conserved regions and, in particular, additional sequences in enzymes with quantitatively higher specificity (Turner, R. et al. (2002) *J. Biol. Chem.*, 277, 33068-33074) are preferred insertion sites for SDRs.

For proteases currently five families are known, namely aspartic-, cysteine-, serine-, metallo- and threonine proteases. Each family includes groups of proteases that share a similar fold. Crystallographic structures of members of these groups have been solved and are accessible through public databases, e.g. the Brookhaven protein database (H.M. Berman et al. *Nucleic Acids Research*, 28 pp. 235-242 (2000)). Such databases also include structural homologs in other enzyme classes and nonenzymatically active proteins of each class. Several tools are available to search public databases for structural homologues: SCOP - a structural classification of proteins database for the investigation of sequences and structures. (Murzin A. G. et al. (1995) *J. Mol. Biol.* 247, 536-540); CATH - Class, Architecture, Topology and Homologous superfamily: a hierarchical classification of protein domain structures (Orengo et al. (1997) *Structure* 5(8) 1093-1108); FSSP - Fold classification based on structure-structure alignment of proteins (Holm and Sander (1998) *Nucl. Acids Res.* 26 316-319); or VAST - Vector alignment search tool (Gibrat, Madej and Bryant (1996) *Current Opinion in Structural Biology* 6, 377-385).

In the above described approaches, members of structural classes are compared in order to identify insertion sites for SDRs.

In a preferred variant of these approaches serine proteases of the structural class S1 are compared with each other. Trypsin represents a member with low substrate specificity, as it requires only an arginine or lysine residue at the P<sub>1</sub> position. On the other hand, thrombin, tissue-type plasminogen activator or enterokinase all have a high specificity towards their substrate sequences, i.e. (L/I/V/F)XPR<sup>NA</sup>, CPGR<sup>NA</sup> VVGG and DDDK<sup>NA</sup>, respectively (Perona, J. & Craik, C. (1997) *J. Biol. Chem.*, 272, 29987-29990; Perona, J. & Craik, C (1995) *Protein Science*, 4, 337-360). An alignment of the amino acid sequences of these proteases is described in example 1 (Figure 2) along with the identification of SDRs.

A further example within the family of serine proteases is given by members of the structural class S8 (subtilisin fold). Subtilisin is the type protease for this class and represents an unspecific protease (Ottesen, M. & Svendsen, A. (1998) *Methods Enzymol.* 19, 199-215). Furin, PC1 and PC5 are proteases of the same structural class involved in the processing of propeptides and have a high substrate specificity (Seidah, N. & Chretien, M. (1997) *Curr. Opin. Biotech.*, 8: 602-607; Bergeron, F. et al. (2000) *J. Mol. Endocrin.*, 24:1-22). In a preferred variant of the approach alignments of the primary amino acids sequences (Figure 4) are used to identify eleven sequence stretches longer than three amino acids which specific proteases have in addition compared to subtilisin and are therefore potential specificity determining regions. In a further variant of the approach information from the three-dimensional structure of subtilisin can be used in order to further narrow down the selection (Figure 3). Out of the eleven inserted sequence stretches, three are especially close to the active site residues, namely stretch number 7, 8 and 11 which are insertions in PC5, PC1 and all three specific proteases, respectively (Figure 3). In a preferred variant, one or several amino acid stretches of variable length and composition can be inserted into the subtilisin sequence at one or several of the eleven positions. In a more preferred variant of the approach the insertion is performed at regions 7, 8 or 11 or any combination thereof. In another preferred variant of the approach protease scaffolds other than subtilisin from the structural class S8 are used.

In a further preferred variant of this approach, aspartic acid proteases of the structural class A1 are analyzed (Rawlings, N.D. & Barrett, A.J. (1995). *Methods Enzymol.* 248, 105-120; Chitpinitiyol, S. & Crabbe, M.J. (1998), *Food Chemistry*, 61, 395-418). Examples for the A1 structural class of aspartic proteases are pepsin with a low as well as beta-secretase (Grüniger-Leitch, F., et al. (2002) *J. Biol. Chem.* 277, 4687-4693) and renin (Wang, W. & Liang, T.C. (1994) *Biochemistry*, 33, 14636-14641) with relatively high substrate specificities. Retroviral proteases also belong to this class, although the active enzyme is a dimer of two identical subunits. The viral proteases are essential for the correct processing of the polyprotein precursor to generate functional proteins which requires a high substrate specificity in each case (Wu, J. et al. (1998) *Biochemistry*, 37, 4518-4526; Pettit, S. et al. (1991) *J. Biol. Chem.*, 266, 14539-

14547). Pepsin is the type protease for this class and represents an unspecific protease (Kageyama, T. (2002) *Cell. Mol. Life Sci.* 59, 288-306). B-secretase and Cathepsin D (Aguilar, C. F. et al. (1995) *Adv. Exp. Med. Biol.* 362, 155-166) are proteases of the same structural class and have a high substrate specificity. In a preferred variant of the approach alignments of the primary amino acids sequences (Figure 6) are used to identify six sequence stretches longer than three amino acids which are inserted in the specific proteases compared to pepsin and are therefore potential specificity determining regions. In a further variant of the approach information from the three-dimensional structure of b-secretase can be used in order to further narrow down the selection. Out of the six inserted sequence stretches, three are especially close to the active site residues, namely stretch number 1, 3 and 4 which are insertions in cathepsin D and beta-secretase, respectively (Figure 5). In a preferred variant of the approach, one or several amino acid stretches of variable length and composition can be inserted into the pepsin sequence at one or several of the six positions. In a more preferred embodiment of the invention the insertion is performed at the positions 1, 3 or 4 or any combination thereof. In another preferred embodiment of the invention protease scaffolds other than pepsin are used.

There are cases where a certain structural class does not include known members of bw and high specificity. This is exemplified by the C14 class of caspases which belong to the cysteine protease family (Rawlings, N.D. & Barrett, A.J. (1994) *Methods Enzymol.* 244, 461-486 ) and which all show high specificity for P<sub>4</sub> to P<sub>1</sub> positions. For example, caspase-1, caspase-3 and caspase-9 recognize the sequences YVAD<sup>^</sup>, DEVD<sup>^</sup> or LEHD<sup>^</sup>, respectively. Identification of the regions that differ between the caspases will include the regions responsible for the differences in substrate specificity (Figures 7 and 8).

Finally, non-enzymatic proteins of the same fold as the enzyme scaffold may also contribute to the identification of insertion sites for SDRs. For example, haptoglobin (Arcoleo, J. & Greer, J.; (1982) *J. Biol. Chem.* 257, 10063-10068) and azurocidin (Almeida, R. et al. (1991) *Biochem. Biophys. Res. Commun.* 177, 688-695) share the same chymotrypsin-like fold with all S1 proteases. Due to substitutions in the active site residues these proteins do not possess any proteolytic function, yet they show high homology with active proteases.

Differences between these proteins and specific proteases include regions that can serve as insertion sites for SDRs.

In a fourth approach, insertion sites for SDRs are identified experimentally by techniques such as alanine scanning, random mutagenesis, random insertion or random deletion. In contrast to the approach disclosed above, this approach does not require detailed knowledge about the three-dimensional structure of the scaffold protein. In one preferred variant of this approach, random mutagenesis of enzymes with relatively high specificity from the same structural class as the protein scaffold and screening for loss or change of specificity can be used to identify insertion sites for SDRs in the protein scaffold.

Random mutagenesis, alanine scanning, random insertion or random deletion are all done on the level of the polynucleotides encoding the enzymes. There are a variety of protocols known in the literature (e.g. Sambrook, J.F; Fritsch, E.F.; Maniatis, T.; Cold Spring Harbor Laboratory Press, Second Edition, 1989, New York). For example, random mutagenesis can be achieved by the use of a polymerase as described in patent WO 9218645. According to this patent, the one or more genes encoding the one or more proteases are amplified by use of a DNA polymerase with a high error rate or under conditions that increase the rate of misincorporations. For example the method of Cadwell and Joyce can be employed (Cadwell, R.C. and Joyce, G.F., PCR methods. Appl. 2 (1992) 28-33). Other methods of random mutagenesis such as, but not limited to, the use of mutator stains, chemical mutagens or UV-radiation can be employed as well.

Alternatively, oligonucleotides can be used for mutagenesis that substitute randomly distributed amino acid residues with an alanine. This method is generally referred to as alanine scanning mutagenesis (Fersht, A.R. Biochemistry (1989) 8031-8036). As a further alternative, modifications of the alanine scanning mutagenesis such as binominal mutagenesis (Gregoret, L.M. and Sauer, R.T. PNAS (1993) 4246-4250) or combinatorial alanine scanning (Weiss et al., PNAS (2000) 8950-8954) can be employed.

In order to express engineered enzymes, the DNA encoding such engineered proteins is ligated into a suitable expression vector by standard molecular cloning techniques (e.g. Sambrook, J.F; Fritsch, E.F.; Maniatis, T.; Cold Spring Harbor Laboratory Press, Second Edition, 1989, New York). The vector is introduced in a



suitable expression host cell, which expresses the corresponding engineered enzyme variant. Particularly suitable expression hosts are bacterial expression hosts such as *Escherichia coli* or *Bacillus subtilis*, or yeast expression hosts such as *Saccharomyces cerevisiae* or *Pichia pastoris*, or mammalian expression hosts such as Chinese Hamster Ovary (CHO) or Baby Hamster Kidney (BHK) cell lines, or viral expression systems such as bacteriophages like M13 or Lambda, or viruses such as the Baculovirus expression system. As a further alternative, systems for *in vitro* protein expression can be used. Typically, the DNA is ligated into an expression vector behind a suitable signal sequence that leads to secretion of the enzyme variants into the extracellular space, thereby allowing direct detection of protease activity in the cell supernatant. Particularly suitable signal sequences for *Escherichia coli* are HlyA, for *Bacillus subtilis* AprE, NprB, Mpr, AmyA, AmyE, Blac, SacB, and for *S. cerevisiae* Bar1, Suc2, Mat $\alpha$ , Inu1A, Ggplp. Alternatively, the enzyme variants are expressed intracellularly and the substrates are expressed also intracellularly. Preferably, this is done essentially as described in patent application WO 0212543, using a fusion peptide substrate comprising two auto-fluorescent proteins linked by the substrate amino-acid sequence. As a further alternative, after intracellular expression of the enzyme variants, or secretion into the periplasmatic space using signal sequences such as DsbA, PhoA, PelB, OmpA, OmpT or gIII for *Escherichia coli*, a permeabilisation or lysis step releases the enzyme variants into the supernatant. The destruction of the membrane barrier can be forced by the use of mechanical means such as ultrasonic, French press, or the use of membrane-digesting enzymes such as lysozyme. As another, further alternative, the genes encoding the enzyme variants are expressed cell-free by the use of a suitable cell-free expression system. For example, the S30 extract from *Escherichia coli* cells is used for this purpose as described by Lesly et al. (Methods in Molecular Biology 37 (1995) 265-278).

The ensemble of gene variants generated and expressed by any of the above methods are analyzed with respect to their affinity, substrate specificity or activity by appropriate assay and screening methods as described in detail for example in patent application PCT/EP03/04864. Genes from catalytically active variants having reduced specificity in comparison to the original enzyme are analyzed by sequencing. Sites at which mutations and/or insertions and/or

deletions occurred are preferred insertion sites at which SDRs can be inserted site-specifically.

In a second embodiment, the one or more fully or partially random peptide sequences are inserted at random sites in the protein scaffold. This modification is usually done on the polynucleotide level, i.e. by inserting nucleotide sequences into the gene that encodes the protein scaffold. Several methods are available that enable the random insertion of nucleotide sequences. Systems that can be used for random insertion are for example ligation based systems (Murakami et al. Nature Biotechnology 20 (2002) 76-81), systems based on DNA polymerisation and transposon based systems (e.g. GPS-M<sup>TM</sup> mutagenesis system, NEB Biolabs; MGS<sup>TM</sup> mutation generation system, Finnzymes). The transposon-based methods employ a transposase-mediated insertion of a selectable marker gene that contains at its termini recognition sequences for the transposase as well as two sites for a rare cutting restriction endonuclease. Using the latter endonuclease one usually releases the selection marker and after religation obtains an insertion. Instead of performing the religation one can alternatively insert a fragment that has terminal recognition sequences for one or two outside cutting restriction endonuclease as well as a selectable marker. After ligation, one releases this fragment using the one or two outside cutting endonucleases. After creating blunt ends by standard methods one inserts blunt ended random fragments at random positions into the gene.

In a further preferred embodiment, methods for homologous in-vitro recombination are used to combine the mutations introduced by the above mentioned methods to generate enzyme populations. Examples of methods that can be applied are the Recombination Chain Reaction (RCR) according to patent application WO 0134835, the DNA-Shuffling method according to the patent application WO 9522625, the Staggered Extension method according to patent WO 9842728, or the Random Priming recombination according to patent application WO9842728. Furthermore, also methods for non-homologous recombination such as the Itchy method can be applied (Ostermeier, M. et al. Nature Biotechnology 17 (1999) 1205-1209).

Upon random insertion of a nucleotide sequence into the protein scaffold one obtains a library of different genes encoding enzyme variants. The polynucleotide library is subsequently transferred to an appropriate expression vector. Upon

expression in a suitable host or by use of an in vitro expression system, a library of enzymes containing randomly inserted stretches of amino acids is obtained.

According to step (b) of this third aspect of the invention, one or more fully or partially random peptide sequences are inserted into the protein scaffold. The actual number of such inserted SDRs is determined by the intended quantitative specificity following the relation: the higher the intended specificity is, the more SDRs are inserted. Whereas a single SDR enables the generation of moderately specific enzymes, two SDRs enable already the generation of significantly specific enzymes. However, up to six and more SDRs can be inserted into a protein scaffold. A similar relation is valid for the length of the SDRs: the higher the intended specificity is, the longer are the SDRs that are to be inserted. SDRs can be as short as one to four amino acid residues. They can, however, also be as long as 50 amino acid residues. Significant specificity can already be generated by the use of SDRs of a length of four to six amino acid residues.

The peptid sequences that are inserted can be fully or partially random. In this context, fully random means that a set of sequences are inserted in parallel that includes sequences that differ from each other in each and every position. Partially random means that a set of sequences are inserted in parallel that includes sequences that differ from each other in at least one position. This difference can be either pair-wise or with respect to a single sequence. For example, when regarding an insertion of the length of four amino acids, partial random could be a set (i) that includes AGGG, GVGG, GGLG, GGGI, or (ii) that includes AGGG, VGGG, LGGG and IGGG. Alternatively, random sequences also comprises sequences that differ from each other in length. Randomization of the peptide sequences is achieved by randomization of the nucleotide sequences that are inserted into the gene at the respective sites. Thereby, randomization can be achieved by employing mixtures of nucleobases as monomers during chemical synthesis of the oligonucleotides. A particularly preferred mixture of monomers for a fully random codon that in addition minimizes the probability of stop codons is NN(GTC). Alternatively, random oligonucleotides can be obtained by fragmentation of DNA into short fragments that are inserted into the gene at the respective sites. The source of the DNA to be fragmented may be a synthetic oligonucleotide but alternatively may originate from cloned genes, cDNAs, or

genomic DNA. Preferably, the DNA is a gene encoding an enzyme. The fragmentation can, for example, be achieved by random endonucleolytic digestion of DNA. Preferably, an unspecific endonuclease such as DNase I (e.g. from bovine pancreas) is employed for the endonucleolytic digestion.

If steps (a) – (c) of the inventive method are repeated cyclically, there are different alternatives for obtaining random peptide sequences that are inserted in consecutive rounds. Preferably, SDRs that were identified in one round as leading to increased specificity of enzyme are used as templates for the random peptide sequences that are inserted in the following round.

In a preferred alternative, the sequences selected in one round are analysed and randomized oligonucleotides are generated based on these sequences. This can, for example, be achieved by using in addition to the original nucleotide with a certain percentage mixtures of the other three nucleotides monomers at each position in the oligonucleotide synthesis. If, for example, in a first round an SDR is identified that has the amino acid sequence ARLT, e.g. encoded by the nucleotide sequence GCG CGC CTT ACC, a random peptide sequence inserted in this SDR site could be encoded by an oligonucleotide with 70% G, 10% A, 10% T and 10% C at the first position, 70% C, 10% G, 10% T and 10% A at the second position, etc. This leads at each position approximately in 1 of 3 cases to the template amino acid and in 2 of 3 cases to another amino acid.

In another preferred alternative, the sequences selected in one round are analyzed and a consensus library is generated based on these sequences. This can, for example, be achieved by using defined mixtures of nucleotides at each position in the oligonucleotide synthesis in a way that leads to mixtures of the amino acid residues that were identified at each position of the SDR selected in the previous round. If, for example, in a first round two SDRs are identified that have the amino acid sequences ARLT and VPGS, a consensus library inserted in this SDR site in the following round could be encoded by an oligonucleotide with the sequence G(C/T)G C(G/C)C (G/T)(G/T)G (A/T)CC. This would correspond to the random peptide sequence (A/V)(R/P)(L/G/V/W)(T/S), thereby allowing all combinations of the amino acid residues identified in the first round, and, due to the degeneracy of the genetic code, allowing in addition to a lower degree alternative amino acid residues at some positions.

In another preferred alternative, the sequences selected in one round are, without previous analysis, recombined using methods for the *in vitro* recombination of polynucleotides, such as the methods described in WO 01/34835 (the following also provides details of the eighth and ninth aspect of the invention).

After insertion of the partially or fully random sequences into the gene encoding the scaffold protein, and eventually ligation of the resulting gene into a suitable expression vector using standard molecular cloning techniques (Sambrook, J.F.; Fritsch, E.F.; Maniatis, T.; Cold Spring Harbor Laboratory Press, Second Edition, 1989, New York), the vector is introduced in a suitable expression host cell which expresses the corresponding enzyme variant. Particularly suitable expression hosts are bacterial expression hosts such as *Escherichia coli* or *Bacillus subtilis*, or yeast expression hosts such as *Saccharomyces cerevisiae* or *Pichia pastoris*, or mammalian expression hosts such as Chinese Hamster Ovary (CHO) or Baby Hamster Kidney (BHK) cell lines, or viral expression systems such as bacteriophages like M13 T7 phage or Lambda, or viruses such as the Baculovirus expression system. As a further alternative, systems for *in vitro* protein expression can be used. Typically, the DNA is ligated into an expression vector behind a suitable signal sequence that leads to secretion of the enzyme variants into the extracellular space, thereby allowing direct detection of enzyme activity in the cell supernatant. Particularly suitable signal sequences for *Escherichia coli* are ompA, pelB, HlyA, for *Bacillus subtilis* AprE, NprB, Mpr, AmyA, AmyE, Blac, SacB, and for *S. cerevisiae* Bar1, Suc2, Mat $\alpha$ , Inu1A, Ggplp. Alternatively, the enzyme variants are expressed intracellularly and the substrates are expressed also intracellularly. According to protease variants this is done essentially as described in patent application WO 0212543, using a fusion peptide substrate comprising two auto-fluorescent proteins linked by the substrate amino-acid sequence. As a further alternative, after intracellular expression of the enzyme variants, or secretion into the periplasmic space using signal sequences such as DsbA, PhoA, PelB, OmpA, OmpT or gIII for *Escherichia coli*, a permeabilisation or lysis step releases the enzyme variants into the supernatant. The destruction of the membrane barrier can be forced by the use of mechanical means such as ultrasonic, French press, or the use of membrane-digesting enzymes such as

lysozyme. As another, further alternative, the genes encoding the enzyme variants are expressed cell-free by the use of a suitable cell-free expression system. For example, the S30 extract from *Escherichia coli* cells is used for this purpose as described by Lesly et al. (Methods in Molecular Biology 37 (1995) 265-278).

After introduction of the vector into host cells, these cells are screened for the expression of enzymes with specificity for the intended target substrate. Such screening is typically done by separating the cells from each other, in order to enable the correlation of genotype and phenotype, and assaying the activity of each cell clone after a growth and expression period. Such separation can for example be done by distribution of the cells into the compartments of sample carriers, e.g. as described in WO 01/24933. Alternatively, the cells are separated by streaking on agar plates, by enclosing in a polymer such as agarose, by filling into capillaries, or by similar methods.

Identification of variants with the intended specificity can be done by different approaches. In the case of proteases, preferably assays using peptide substrates essentially as described in PCT/EP03/04864 are employed.

Regardless of the expression format, selection of enzyme variants is done under conditions that allow identification of enzymes that recognize and convert the target sequence preferably. As a first alternative, enzymes that recognize and convert the target sequence preferably are identified by screening for enzymes with a high affinity for the target substrate sequence. High affinity corresponds to a low  $K_M$  which is selected by screening at target substrate concentrations substantially below the  $K_M$  of the first enzyme. Preferably, the substrates that are used are linked to one or more fluorophores that enable the detection of the modification of the substrate at concentrations below 10  $\mu$ M, preferably below 1  $\mu$ M, more preferably below 100 nM, and most preferably below 10 nM.

As a second alternative, enzymes that recognize and convert the target substrate preferably are identified by employing two or more substrates in the assay and screening for activity on these two or more substrates in comparison. Preferably, the two or more substrates employed are linked to different marker molecules,

thereby enabling the detection of the modification of the two or more substrates consecutively or in parallel. In the case of proteases, particularly preferably two peptide substrates are employed, one peptide substrate having an arbitrarily chosen or even partially or fully random amino-acid sequence thereby enabling to monitor the activity on an arbitrary substrate, and the other peptide substrate having an amino-acid sequence identical to or resembling the intended target substrate sequence thereby enabling to monitor the activity on the target substrate. Especially preferably, these two peptide substrates are linked to fluorescent marker molecules, and the fluorescent properties of the two peptide substrates are sufficiently different in order to distinguish both activities when measured consecutively or in parallel. For example, a fusion protein comprising a first autofluorescent protein, a peptide, and a second autofluorescent protein according to patent application WO 0212543 can be used for this purpose. Alternatively, fluorophores such as rhodamines are linked chemically to the peptide substrates.

As a third alternative, enzymes that recognize and convert the target substrate preferably are identified by employing one or more substrates resembling the target substrate together with competing substrates in high excess. Screening with respect to activity on the substrates resembling the target substrate is then done in the presence of the competing substrates. Enzymes having a specificity which corresponds qualitatively to the target specificity, but having only a low quantitative specificity are identified as negative samples in such a screen. Whereas enzymes having a specificity which corresponds qualitatively and quantitatively to the target specificity are identified positively. Preferably, the one or more substrates resembling the target substrate are linked to marker molecules, thereby enabling the detection of their modifications, whereas the competing substrates do not carry marker molecules. The competing substrates have arbitrarily chosen or random amino-acid sequences, thereby acting as competitive inhibitors for the hydrolysis of the marker-carrying substrates. For example, protein hydrolysates such as Trypton can serve as competing substrates for engineered proteolytic enzymes according to the invention.

As a fourth alternative, enzymes that recognize and convert the target substrate preferably are identified and selected by an amplification-coupled or growth-coupled selection step. Furthermore, the activity can be measured intracellularly

and the selection can be done by a cell sorter, such as a fluorescence-activated cell sorter.

As a further alternative, enzymes that recognize and convert the target substrate are identified by first selecting enzymes that preferentially bind to the target substrate, and secondly selecting out of this subgroup of enzyme variants those enzymes that convert the target substrate. Selection for enzymes that preferentially bind the target substrate can be either done by selection of binders to the target substrate or by counter-selection of enzymes that bind to other substrates. Methods for the selection of binders or for the counter-selection of non-binders is known in the art. Such methods typically require phenotype-genotype coupling which can be solved by using surface display expression methods. Such methods include, for example, phage or viral display, cell surface display and in vitro display. Phage or viral display typically involves fusion of the protein of interest to a viral/phage protein. Cell surface display, i.e. either bacterial or eukaryotic cell display, typically involves fusion of the protein of interest to a peptide or protein that is located at the cell surface. In in-vitro display, the protein is typically made in vitro and linked directly or indirectly to the mRNA encoding the protein (DE 19646372).

The invention also provides for a composition or pharmaceutical composition comprising one or more engineered enzymes according to the first aspect of the invention as defined herein before. The composition may optionally comprise an acceptable carrier, excipient and/or auxiliary agent.

Pharmaceutical compositions according to the invention may optionally comprise a pharmaceutically acceptable carrier. Pharmaceutical formulations are well known and pharmaceutical compositions may be routinely formulated by one having ordinary skill in the art. The composition can be formulated as a solution, suspension, emulsion, or lyophilized powder in association with a pharmaceutically acceptable vehicle. Examples of such vehicles are water, saline, Ringer's solution, dextrose solution, and human serum albumin. Liposomes and nonaqueous vehicles such as fixed oils may also be used. The vehicle or lyophilized powder may contain additives that maintain isotonicity (e.g. sodium



chloride, mannitol) and chemical stability (e.g. buffers and preservatives). The composition is sterilized by commonly used techniques.

The pharmaceutical composition of the present invention may be administered by any means that enables the active agent to reach the agent's site of action in the body of a mammal. Pharmaceutical compositions may be administered parentally, i.e. intravenous, subcutaneous, intramuscular.

Dosage varies depending upon known factors such as the pharmacodynamic characteristics of the particular agent, and its mode and route of administration ; age, health, and weight of the recipient; nature and extent of symptoms, kind of concurrent treatment, frequency of treatment, and the effect desired.

Non-pharmaceutical compositions as defined herein are research composition, nutritional composition, cleaning composition, disinfection composition, cosmetic composition or composition for personal care. Moreover, DNA sequences coding for the engineered enzyme as defined herein before and vectors containing said DNA sequences are also provided. Finally, transformed host cells (prokaryotic or eukaryotic) or transgenic organisms containing such DNA sequences and/or vectors, as well as a method utilizing such host cells or transgenic animals for producing the engineered enzyme of the first aspect of the invention are also contemplated.

#### Detailed description of the figures

Figure 1: Three-dimensional structure of human trypsin I with the active site residues shown in "ball-and-stick" representation and with the marked regions indicating potential SDR insertion sites.

Figure 2: Alignment of the primary amino acid sequences of the human proteases trypsin I, alpha-thrombin and enteropeptidase all of which belong to the structural class S1 of the serine protease family. Trypsin represents an unspecific protease of this structural class, while alpha-thrombin and enteropeptidase are proteases with high substrate specificity. Compared to trypsin several regions of insertions of three or more amino acids into the primary sequence of  $\alpha$ -thrombin and enterokinase are seen. The region marked with (-1-) and the region marked with (-3-) are preferred SDR insertion sites. In

the tertiary structure of alpha-thrombin both regions are in the vicinity of the substrate binding site. These regions therefore fulfill two criteria to be selected as candidates for SDRs: firstly, they represent insertions in the specific proteases compared to the unspecific one and, secondly, they are close to the substrate binding site. A representation of the three-dimensional structure is given in figure 3.

Figure 3: Three-dimensional structure of subtilisin with the active site residues being shown in "ball-and-stick" representation and with the numbered regions indicating potential SDR insertion sites.

Figure 4: Alignment of the primary amino acid sequences of subtilisin E, furin, PC1 and PC5 all of which belong to the structural class S8 of the serine protease family. Subtilisin E represents an unspecific protease of this structural class, while furin, PC1 and PC5 are proteases with high substrate specificity. Compared to subtilisin several regions of insertions of three or more amino acids into the primary sequence of furin, PC1 and PC5 are seen. The regions marked with (-4-), (-5-), (-7-), (-9-) and (-11-) are preferred SDR insertion sites. These regions stretches fulfill two criteria to be selected as candidates for SDRs: firstly, they represent insertions in the specific proteases compared to the unspecific one and, secondly, they are close to the active site residues.

Figure 5: Three-dimensional structure of beta-secretase with the active site residues being shown in "ball-and-stick" representation and with the numbered regions indicating potential SDR insertion sites.

Figure 6: Alignment of the primary amino acid sequences of pepsin, b-secretase and cathepsin D, all of which belong to the structural class A1 of the aspartic protease family. Pepsin represents an unspecific protease of this structural class, while b-secretase and cathepsin D are proteases with high substrate specificity. Compared to pepsin several regions of insertions of three or more amino acids into the primary sequence of b-secretase and cathepsin D are seen. The regions marked with -1- to -11- correspond to possible SDR combining sites and are also marked in Fig.5.

Figure 7: illustrates the three-dimensional structure of caspase 7 with the active site residues being shown in "ball-and-stick" representation and with the numbered regions indicating potential SDR insertion sites.

Figure 8: shows the primary amino acid sequence of caspase 7 as a member of the cysteine protease class C14 family (see also SEQ ID NO: 14).

Figure 9: Schematic representation of method according to the third aspect of the invention.

Figure 10: Western blot analysis of trypsin expression. Supernatant of cell cultures expressing variants of trypsin are compared to negative controls. Lane 1: molecular weight standard; lane 2: negative control; lane 3: supernatant of variant a; lane 4: negative control; lane 5: supernatant of variant b. A primary antibody specific to the expressed protein and a secondary antibody for generation of the signal were used.

Figure 11: Time course of the proteolytic cleavage of a target substrate. Supernatant of cells containing the vector with the gene for human trypsin and that of cells containing the vector without the gene was incubated with the peptide substrate described in the text. Cleavage of the peptide results in a decreased read out value. Proteolytic activity is confirmed for the positive clone.

Figure 12: Relative activity of three engineered proteolytic enzymes in comparison with human trypsin I on two different peptide substrates. A time course of the proteolytic digestion of the two substrates was performed and evaluated. Substrate B was used for screening and substrate A is a closely related sequence. Relative activity of the three variants was normalized to the activity of human trypsin I. Variant 1 and 2 clearly show increased specificity towards the target substrate. Variant 3, on the other hand, serves as a negative control with similar activities as the human trypsin I.

Figure 13: Relative specificities of trypsin and variants of engineered proteolytic enzymes with one or two SDRs, respectively. Activity of the proteases was determined in the presence and absence of competitor substrate, i.e. peptone at a concentration of 10mg/ml. Time courses for the proteolytic cleavage were

recorded and the time constants  $k$  determined. The ratios between the time constants with and without competitor were formed and represent a quantitative measure for the specificity of the protease. The ratios were normalized to trypsin. The specificity of the variant containing two SDRs is 2.5 fold higher than that of the variant with SDR2 alone.

Figure 14: Shows the relative specificities of protease variants in absence and presence of competitor substrate. The protease variants containing two inserts with different sequences and the non-modified scaffold human trypsin I were expressed in a suitable host. Activity of the protease variants was determined as the cleavage rate of a peptide with the desired target sequence of TNF-alpha in the absence and presence of competitor substrate. Specificity is expressed as the ratio of cleavage rates in the presence and absence of competitor.

Figure 15: The figure shows the reduction of cytotoxicity induced by human TNF-alpha when incubating the human TNF-alpha with concentrated supernatant from cultures expressing the inventive engineered proteolytic enzymes being specific for human TNF-alpha. This indicates the efficacy of the inventive engineered proteolytic enzymes.

Figure 16: The figure shows the reduction of cytotoxicity induced by human TNF-alpha when incubating the human TNF-alpha with different concentrations of purified inventive engineered proteolytic enzyme being specific for human TNF-alpha. Variant g comprises SEQ ID NO:72 as SDR1 and SEQ ID NO:73 as SDR2. This indicates the efficacy of the inventive engineered proteolytic enzymes.

Figure 17: The figure compares the activity of inventive engineered proteolytic enzymes being specific for human TNF-alpha with the activity of human trypsin I on two protein substrates: (a) human TNF-alpha; (b) mixture of human serum proteins. This indicates the safety of the inventive engineered proteolytic enzymes. Variant x corresponds to Seq ID No: 75 comprising the SDRs according to Seq ID No. 89 (SDR1) and 95 (SDR2). Variants xi and xii correspond to derivatives thereof comprising the same SDR sequences.

Figure 18: Specific hydrolysis of human VEGF by an engineered proteolytic enzyme derived from human trypsin.

### Examples

In the following examples, materials and methods of the present invention are provided including the determination of catalytic properties of enzymes obtained by the method. It should be understood that these examples are for illustrative purpose only and are not to be construed as limiting this invention in any manner. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

In the experimental examples described below, standard techniques of recombinant DNA technology were used that were described in various publications, e.g. Sambrook et al. (1989), Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, or Ausubel et al. (1987), Current Protocols in Molecular Biology 1987-1988, Wiley Interscience. Unless otherwise indicated, restriction enzymes, polymerases and other enzymes as well as DNA purification kits were used according to the manufacturers specifications.

#### Example I: Identification of SDR sites in human trypsin

Insertion sites for SDRs have been identified in the serine protease human trypsin I (structural class S1) by comparison with members of the same structural class having a higher sequence specificity. Trypsin represents a member with low substrate specificity, as it requires only an arginine or lysine residue at the P<sub>1</sub> position. On the other hand, thrombin, tissue-type plasminogen activator or enterokinase all have a high specificity towards their substrate sequences, i.e. (L/I/V/F)XPR<sup>^</sup> NA, CPGR<sup>^</sup> VVGG and DDDK<sup>^</sup>, respectively. The primary sequences and tertiary structures of these and further S1 serine proteases have been aligned in order to determine regions of low and high sequence and structure homology and especially regions that correspond to insertions in the sequences of the more specific proteases (Figure 2). Several regions of insertions equal or longer than 3 amino acids representing potential SDR sites have been identified as indicated in Figure 1. These regions were chosen as target sites for the insertion of SDRs in the examples below, e.g. SDR1 (region one in figure 2, after amino acid 42 according to SEQ ID NO:1) with a length of six and SDR2 (region three in figure 2, after amino acid 123 according to SEQ ID NO:1) with a length of five amino acids, respectively.

Example II: Molecular cloning of the human trypsin I gene to be used as scaffold protein and expression of the mature protease in *B. subtilis*

The gene encoding the unspecific protease human trypsinogen I was cloned into the vector pUC18. Cloning was done as follows: the coding sequence of the protein was amplified by PCR using primers that introduced a KpnI site at the 5' end and a BamHI site at the 3' end. This PCR fragment was cloned into the appropriate sites of the vector pUC18. Identity was confirmed by sequencing. After sequencing the coding sequence of the mature protein was amplified by PCR using primers that introduced different BglI sites at the 5' end and the 3' end.

This PCR fragment was cloned into the appropriate sites of an *E. coli* – *B. subtilis* shuttle vector. The vector contains a pMB1 origin for amplification in *E. coli*, a neomycin resistance marker for selection in *E. coli*, as well as a P43 promoter for the constitutive expression in *B. subtilis*. A 87 bp fragment that contains the leader sequence encoding the signal peptide from the *sacB* gene of *B. subtilis* was introduced behind the P43 promoter. Different BglI restriction sites serve as insertion sites for heterologous genes to be expressed.

Expression of human trypsin I was confirmed by measurement of the proteolytic activity in supernatant of cells containing the vector with the gene in comparison to a negative control. A peptide including an arginine cleavage site was chosen as a substrate. The peptide was Nterminally biotinylated and labeled with a fluorophore at the C-terminus. After incubation of the peptide with culture supernatant streptavidin was added. Uncleaved peptide associate with streptavidin and lead to a high read out value while cleavage results in low read out values. Figure 11 shows the time course of a proteolytic digestion of *B. subtilis* cells containing the vector with the trypsin I gene in comparison to *B. subtilis* cells containing the vector without the trypsin I gene (negative control).

As a further confirmation of expression of the protease, supernatants of cells containing the vector with the gene and control cells were analyzed by polyacrylamid gel electrophoreses and subsequent western blot using an antibody specific to the target protease. The procedure was performed according to standard methods (Sambrook, J.F; Fritsch, E.F.; Maniatis, T.; Cold Spring Harbor Laboratory Press, Second Edition, 1989, New York). Figure 8 confirms expression of the protein only in the cells harbouring the vector with the gene for trypsin.

Example III: Providing a scaffold protein

In this example, human trypsin I was used as the scaffold protein. The gene was either used in its natural form, or, alternatively, was modified to result in a scaffold protein with increased catalytic activity or further improved characteristics.

The modification was done by random modification of the gene, followed by expression of the enzyme and subsequent selection for increased activity. First, the gene was PCR amplified under error-prone conditions, essentially as described by Cadwell, R.C and Joyce, G.F. (PCR Methods Appl. 2 (1992) 28-33). Error-prone PCR was done using 30 pmol of each primer, 20 nmol dGTP and dATP, 100 nmol dCTP and dTTP, 20 fmol template, and 5 U Taq DNA polymerase in 10 mM Tris HCl pH 7.6, 50 mM KCl, 7 mM MgCl<sub>2</sub>, 0.5 mM MnCl<sub>2</sub>, 0.01 % gelatin for 20 cycles of 1 min at 94 °C, 1 min at 65 °C and 1 min at 72 °C. The resulting DNA library was purified using the Qiaquick PCR Purification Kit following the suppliers' instructions. The PCR product was digested with the restriction enzyme *Bgl*I and purified. Afterwards, the PCR product was ligated into the E. coli – B. subtilis shuttle vector described above which was digested with *Bgl*I and dephosphorylated. The ligation products were transformed into E. coli, amplified in LB, and the plasmids were purified using the Qiagen Plasmid Purification Kit following the suppliers' instructions. Resulting plasmids were transformed into B. subtilis cells.

Alternatively, or in addition to random mutagenesis, variants of the gene were statistically recombined at homologous positions by use of the Recombination Chain Reaction, essentially as described in WO 0134835. PCR products of the genes encoding the protease variants were purified using the QIAquick PCR Purification Kit following the suppliers' instructions, checked for correct size by agarose gel electrophoresis and mixed together in equimolar amounts. 80 µg of this PCR mix in 150 mM TrisHCl pH 7.6, 6.6 mM MgCl<sub>2</sub> were heated for 5 min at 94 °C and subsequently cooled down to 37 °C at 0.05 °C/s in order to re-anneal strands and thereby produce heteroduplexes in a stochastic manner. Then, 2.5 U Exonuclease III per µg DNA were added and incubated for 20, 40 or 60 min at 37 °C in order to digest different lengths from both 3' ends of the heteroduplexes. The partly digested PCR products were refilled with 0.6 U Pfu polymerase per µg DNA by incubating for 15 min at 72 °C in 0.17 mM dNTPs and Pfu polymerase

buffer according to the suppliers' instructions. After performing a single PCR cycle, the resulting DNA was purified using the QIAquick PCR Purification Kit following the suppliers' instructions, digested with BglI and ligated into the linearized vector. The ligation products were transformed into E. coli, amplified in LB containing ampicillin as marker, and the plasmids were purified using the Qiagen Plasmid Purification Kit following the suppliers' instructions. Resulting plasmids were transformed into B. subtilis cells.

Example IV: Insertion of SDRs into the protein scaffold of human trypsin I and generation of an engineered proteolytic enzyme with specificity for a peptide substrate having the sequence KKWLGRVPGGPV.

In order to create insertion sites for SDRs in human trypsin I, two pairs of different restriction sites were introduced into the gene at sites that were identified as potential SDR sites (see Example I above) without changing the amino acid sequence. The insertion of the restriction sites was done by overlap extension PCR. Primers restr1 and restr2 were used for the introduction of SacI and BamHI restriction sites, restr3 and restr4 were used for the introduction of KpnI and NheI restriction sites. The sequences of the primers were as follows:

Binding site for restr1 and restr2 and the corresponding amino acid sequence (SEQ ID NO:54):

5'-GGTGGTATCAGCAGGCCACTGCTACAAGTCCCGCATCCAGGT-3'  
           V  V  S  A  G  H  C  Y  K  S  R  I  Q

Forward primer restr1 (SEQ ID NO:56):

5'-GGTGGTATCCGCGGGCCACTGCTACAAGTCCCGGATCCAGGT-3'

Reverse primer restr2 (SEQ ID NO:57):

5'-ACCTGGATCCGGGACTTGCTAGCAGTGGCCCGCGGATACCACC-3'

Binding site for restr3 and restr4 and the corresponding amino acid sequence (SEQ ID NO:58):

5'-CCACTGGCACGAAGTGCCTCATCTCTGGCTGGGGCAACACTGCGAGCTCT-3'  
           T  G  T  K  C  L  I  S  G  W  G  N  T  A  S  S

Forward primer restr3 (SEQ ID NO:60):



5'-CCACTGGCACGAAGTGCCTCATCTCTGGCTGGGGCAACACTGCGAGCTCT-3'

Reverse primer restr4 (SEQ ID NO:61):

5'-AGAGCTAGCAGTGTGCCCCAGCCAGAGATGAGGCACTTGGTACCAGTGG-3'

In a first overlap extension PCR, the SacII/BamHI sites were introduced, enabling to insert SDR1, and in a second overlap extension PCR the KpnI/NheI sites, enabling the insertion of SDR2. The product of the overlap extension PCR was amplified using primers pUC-forward and pUC-reverse. The sequences of pUC-forward and pUC-reverse are as follows:

pUC-forward (SEQ ID NO:62): 5'-GGGGTACCCACCACCATGAATCCACTCCT-3'

pUC-reverse (SEQ ID NO:63): 5'-CGGGATCCGGTATAGAGACTGAAGAGATAC-3'

The restriction sites generated thereby were subsequently used to insert defined or random oligonucleotides into the SDR1 and SDR2 insertion sites by standard restriction and ligation methods. Typically, two complementary synthetic 5'-phosphorylated oligonucleotides were annealed and ligated into a vector carrying the modified human trypsin I gene that was cleaved with the respective restriction enzymes. Oligonucleotides encoding SDR1 were inserted via the SacII/BamHI sites whereas oligonucleotides encoding SDR2 were inserted via the KpnI/NheI sites. For each insertion an oligonucleotide pair according to the following general sequences was used ([P] indicating 5'-phosphorylation, N and X indicating any nucleotide or amino acid residue, respectively):

oligox-SDR1f (SEQ ID NO:64):

5'-[P]-GGGCCACTGCTACNNNNNNNNNNNNNNNNNNNAAGTCCCG-3'

oligox-SDR1r (SEQ ID NO:66):

3'-CGCCCGGTGACGATGNNNNNNNNNNNNNNNNNNNTTCAGGGCCTAG-[P]-5'

G H C Y X X X X X X X K S

oligox-SDR2f (SEQ ID NO:67):

5'-[P]-CAAGTGCCTCATCTCTGGCTGGGGCAACNNNNNNNNNNNNNNNNNACTG-3'

oligox-SDR2r (SEQ ID NO:69):

3'-CATGGTTCACGGAGTAGAGACCGACCCGTTGNNNNNNNNNNNNNNNNNTGACGATC-[P]-5'

K C L I S G W G N X X X X X X T

As an alternative to the above method, a PCR based method was used for the integration of random-sequences into the SDR1 and SDR2 insertion sites in the modified human trypsin I. For each SDR, one primer was used where the SDR region is fully randomized. Sequences of the primers were as follows (N = A/C/G/T, B = C/G/T, V = A/C/G):

Primer SDR1-mutnnb-forward (SEQ ID NO:70):

5'-TGGTATCCGCGGGCCACTGCTACNNBNNBNNBNNBNNBNNBAAGTCCCGGATCCAGGTG-3'

Primer SDR2-mutnnb-reverse (SEQ ID NO:71):

5'-GGCGCCAGAGCTAGCAGTVNNVNNVNNVNNVNNNGTTGCCCCAGCCAGAGATG-3'

The codon NNB, or VNN in the reverse strand, allows all 20 amino acids to be made, but reduces the probability of encoding a stop codon from 0.047 to 0.021.

As a further alternative, after identification of SDRs that lead to increased specificity, these SDRs were used as templates for further randomization. Thereby, random peptide sequences were inserted that were partially randomized at each position and partially identical at each position to the original sequence.

As an example, random peptide sequences that have in approximately 1 of 3 cases the template amino acid residue and in approximately 2 of 3 cases any other amino acid residue at each position were inserted into the two SDR insertion sites of the modified human trypsin I. For this purpose, primers that contain at each nucleotide position of the SDR approximately 70% of the template bases and 30% of a mixture of the three other bases were used.

With each primer pair a PCR was performed under standard conditions using the human trypsin I gene as template. The resulting DNA was purified using the QIAquick PCR Purification Kit following the suppliers' instructions and digested with SacII and NheI. After digestion the DNA was purified and ligated into the SacII and NheI digested and dephosphorylated vector. The ligation products were transformed into *E. coli*, amplified in LB containing the respective marker, and the plasmids were purified using the Qiagen Plasmid Purification Kit following the suppliers' instructions. Resulting plasmids were transformed into *B. subtilis* cells. These cells were then separated to single cells, grown to clones, and after expression of the protease gene screened for proteolytic activity.

The following substrates were employed for screening for proteolytic activity (SEQ ID NOs:76 and 77):

substrate A	L	L	W	L	G	R	V	V	G	G	P	V
substrate B	K	K	W	L	G	R	V	P	G	G	P	V

Protease variants were screened on substrate B at complexities of  $10^6$  variants by confocal fluorescence spectroscopy. The substrate was a peptide biotinylated at the N-terminus and fluorescently labeled at the C-terminus. After incubation of the peptide with supernatant of cells expressing different variants of the protease, streptavidin is added and the samples are analysed by confocal fluorimetry. The low concentration of the peptide (20nM) leads to a preferential cleavage by proteases with a high  $k_{cat}/K_M$  value, i.e. proteases with high specificity towards the target sequence.

Variants selected in the screening procedure were further evaluated for their specificity towards substrate B and closely related substrate A by measuring time courses of the proteolytic digestion and determining the rate constants which are proportional to the  $k_{cat}/K_M$  values. Clearly, compared to the human trypsin that was used as scaffold protein, the specific activity of variants 1 and 2 is shifted (SEQ ID NOs: 2 and 3, respectively) towards substrate B. Variant 3 (SEQ ID NO:4), on the other hand, serves as a negative control with similar activities as the human trypsin I. Sequencing of the genes of the three variants revealed the following amino acid sequences in the SDRs.

**Table 2:** Sequences of the two SDRs in three different variants selected for specific hydrolysis of substrate B (SEQ ID NOs:78-83).

	SDR 1						SDR 2					
Trypsin	-	-	-	-	-	-	-	-	-	-	-	-
Variant 1	D	A	V	G	R	D	T	I	T	N	S	
Variant 2	N	G	R	D	L	E	V	R	G	T	W	
Variant 3	G	F	V	M	F	N	R	S	P	L	T	

In a further experiment a pool of variants containing different numbers of SDRs per gene were screened for increased specificity using a mixture of the defined substrate and pepton as a competing substrate. Variants containing one or two

SDRs per gene have been analyzed further. As a measure for the specificity the activity in the peptide cleavage assay was compared with and without the presence of the competing substrate. The concentration of the competing substrate was 10mg/ml. Under these conditions, unspecific proteases show, compared to specific proteases, a stronger decrease in activity with increasing competitor concentrations (range between 0 and 100mg/ml). The ratio of proteolytic activity with and without substrate is a quantitative measure for the specificity of the proteases. Figure 9 shows the relative activities with and without competing substrate. Human trypsin I that was used as the scaffold protein and two variants, one containing only SDR2, and one containing both SDRs, were compared. The specificity of the variant with both SDRs is by a factor of 2.5 higher than that of the variant with SDR2 only, confirming that there is a direct relation between the number of SDRs and the quantitative specificity of resulting engineered proteolytic enzymes.

Example V: Generation of an engineered proteolytic enzyme that specifically inactivates human TNF-alpha

Human trypsin alpha I or a derivative comprising one or more of the following amino acid substitutions E56G; R78W; Y131F; A146T; C183R was used as protein scaffold for the generation of an engineered proteolytic enzyme with high specificity towards human TNF-alpha. The identification of SDR sites in human trypsin I or derivatives thereof was done as described above. Two insertion sites within the scaffold were chosen for SDRs. The protease variants containing two inserts with different sequences and also the human trypsin I itself with no inserts were expressed in a *Bacillus subtilis* cells. The variant protease cells were separated to single cell clones and the protease expressing variants were screened for proteolytic activity on peptides with the desired target sequence of TNF-alpha. The activity of the protease variants was determined as the cleavage rate of a peptide with the desired target sequence of TNF-alpha in the absence and presence of competitor substrate. The specificity is expressed as the ratio of cleavage rates in the presence and absence of competitor (Fig. 14).

Table 3: Relative specificity of variants of engineered proteolytic enzymes with different SDR sequences in absence and presence of competitor substrate (SEQ ID NOs:84-95).

	k with comp / k without comp	Seq. of SDR 1	Seq. of SDR 2
--	---------------------------------	------------------	------------------

scaffold (no SDRs)	0.092	---	----
variant a	0.130	RPWDPS	VHPTS
variant b	0.187	GFVMFN	RSPLT
variant c	0.235	EIANRE	RGART
variant d	0.310	KAVVGT	RTPIS
variant e	0.374	VNIMAA	TTARK
variant f	0.487	AAFNGD	RKDFW

The antagonistic effect of three inventive protease variants on human TNF- $\alpha$  is shown in Figure 15. By the use of the variants, the induction of apoptosis is almost completely eliminated indicating the anti-inflammatory efficacy of the inventive proteases to initiate TNF- $\alpha$  break down. TNF- $\alpha$  has been incubated with concentrated supernatant from cultures expressing the variants i to iii for 2 hours. The resulting TNF- $\alpha$  has been incubated with non-modified cells for 4 hours. The effect of the remaining TNF- $\alpha$  activity was determined as the extent of apoptosis induction by detection of activated caspase-3 as marker for apoptotic cells. For the controls either no protease was added with the human TNF- $\alpha$  (dead cells) or buffer instead of human TNF- $\alpha$  (live cells) was used, respectively. An analogous experiment is shown in Figure 16 using purified variant xiii. TNF- $\alpha$  was incubated with different concentrations of the purified inventive protease variant.

To demonstrate the specificity of the inventive protease variants, proteins from human blood serum or purified human TNF- $\alpha$  have been incubated with human trypsin I or the inventive engineered proteolytic enzyme variants, respectively. Here, variant x corresponds to Seq ID No: 75 comprising the same SDRs as variant f, i.e. SDRs according to Seq ID No. 89 (SDR1) and 95 (SDR2). Variants xi and xii correspond to derivatives thereof comprising the same SDR sequences. Remaining intact protein was determined as a function of time. While the variants as well as human trypsin I digest human TNF- $\alpha$ , only trypsin shows activity on serum protein (Figure 17 a and b). This demonstrates the high TNF- $\alpha$  specificity of the inventive proteolytic enzymes and indicates their safety and accordingly their low side effects for therapeutic use.

Example VI: Generation of an engineered proteolytic enzyme that specifically hydrolysis human VEGF.

Human trypsin I was used as protein scaffold for the generation of an engineered proteolytic enzyme with high specificity towards human VEGF. The identification of SDR sites in human trypsin I was done as described above. Two insertion sites within the scaffold were chosen for SDRs. The protease variants containing two inserts with different sequences were expressed in *Bacillus subtilis* cells. The variant protease cells were separated to single cell clones and the protease expressing variants were screened as described above. The activity of the protease variants was determined as the rate of VEGF cleavage. 4 µg of recombinant human VEGF<sub>165</sub> was incubated with 0.18 µg of purified protease in PBS / pH 7.4 at room temperature. Aliquots were taken at the indicated time points and analysed on a polyacrylamide gel. The extent of cleavage was quantified by densitometric analysis of the bands. The activity is plotted over incubation time in Figure 18. Specific cleavage was controlled by further SDS polyacrylamide gel analyses.

## Claims

1. Use of a protease with defined specificity for a target substrate for preparing a medicament for the treatment of a specific disease related to said target substrate.

2. The use according to claim 1, wherein

(I) the protease hydrolyzes the target substrate and thereby

- (i) eliminates or reduces one or more biological activities or physico-chemical properties or pharmacological properties of the target protein, and/or
- (ii) activates or increases one or more biological activities or physico-chemical properties or pharmacological properties of the target protein, and/or
- (iii) adds one or more biological activities or physico-chemical properties or pharmacological properties to the target protein;

and/or

(II) the target substrate hydrolyzed by the protease is a soluble protein, in particular a cytokine, such as the TNF-superfamily proteins, interleukines, interferons, chemokines and growth factors; a hormone; a toxin; an enzyme, such as oxidoreductases, transferases, hydrolases, lyases, isomerases and ligases; a structural protein, such as collagens; or an immunoglobulin; or a membrane associated protein, in particular a single pass transmembrane protein; a multipass transmembrane protein, such as G-protein coupled receptors, ion channels and transporters; a lipid-anchored membrane protein or a GPI-anchored membrane protein.

3. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a TNF-superfamily protein.

4. The use according to claim 3, wherein

(i) the protease is capable of hydrolysing peptide bonds in human tumor necrosis factor-alpha (hTNF- $\alpha$ , SEQ ID NO:96) or related molecules of the same structure class, preferably the peptide bonds between positions 31/32, 32/33, 44/45, 45/46, 87/88, 128/129, 130/131, 140/141 and/or 141/142, most preferably

between positions 31/32, 32/33 and/or 45/46 of hTNF- $\alpha$  or between analogous positions in said related molecules; and/or

(i) the medicament is suitable for the treatment of rheumatoid arthritis, inflammatory bowel diseases, psoriasis, Crohn's disease, Ulcerative colitis, diabetes type II, classical Hodgkin's Lymphoma (cHL), Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, systemic lupus erythematosus, multiple sclerosis, Systemic inflammatory response syndrome (SIRS), multiple organ dysfunction syndrome (MODS), eosinophilia, neurodegenerative disease, stroke, closed head injury, encephalitis, CNS disorders, asthma, rheumatoid arthritis, sepsis, vasodilation, intravascular coagulation, multiple organ failure or other diseases connected with hTNF- $\alpha$ .

5. The use according to claim 4, wherein the protease is derived from a serine protease of the structural class S1, and preferably wherein the protease is derived from human trypsin I (SEQ ID NO:1).

6. The use according to claim 4, wherein the protease has the a sequence shown in SEQ ID NO:74 or 75 or a derivative thereof and is capable of hydrolysing hTNF- $\alpha$  at positions 31/32 and/or 32/33.

7. The use according to claim 3, wherein

(i) the protease is capable of hydrolysing peptide bonds in human Tumor necrosis factor ligand superfamily member 5 (CD40-L) or related molecules of the same structural class, preferably the peptide bonds between positions 117/118, 133/134, 145/146, 165/166, 200/201, 201/202, 207/208, 216/217 and/or 243/244, most preferred between positions 133/134, 165/166, 201/202 and/or 216/217 of CD40-L (SEQ ID NO:143), or between analogous positions in related molecules; and/or

(ii) the medicament is suitable for the treatment of systemic lupus erythematosus, classical Hodgkin's Lymphoma (cHL) or other diseases connected with CD40-L.

8. The use according to claim 3, wherein

(i) the engineered protease is capable of hydrolysing peptide bonds in human Macrophage migration inhibitory factor (hMIF) or related molecules of the same



structural class, preferably the peptide bonds between positions 16/17, 44/45, 66/67, 73/74, 77/78, 88/89, 92/93 and/or 100/101, most preferred between positions 16/17 and/or 92/93 of hMIF (SEQ ID NO:109) or between analogous positions in related molecules; and/or

(ii) the medicament is suitable for the treatment of inflammatory diseases, or other diseases connected with hMIF.

9. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is an interleukine.

10. The use according to claim 9, wherein

(i) the protease is capable of hydrolysing peptide bonds in Interleukin-1 beta (IL-1 $\beta$ ) or related molecules of the same structural class, preferably the peptide bonds between positions 24/25, 35/36, 46/47, 54/55, 74/75, 75/76, 76/77, 77/78, 86/87, 88/89, 93/94, 94/95, 97/98 and/or 150/151, most preferred between positions 35/36, 75/76, 76/77, 88/89, 93/94, 94/95 and/or 150/151 of IL-1 $\beta$  (SEQ ID NO:112), or between analogous positions in related molecules; and/or

(iii) the medicament is suitable for the treatment of diabetes, brain inflammation in cancer, arthritis, autoimmune and inflammatory diseases or other diseases connected with IL-1 $\beta$ .

11. The use according to claim 9, wherein

(i) the protease is capable of hydrolysing peptide bonds in human Interleukin 2 (hIL-2) or related molecules of the same structural class, preferably the peptide bonds between positions 20/21, 32/33, 38/39, 43/44, 45/46, 48/49, 49/50, 54/55, 64/65, 76/77, 83/84, 84/85, 107/108, 109/110 and/or 120/121, most preferred between positions 109/110 of hIL-2 (SEQ ID NO:99) or between analogous positions in related molecules; and/or

(ii) the medicament is suitable for the treatment of T-cell leukemia, hairy cell leukemia, Crohn's disease, Ulcerative colitis, Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, systemic lupus erythematosus, multiple sclerosis, asthma and chronic obstructive pulmonary, classical Hodgkin's Lymphoma (cHL) or other diseases connected with hIL-2.

12. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin 3 (hIL-3) or related molecules of the same structural class, preferably peptide bonds between positions 21/22, 28/29, 36/37, 44/45, 46/47, 51/52, 63/64, 66/67, 79/80, 94/95, 101/102, 108/109 and/or 109/110, most preferred between positions 21/22, 28/29, 46/47, 63/64, 66/67, 79/80 and/or 101/102 of hIL-3 (SEQ ID NO:148), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), eosinophilia or other diseases connected with hIL-3.

13. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin 4 (hIL-4) or related molecules of the same structural class, preferably the peptide bonds between positions 4/5, 12/13, 31/32, 37/38, 61/62, 62/63, 64/65, 91/92, 102/103, 121/122 and/or 126/127, most preferred between positions 4/5, 61/62, 62/63, 64/65 and/or 121/122 of hIL-4 (SEQ ID NO:118), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, Asthma, chronic obstructive pulmonary disease, allergic inflammatory reactions or other diseases connected with hIL-4.

14. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin-5 (hIL-5) or related molecules of the same structural class, preferably the peptide bonds between positions 12/13, 32/33, 67/68, 76/77, 77/78, 80/81, 83/84, 84/85, 85/86, 90/91, 91/92, 92/93 and/or 98/99, most preferred between positions 90/91, 91/92, 92/93 and/or 98/99 of hIL-5 (SEQ ID NO:133), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), asthma, chronic obstructive pulmonary disease,

eosinophilia, allergic inflammatory diseases or other diseases connected with hIL-5.

15. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin-6 (hIL-6) or related molecules of the same structural class, preferably the peptide bonds between positions 32/33, 35/36, 55/56, 71/72, 129/130, 130/131, 132/133, 135/136, 141/142, 161/162, 180/181 and/or 183/184, most preferred between positions 135/136 and/or 141/142 of hIL-6 (SEQ ID NO:134), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), breast cancer, renal cell carcinoma, multiple myeloma, lymphoma, leukemia, Grave's disease, Hashimoto's thyroiditis, Sjogren's syndrome, systemic lupus erythematosus, Systemic inflammatory response syndrome (SIRS) which leads to distant organ damage and multiple organ dysfunction syndrome (MODS), chronic obstructive pulmonary disease (COPD), Castleman's diseases, inflammatory bowel diseases, Crohn's disease or other diseases connected with hIL-6.

16. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin 8 (hIL-8) or related molecules of the same structural class, preferably the peptide bonds between positions 11/12, 15/16, 45/46, 47/48, 52/53, 54/55, 60/61, 64/65 and/or 67/68, most preferred between positions 45/46 of hIL-8 (SEQ ID NO:100), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of Crohn's disease, Ulcerative colitis, classical Hodgkin's Lymphoma (cHL), Systemic inflammatory response syndrome (SIRS) which leads to distant organ damage and multiple organ dysfunction syndrome (MODS), chronic obstructive pulmonary disease (COPD), endometriosis, psoriasis, atherosclerotic lesions or other diseases connected with hIL-8.

17. The use according claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin-10 (hIL-10) or related molecules of the same structural class, preferably the peptide bonds between positions 24/25, 25/26, 27/28, 28/29, 40/41, 44/45, 49/50, 57/58, 59/60, 84/85, 86/87, 106/107, 107/108, 110/111, 130/131, 134/135, 137/138, 138/139 and/or 144/145, most preferred between positions 24/25, 27/28, 44/45, 49/50, 86/87, 137/138 and/or 144/145 of hIL-10 (SEQ ID NO:135), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), diseases related to the suppression of cytotoxic T-cells or other diseases connected with hIL-10.

18. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin 12 beta chain (hIL-12 $\beta$ ) or related molecules of the same structural class, preferably the peptide bonds between positions 14/15, 18/19, 29/30, 34/35, 87/88, 99/100, 102/103, 104/105, 161/162, 174/175, 222/223, 225/226, 228/229, 238/239, 268/269 and/or 293/294, most preferred between positions 18/19, 34/35, 87/88 and/or 161/162 of hIL-12 $\beta$  (SEQ ID NO:97) or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of Crohn's disease, classical Hodgkin's Lymphoma (cHL) or other diseases connected with hIL-12 $\beta$ .

19. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin 13 (hIL-13) or related molecules of the same structural class, preferably the peptide bonds between positions 25/26, 62/63, 65/66, 86/87, 87/88, 98/99, 108/109 and/or 111/112, most preferred between positions 87/88 of hIL-13 (SEQ ID NO:119), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of cancer, classical Hodgkin's Lymphoma (cHL), eosinophilia, asthma, chronic obstructive pulmonary disease, fibrosis, psoriasis, atopic dermatitis, Ulcerative colitis or other diseases connected with hIL-13.

20. The use according to claim 9, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interleukin 18 (hIL-18) or related molecules of the same structural class, preferably the peptide bonds between positions 17/18, 32/33, 37/38, 39/40, 40/41, 53/54, 58/59, 79/80, 90/91, 93/94, 98/99, 110/111, 120/121, 123/124, 131/132, 132/133, 142/143, 147/148 and/or 157/158, most preferred between positions 37/38, 132/133, 142/143 and/or 157/158 of hIL-18 (SEQ ID NO:98) or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of Crohn's disease, inflammation liver injuries, pulmonary tuberculosis, plural tuberculosis, rheumatoid arthritis or other diseases connected with hIL-18.

21. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is an interferone.

22. The use according to claim 21, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interferon-gamma (hIFN-?) or related molecules of the same structural class, preferably the peptide bonds between positions 2/3, 6/7, 13/14, 21/22, 24/25, 34/35, 36/37, 37/38, 62/63, 68/69, 83/84, 86/87, 90/91, 102/103, 107/108 and/or 108/109, most preferred between positions 13/14, 24/25, 37/38, 62/63, 68/69, 102/103 and/or 107/108 of hIFN-? (SEQ ID NO:137), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), Crohn's disease, type I diabetes or other diseases connected with IFN-?.

23. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a chemokine.

24. The use according to claim 23, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human small inducible cytokine A2 (hCCL2) or related molecules of the same structural class, preferably the peptide bonds between positions 3/4, 13/14, 18/19, 19/20, 24/25, 29/30, 38/39, 54/55, 56/57, 58/59, 62/63, 65/66 and/or 68/69, most preferred between positions 19/20, 29/30, 38/39, 54/55 and/or 62/63 of hCCL2 (SEQ ID NO:102), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of Crohn's disease, Ulcerative colitis, or other diseases connected with hCCL2.

25. The use according to claim 23, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Eotaxin (hCCL11) or related molecules of the same structural class, preferably the peptide bonds between positions 11/12, 16/17, 17/18, 22/23, 27/28, 33/34, 44/45, 47/48, 48/49, 52/53, 54/55, 56/57, 60/61, 66/67, and/or 73/74, most preferred between positions 48/49 and/or 66/67 of hCCL11 (SEQ ID NO:101), or between analogous positions in related molecules; and/or
- (ii) wherein the medicaments is suitable for the treatment of Crohn's disease and Ulcerative colitis, classical Hodgkin's Lymphoma (cHL), chronic pathophysiologic dysfunction, characterized by an influx mainly of Th2 cells, eosinophilia or other diseases connected with hCCL11.

26. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a growth factor.

27. The use according to claim 26, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Vascular endothelial growth factor (hVEGF) or related molecules of

the same structural class, preferably the peptide bonds between positions 16/17, 19/20, 23/24, 34/35, 41/42, 56/57, 62/63, 63/64, 64/65, 65/66, 82/83, and/or 84/85, most preferred between positions 23/24, 41/42, 63/64, 82/83 and/or 84/85 of hVEGF (SEQ ID NO:103), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of all solid tumors and metastatic solid tumors, inflammatory breast cancer or other diseases connected with hVEGF.

28. The use according to claim 26, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Transforming growth factor beta 1 (hTGF- $\beta$ 1) or related molecules of the same structural class, preferably the peptide bonds between positions 23/24, 25/26, 26/27, 27/28, 37/38, 55/56 and/or 94/95, most preferred between positions 25/26, 55/56 and/or 94/95 of hTGF- $\beta$ 1 (SEQ ID NO:104), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of a variety of cancers including breast cancer, colorectal cancer and classical Hodgkin's Lymphoma (cHL); fibrosis, suppression of cell-mediated immunity, glaucoma, diffuse systemic sclerosis or other diseases connected with hTGF- $\beta$ 1.

29. The use according to claim 26, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Somatotropin (hGrowth hormone; hGH) or related molecules of the same structural class, preferably the peptide bonds between positions 8/9, 16/17, 19/20, 26/27, 33/34, 38/39, 41/42, 70/71, 77/78, 94/95, 103/104, 112/113, 115/116, 116/117, 130/131, 147/148, 154/155 and/or 178/179, most preferred between positions 112/113, 147/148 and/or 154/155 of GH (SEQ ID NO:121), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of acromegaly, diabetes and diabetic kidney disease including renal hypertrophy and glomerular enlargement, cardiovascular disorders or other diseases connected with hGH.

30. The use according to claim 26, wherein

- (i) the protease is capable of hydrolysing peptide bonds in Insulin-like growth factor II (hIGF-II) or related molecules of the same structural class, preferably the peptide bonds between positions 15/16, 23/24, 24/25, 34/35, 37/38, 38/39, 48/49 and/or 49/50, most preferred between positions 23/24 of hIGF-II (SEQ ID NO:122), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of diabetes, diabetic kidney disease or other diseases connected with hIGF-II.

31. The use according to claim 26, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Hepatocyte growth factor (hHGF) or related molecules of the same structural class, preferably the peptide bonds between positions 54/55, 60/61, 62/63, 63/64, 68/69, 76/77, 112/113, 123/124, 134/135, 168/169, 198/199 and/or 202/203, most preferred between positions 63/64, 68/69, 76/77, 168/169 and/or 202/203 of hHGF (SEQ ID NO:120), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of angiogenic disorders, hepatocellular carcinoma or other diseases connected with hHGF.

32. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is soluble hormone.

33. The use according to claim 32, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Insulin (hInsulin) or related molecules of the same structural class,



- preferably the peptide bonds between positions 16/17 and/or 22/23 of hInsulin B chain (SEQ ID NO:105), and/or between positions 14/15 of Insulin A chain (SEQ ID NO:106) or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of insulin overdose or other diseases connected with hInsulin.

34. The use according to claim 32, wherein

- (i) the engineered protease is capable of hydrolysing peptide bonds in human Ghrelin (Ghrelin) or related molecules of the same structural class, preferably the peptide bonds between positions 1/2, 2/3, 3/4 and/or 4/5 of hGhrelin (SEQ ID NO:107), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of obesity or other diseases connected with hGhrelin.

35. The use according to claim 32, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human angiotensinogen (angiotensin) or related molecules of the same structural class, preferably the peptide bonds between positions 1/2, 3/4, and/or 7/8, most preferred between positions 3/4 of angiotensin (SEQ ID NO:108), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of essential hypertension and other diseases connected with angiotensin.

36. The use according to claim 32, wherein

- (i) the engineered protease is capable of hydrolysing peptide bonds in human leptin (leptin) or related molecules of the same structural class, preferably the peptide bonds between positions 8/9, 9/10, 15/16, 23/24, 40/41, 53/54, 71/72, 85/86, 94/95, 108/109 and/or 141/142, most preferred between positions 9/10, 40/41, 71/72, 94/95 and/or 108/109 of leptin (SEQ ID NO:127), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of obesity or other diseases connected with leptin.

37. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a binary toxin.

38. The use according to claim 37, wherein

- (i) the engineered protease is capable of hydrolysing peptide bonds in Protective antigen precursor (PA-83) or related molecules of the same structural class, preferably the peptide bonds between positions 72/73, 73/74, 92/93, 93/94, 131/132, 149/150, 178/179, 213/214, 214/215, 387/388, 425/426, 426/427, 427/428, 453/454, 520/521, 608/609, 617/618, 671/672, 679/680, 680/681, 683/684 and/or 684/685, most preferred between positions 72/73, 73/74, 93/94, 149/150, 387/388, 425/426, 427/428 and/or 683/684 of hPA-83 (SEQ ID NO:123), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of anthrax infection or other diseases connected with PA-83.

39. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a protease, in particular a serine, cysteine or metallo protease.

40. The use according to claim 39, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human plasminogen (plasminogen) or related molecules of the same structural class, preferably the peptide bonds between position 580/581 of plasminogen (SEQ ID NO:140), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of thrombosis or other diseases connected with plasminogen.

41. The use according to claim 39, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Prothrombin (thrombin) or related molecules of the same structural

- class, preferably the peptide bonds between positions 198/199, 327/328, 363/364, most preferred between positions 327/328 and/or 363/364 of thrombin (SEQ ID NO:149), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of bleeding or other diseases connected with thrombin.

42. The use according to claim 39, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human beta-secretase or related molecules of the same structural class, preferably the peptide bonds between positions 61/62, 64/65, 130/131, 131/132, 159/160, 216/217, 238/239, 239/240, 246/247, 256/257, 259/260, 330/331, 365/366, 378/379 and/or 381/382, most preferred between positions 61/62, 131/132, 246/247, 259/260, 365/366 and/or 378/379 of beta-secretase (SEQ ID NO:139), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of Alzheimer or other diseases connected with beta-secretase precursor.

43. The use according to claim 39, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human matrix metalloproteinase-2 (hMMP-2) or related molecules of the same structural class, preferably the peptide bonds between positions 62/63, 68/69, 75/76, 76/77, 79/80, 88/89, 110/111, 112/113, 115/116, 120/121, 164/165, 254/255, 267/268, 296/297, 324/325, 325/326, 382/383, 383/384, 470/471, 500/501, 550/551, 564/565, 595/596, 597/598, 608/609, 646/647, 649/650 and/or 650/651, most preferred between positions 68/69, 115/116, 120/121, 164/165, 325/326, 383/384, 470/471, 500/501, 595/596, 608/609 and/or 650/651 of hMMP-2 (SEQ ID NO:131), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of a variety of cancers including bladder cancer, breast tumor cancer, gastric cancer, lung cancer or other diseases connected with hMMP-2.

44. The use according to claim 39, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human matrix metalloproteinase-9 (hMMP-9) or related molecules of the same structural class, preferably the peptide bonds between positions 41/42, 42/43, 106/107, 113/114, 134/135, 160/161, 162/163, 163/164, 222/223, 226/227, 265/266, 266/267, 267/268, 284/285, 309/310, 321/322, 322/323, 324/325, 356/357, 380/381, 433/434 and/or 440/441, most preferred between positions 160/161, 163/164, 226/227, 284/285, 321/322, 322/323 and/or 433/434 of hMMP-9 (SEQ ID NO:132), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of a variety of cancers including bladder cancer, breast tumor cancer, gastric cancer, lung cancer or other diseases connected with hMMP-9.

45. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a type-1 single pass transmembrane protein.

46. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in HIV membrane glycoprotein (GP120) or related molecules of the same structural class, preferably the peptide bonds between positions 97/98, 99/100, 107/108, 113/114, 117/118, 227/228, 231/233, 279/280, 335/336, 337/338, 368/369, 412/413, 419/420, 429/430, 444/445, 457/458, 474/475, 476/477, 477/478, 485/486 and/or 490/491, most preferred between positions 99/100, 368/369, 412/413, 419/420, 444/445 and/or 490/491 of GP120 (SEQ ID NO:124) or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of AIDS or HIV infection or other diseases connected with GP120.

47. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds human Cytotoxic T-lymphocyte protein 4 (CTLA-4) or related molecules of the same structural class, preferably the peptide bonds between positions 14/15, 28/29, 33/34, 38/39, 41/42, 62/63, 72/73, 85/86, 95/96, 100/101, 105/106, 119/120, 125/126 and/or 127/128, most preferred between positions 14/15, 28/29, 38/39, 41/42, 62/63 and/or 85/86 of CTLA-4 (SEQ ID NO:144), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of breast cancer or other diseases connected with CTLA-4.

48. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Integrin alpha-2 (hVLA-2) or related molecules of the same structural class, preferably the peptide bonds between positions 160/161, 174/175, 201/202, 219/220, 231/232, 232/233, 233/234, 243/244, 259/260, 264/265, 268/269, 288/289, 292/293, 294/295, 298/299, 301/302, 310/311 and/or 317/318, most preferred between positions 160/161, 174/175, 201/202, 219/220, 243/244, 264/265, 292/293 and/or 294/295 of hVLA-2 (SEQ ID NO:147), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of renal tumors, uveal melanomas, gastrointestinal tumors or other diseases connected with hVLA-2.

49. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Vascular endothelial growth factor receptor 1 (hVEGFR 1) or related molecules of the same structural class, preferably the peptide bonds between positions 175/176, 180/181, 187/188, 189/190, 190/191, 224/225 and/or 331/332, most preferred between positions 189/190 and/or 331/332 of hVEGFR 1 (SEQ ID NO:114), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of solid tumors and metastatic solid tumors, astrocytic brain tumors, pancreatic cancer, metastatic renal cancer or other diseases connected with hVEGFR 1.

50. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Vascular endothelial growth factor receptor 2 (hVEGFR 2) or related molecules of the same structural class, preferably the peptide bonds between positions 214/215, and/or 323/324, most preferred between positions 214/215 of hVEGFR 2 (SEQ ID NO:115), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of solid tumors and metastatic solid tumors pancreatic cancer, metastatic renal cancer, metastatic CRC, or other diseases connected with hVEGFR 2.

51. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Epidermal growth factor receptor (hEGFr) or related molecules of the same structural class, preferably the peptide bonds between positions 20/21, 29/30, 48/49, 74/75, 155/156, 165/166, 167/168, 202/203, 206/207, 220/221, 223/224, 246/247, 251/252, 254/255, 269/270, 270/271, 297/298, 304/305, 305/306, 357/358, 364/365, 369/370, 430/431, 443/444, 454/455, 455/456, 463/464, 465/466, 476/477, 507/508 and/or 509/510, most preferred between positions 155/156, 246/247, 251/252, 254/255, 269/270, 270/271, 297/298, 304/305, 306/307, 364/365 and/or 454/455 of hEGFr (SEQ ID NO:116), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of bladder cancer, breast cancer, cervical cancer, colorectal cancer, endometrial cancer, oesophageal cancer, head and neck cancer, gastric cancer, non-small-cell lung carcinoma, ovarian cancer or other diseases connected with hEGFr.

52. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Epithelial cell adhesion molecule (Ep-CAM) or related molecules of the same structural class, preferably the peptide bonds between positions 14/15, 19/20, 25/26, 29/30, 30/31, 33/34, 44/45, 55/56, 67/68, 70/71, 90/91 and/or 100/101, most preferred between positions 14/15, 30/31, 44/45, 70/71 and/or 100/101 of Ep-CAM (SEQ ID NO:125) or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of colorectal cancer or other diseases connected with Ep-CAM.

53. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Insulin-like growth factor I receptor (hIGF-1r) or related molecules of the same structural class, preferably the peptide bonds between positions 39/40, 59/60, 115/116, 132/133, 146/147, 171/172, 191/192, 250/251, 262/263, 270/271, 290/291, 306/307, 307/308, 335/336, 336/337, 403/404, 405/406, 455/456 and/or 470/471, most preferred between positions 39/40, 262/263, 306/307, 307/308, 335/336, 405/406 and/or 470/471 of hIGF-1r (SEQ ID NO:126) or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of a variety of cancers including breast cancer or other diseases connected with hIGF-1r.

54. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human T-cell surface antigen CD2 (hCD2) or related molecules of the same structural class, preferably the peptide bonds between positions 16/17, 20/21, 28/29, 29/30, 40/41, 42/43, 43/44, 48/49, 49/50, 51/52, 54/55, 58/59, 63/64, 69/70, 76/77, 89/90 and/or 91/92, most preferred between positions 28/29, 40/41, 43/44, 51/52, 76/77 and/or 89/90 of hCD2 (SEQ ID NO:128) or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of psoriasis or other diseases connected with hCD2.

55. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human T-cell surface glycoprotein CD4 (hCD4) or related molecules of the same structural class, preferably the peptide bonds between positions 53/54, 63/64, 88/89, 166/167, 167/168, 173/174, 206/207, 219/220, 224/225, 226/227, 230/231, 244/245, 251/252, 252/253, 322/323, 329/330 and/or 334/335, most preferred between positions 88/89, 173/174, 206/207, 219/220, 230/231, 251/252 and/or 252/253 of hCD4 (SEQ ID NO:129), or between analogous positions in related molecules;
- (ii) preferably the medicament is suitable for the treatment of psoriasis, transplant rejection, graft-versus-host colitis, autoimmune disorders, rheumatoid arthritis or other diseases connected with hCD4.

56. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Integrin alpha-L (hCD11a) or related molecules of the same structural class, preferably the peptide bonds between positions 145/146, 152/153, 156/157, 159/160, 160/161, 177/178, 178/179, 189/190, 190/191, 191/192, 193/194, 197/198, 200/201, 221/222, 229/230, 249/250, 253/254, 268/269, 290/291, 297/298, 304/305 and/or 305/306, most preferred between positions 145/146, 159/160, 160/161, 189/190, 229/230, 249/250, 268/269, 297/298, 304/305 and/or 305/306 of hCD11a (SEQ ID NO:130), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of psoriasis or other diseases connected with hCD11a.

57. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Interferon-gamma receptor alpha chain (hIFN- $\gamma$ -R1) or related



molecules of the same structural class, preferably the peptide bonds between positions 49/50, 52/53, 58/59, 62/63, 72/73, 76/77, 106/107, 107/108, 116/117, 122/123, 174/175, 176/177, 179/180, 215/216 and/or 222/223, most preferred between positions 49/50, 72/73, 116/117, 122/123, 174/175, 176/177 and/or 215/216 of hIFN- $\gamma$ -R1 (SEQ ID NO:136), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of classical Hodgkin's Lymphoma (cHL), type I diabetes or other diseases connected with hIFN- $\gamma$ -R1.

58. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Platelet membrane glycoprotein IIb/IIIa (GPIIb/IIIa) or related molecules of the same structural class, preferably the peptide bonds between positions 67/68, 76/77, 91/92, 129/130, 143/144, 144/145, 179/180, 181/182, 198/199, 208/209, 209/210, 216/217, 239/240, 261/262, 410/411, 434/435, 532/533, 556/557, 557/558, 596/597, 597/598, 621/622, 650/651, 651/652, 661/662, 662/663 and/or 689/690, most preferred between positions 67/68, 76/77, 179/180, 261/262, 410/411, 434/435, 650/651, 662/663 and/or 689/690 of GPIIb/IIIa (SEQ ID NO:141), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of unstable angina, carotid stenting, ischemic stroke, peripheral vascular diseases, angiogenesis-related diseases, disseminating tumors or other diseases connected with GPIIb/IIIa.

59. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Intercellular adhesion molecule-1 (ICAM-1) or related molecules of the same structural class, preferably the peptide bonds between positions 26/27, 40/41, 60/61, 71/72, 88/89, 97/98, 102/103, 128/129, 131/132, 132/133, 149/150, 150/151, 151/152, 160/161, 164/165 and/or 166/167, most preferred between positions 71/72,

88/89, 102/103, 150/151, 151/152, 160/161 and/or 166/167 of ICAM-1 (SEQ ID NO:142), or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of Crohn's disease or other diseases connected with ICAM-1.

60. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds human TGF-beta receptor type II (hTGF- $\beta$  RII) or related molecules of the same structural class, preferably the peptide bonds between positions 32/33, 34/35, 35/36, 66/67, 67/68, 69/70, 82/83, 103/104, 104/105, 105/106, 118/119, 122/123 and/or 130/131, most preferred between positions 32/33, 34/35, 66/67, 69/70, 104/105, 122/123 and/or 130/131 of hTGF- $\beta$  RII (SEQ ID NO:145), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of diffuse systemic sclerosis or other diseases connected with hTGF- $\beta$  RII.

61. The use according to claim 45, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Membrane cofactor protein (hMCP) or related molecules of the same structural class, preferably the peptide bonds between positions 15/16, 17/18, 25/26, 27/28, 31/32, 32/33, 35/36, 47/48, 48/49, 58/59, 67/68, 69/70, 70/71, 110/111, 119/120, 125/126 and/or 130/131, most preferred between positions 15/16, 32/33, 47/48, 48/49, 70/71, 119/120 and/or 125/126 of hMCP (SEQ ID NO:146), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of renal tumors, uveal melanomas, gastrointestinal tumors or other diseases connected with hMCP.

62. The use according to claim 1 or 2, wherein the target substrate hydrolyzed by the protease is a protease-activated receptor or hydroxytryptamine receptor.

63. The use according to claim 62, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Protease activated receptor 1 (hPAR1) or related molecules of the same structural class, preferably the peptide bonds between positions 46/47, 50/51, 51/52, 52/53 and/or 58/59 of hPAR1 (SEQ ID NO:110), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of thrombosis or other diseases connected with hPAR1.

64. The use according to claim 62, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Protease activated receptor 2 (hPAR2) or related molecules of the same structural class, preferably the peptide bonds between positions 41/42, 44/45, 51/52, 59/60 and/or 62/63 of hPAR2 (SEQ ID NO:111), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of Crohn's disease, Ulcerative colitis and Inflammatory bowel disease, asthma, inflammation associated pain and arthritis or other diseases connected with hPAR2.

65. The use according to claim 62, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human Protease activated receptor 4 (hPAR4) or related molecules of the same structural class, preferably the peptide bonds between positions 57/58, 59/60, 68/69, 74/75 and/or 78/79 of hPAR4 (SEQ ID NO:113), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of thrombosis or other diseases connected with hPAR4.

66. The use according to claim 62, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human 5-hydroxytryptamine 1A receptor (h5-HT-1A) or related molecules of the same structural class, preferably the peptide bonds between

positions 101/102, 102/103, 181/182 and/or 370/371 of h5-HT-1A (SEQ ID NO:117) or between analogous positions in related molecules; and/or

- (ii) the medicament is suitable for the treatment of irritable bowel syndrome or other diseases connected with h5-HT-1A.

67. The use according to claim 1 or 2, wherein

- (i) the protease is capable of hydrolysing peptide bonds in human carcinoembryonic antigen (hCEA) or related molecules of the same structural class, preferably the peptide bonds between positions 17/18, 69/70, 71/72, 74/75, 77/78, 98/99, 116/117, 126/127 and/or 128/129 of hCEA (SEQ ID NO:138), or between analogous positions in related molecules; and/or
- (ii) the medicament is suitable for the treatment of colon cancer or other diseases connected with hCEA.

68. The use according to any one of claims 1 to 67, wherein the protease is an engineered protease, preferably an engineered protease characterized by a combination of the following components:

- (a) a protein scaffold capable to catalyze at least one chemical reaction on at least one target substrate and being derived from one or more proteins, and
- (b) one or more specificity determining regions (SDRs) located at sites in the protein scaffold that enable the resulting engineered protein to discriminate between at least one target substrate and one or more different substrates, and wherein the SDRs are essentially synthetic peptide sequences.

69. The use according to claim 68, wherein

- (I) the SDRs (b) have a length between one and 50 amino acid residues, preferably have a length between 2 and 20 amino acid residues, more preferably a length between 2 and 10 amino acid residues, even more preferably a length between 3 and 8 amino acid residues, and wherein the number of SDRs is at least one, preferably more than one, more preferably between two and eleven, most preferably between two and six; and/or
- (II) the protein scaffold (a) is comprised of one or more polypeptide segments being derived from same or different

- (i) proteins encoded by a gene of viral or prokaryotic or eukaryotic origin, and/or
- (ii) native enzymes, mutated variants or truncated derivatives thereof, and/or
- (iii) mammalian enzymes, preferably human enzymes.

70. The use according to claim 68 or 69, wherein the protein scaffold (a) is derived from a protease selected from the group consisting of aspartic, cysteine, serine, metallo and threonine proteases, even more preferably the protein scaffold (a) is derived from a serine protease of the structural class S1, S8, S11, S21, S26, S33 or S51, most preferably from class S1 or S8, a cysteine protease of the structure class C1, C2, C4, C10, C14, C19, C47, C48 or C56, most preferably from class C14, or an aspartic protease of the structural class A1, A2 or A26, most preferably from class A1, or a metalloprotease of the structural class M4 or M10.

71. The use according to any one of claims 68 to 70, wherein

- (i) the protein scaffold (a) is derived from a serine protease of the structural class S1; and/or
- (ii) the SDRs are located at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 18-25, 38-48, 54-63, 73-86, 122-130, 148-156, 165-171 and 194-204 in human trypsin I having the amino acid sequence shown in SEQ ID NO:1, and preferably at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 20-23, 41-45, 57-60, 76-83, 125-128, 150-153, 167-169 and 197-201 in human trypsin I.

72. The use according to claim 71, wherein

- (i) the protein scaffold (a) is derived from the serine protease trypsin, preferably human trypsin I having the amino acid sequence shown in SEQ ID NO:1, or the amino acid sequence SEQ ID NO:1 comprising one or more of the following amino acid substitutions E56G, R78W, Y131F, A146T and C183R; and
- (ii) at least one of two SDRs are located in the scaffold, a first SDR having a length of up to 6 amino acids and being inserted between

residues 42 and 43, and a second SDR having a length of up to 5 amino acids and being inserted between residues 123 and 124 (numbering relative to human trypsin having the amino acid sequence shown in SEQ ID NO: 1).

73. The use according to claim 72, wherein

- (i) one of the peptide sequences of the following group: SEQ ID NO: 72, 78, 79, 80, 84, 85, 86, 87, 88, and 89 is inserted as the first SDR between residues 42 and 43; and/or one of the peptide sequences of the following group: SEQ ID NO: 73, 81, 82, 83, 90, 91, 92, 93, 94, and 95 is inserted as the second SDR between residues 123 and 124; or
- (ii) the engineered enzyme comprises an amino acid sequence as shown in SEQ ID NO: 74, or SEQ ID NO: 75.

74. The use according to any one of claims 68 to 70, wherein

- (i) the protein scaffold (a) is derived from a serine protease of the structural class S8; and/or
- (ii) the SDRs are located at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 6-17, 25-29, 47-55, 59-69, 101-111, 117-125, 129-137, 139-154, 158-169, 185-195 and 204-225 in subtilisin E from *Bacillus subtilis* having the amino acid shown in SEQ ID NO: 7, and preferably at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 59-69, 101-111, 129-137, 158-169 and 204-225 in subtilisin E for *Bacillus subtilis*.

75. The use according to any one of claims 68 to 70, wherein

- (i) the protein scaffold (a) is derived from an aspartic protease of the structural class A1; and/or
- (ii) the SDRs are located at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 6-18, 49-55, 74-83, 91-97, 112-120, 126-137, 159-164, 184-194, 242-247, 262-267 and 277-300

in human pepsin having the amino acid sequence shown in SEQ ID NO:11, and more preferably at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 10-15, 75-80, 114-118, 130-134, 186-191 and 280-296 in human pepsin.

76. The use according to any one of claims 68 to 70, wherein

- (i) the protein scaffold (a) is derived from a cysteine protease of the structural class C14; and/or
- (ii) the SDRs are located at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 78-91, 144-160, 186-198, 226-243 and 271-291 in human caspase 7 having the amino acid sequence of SEQ ID NO:14, and preferably at one or more positions selected from the group of positions that correspond structurally or by amino acid sequence homology to the regions 80-86, 149-157, 190-194 and 233-238 of human caspase 7.

77. The use according to any one of claims 1 to 76, wherein the protease comprises

- (i) at least one further proteinaceous component, preferably being selected from the group consisting of binding domains, receptors, antibodies, regulation domains, pro-sequences, and fragments thereof, and/or
- (ii) at least one further functional component, preferably being selected from the group consisting of polyethylenglycols, carbohydrates, lipids, fatty acids, nucleic acids, metals, metal chelates, and fragments or derivatives thereof.

78. The use according to any one of claims 1 to 77, wherein the protease is obtainable by a method comprising at least the following steps:

- (a) providing a protein scaffold which catalyzes at least one chemical reaction on at least one target substrate,
- (b) generating a library of enzymes or isolated enzymes by combining the protein scaffold from step (a) with variants of one or more fully or partially random peptide sequences at sites in the protein scaffold that enable the

resulting enzyme to discriminate between at least one target substrate and one or more different substrates, and

(c) selecting out of the (library of) enzymes generated in step (b) one or more enzymes that have defined specificities towards at least one target substrate.

79. Use of an enzyme according to any one of claims 1 to 78 for *in vivo* or *in vitro* diagnostic purposes.

80. A pharmaceutical or diagnostic composition comprising one or more enzymes according to any one of claims 1 to 78, said pharmaceutical or diagnostic composition optionally comprising pharmaceutically or diagnostically acceptable carrier(s), excipient(s) and/or auxiliary agent(s).

81. A method for cleaving a target substrate as defined in claims 1 to 67 *in vivo* or *in vitro*, which comprises contacting the target substrate with a protease as defined in claims 1 to 78.

81. A method for treatment of a disease in a patient connected with a specific target substrate as defined in anyone of claims 1 to 67, which comprises administering the patient a suitable amount of a protease with defined specificity for said specific target substrate as defined in anyone of claims 1 to 78.



1/18




Fig. 1

2/18

Trypsin	IVGGYNCEENSVPYQVSL---NSGYHF-CGGSLINEQWVVSAGHCY----
a-Thrombin	IVEGSDAEIGMSPWQVMLFRKSPQELL-CGASLISDRWVLTAAHCLLYPP
Enteropeptidase	IVGGSNAKEGAWPWVVGGL---YYGGRLLCGASLVSSDWLVSAAHCVYGRN
	** * * * * ** ** * * * ----
Trypsin	-----KSRIQVRLGEH---NIEVLEGN-EQFINAAKIIRHPQYD-RKTL
a-Thrombin	WDKNFTENDLLVRIGKH---SRTRYERNIEKISMLEKIYIHPRYNWREN
Enteropeptidase	LE----PSKWTAILGLHMKSNLTSPQTV-PRLLD--EIVINPHYN-RRRK
	-1----- * * * * *
Trypsin	NNDIMLIKLSRAVINARVSTISLPTA----PPAT-----GTKCLISGWG
a-Thrombin	DRDIALMKLKKPVAFSDYIHPVCLPDR----ETAASLLQAGYKGRVTGWG
Enteropeptidase	DNDIAMMHLEFKVNYTDYIQPICLPEENQVFPP-----GRNC SIAGWG
	** * * * * -----2-----* ***
Trypsin	N-----TASSGADYPDELQCLDAPVLSQAKCEASYPG-KITSNMFCVGF
a-Thrombin	NLKETWTANVGKGQPSVLQVVNLPIVERPVCKDSTRI-RIITDNMFCAGYK
Enteropeptidase	T-----VVYQGTT-ANILQEADVPLLSNERCQQQMPEYNI TENMICAGYE
	--3-- * * * * *
Trypsin	-EGGK--DSCQGD SGGPVVCNGQ----LQ-----GVVSWGDGCAQKNKP
a-Thrombin	PDEGKRGDACEGDSGGPFVMKSP----FNNRWYQMGIVSWGEGCDRDGKY
Enteropeptidase	-EGGI--DSCQGD SGGPLMCQENNRWFLA-----GVTSFGYKCALPNRP
	* * * * * -----4-----* * * *
Trypsin	GVYTKVYNYVKWIKNTIAANS-
a-Thrombin	GFYTHVFRLLKKWIKVIDQFGE
Enteropeptidase	GVYARVSRFTEWIQSFLH----
	* * * * *

Fig. 2

3/18




Fig. 3

4/18

```

sub      IAHEYAQS--PY-----GISQ--IKAPALHSQGY-----
furin    VAKRRAKRD--VYQEPTDPKFPQQWYLSGVTQRDLNVKEAWAQGF-----
PC_SK1   EKERSKRSALRDS-----ALNL--FNDFPMWNQQWYLQDTRMTAALPKLDL
PC_SK5   NTHPCQ-----SD--MNIEGAWKRGY-----
          -----1-----2--

sub      -----TGSNVKVAIDSGIDSSHPL--NVRGGAS--FVPSETN-----P--
furin    -----TGHGIVVSILDDGIEKNHPDLAGNYDPGAS--FDVNDQD-----PDPQ
PC_SK1   HVIPVWQKGTGKGVVITVLDGLEWNHTDIYANYDPEASYDFNDNDHD-----P--
PC_SK5   -----TGKNIVVTILDDGIERTHPDL--MQNYDA--LASCDVNGNDLDPMP--
          -----* * * *-----3-----

sub      ----YQ-----DGSS--HGTHVAGTIA--AL--NNSIGVLGVSPSASLYAVKVLDs----
furin    PRYTQM-----NDNR--HGTRCAGEVA--AVANNGVCGVGVAYNARIGGVRLD-----
PC_SK1   ----FPRYDPTNENK--HGTRCAGEIAMQAN--NHKCGV--GVAYNSKVGIGIRMLDG----
PC_SK5   ----RY-----DASNENKHGTRCAGEVA--AAANNSHCTVGIAFNAKIGGVRLDGDVTD
          -----4-----* * * *-----

sub      -TSGSQYSWIINGIE-WAISNNMDVINMSLG-----GPT--GSTA-----LKT--
furin    -GEVTDAVEARS-LGLNPNHIIHYSASW-----GPEDDGKTVDGPARLAE--
PC_SK1   -IVTDAIEASSIGFN--PGHVDIYSASWGPNDGKTVEGP--GRLA-----QKAFE
PC_SK5   MVEAKSVSFNPQHVHIYSASWGPDDDGKTVD-----GPA--PLT-----RQ--
          -5-----6-----7-----8--

sub      --VVDKAVSSG-----IVVAAAAGNEGSS-----GSTSTVGYPKYPSTIAGVAV--
furin    --AFFRGVSQGRGGLGSIFVWASGNGGREHDSNCNCDGYTNSI--YTLSISSATQFQNV--
PC_SK1   YGVKQGRQKKG-----SIFVWASGNGGRQ-----GDNCD--GYTDSIYTISI--
PC_SK5   --AFENGVRMGRRGLGSVFVWASGNGGRSKDHCSNCDGYTNSI--YTISISSTAESGKKPWY
          -----8-----*-----9-----

sub      --N-----SSNQR-----ASFSSAG--SELDVMAPGVSIQSTLPGGTYGAY
furin    --PWYSEACSTLA-----TTYSSGNQNEKQIVTTDLRQKCT-----ESH
PC_SK1   --S-----SASQQGLSPWYAEKCSSTLATSYSYG--DYTDQRTSADLHNDCT-----ETH
PC_SK5   LEE-----CSSTL-----ATTYSYG--ESYDKKI--ITTLRQRTDNH
          -----10-----*-----11--

sub      NGTSMATPHVAGAAALIL--SKHP--TWTNAQVRDRLESTATY--LG--NSFYGKGLINV
furin    TGTSASAPLAAGIIALTLEANKNL--TWRDMQHLVVQTSKPAH--LN--ADDWATNGVGRK
PC_SK1   TGTSASAPLAAGIFALAL--EANP--NLTWDRMQHLVVWTSEYDPLA--NNPGWKKNAGL
PC_SK5   TGTSASAPMAAGIIALAL--EANPFLTWRDVQHVIVRTSRAGH--LNANDWKTNAAGFKV
          ---* * * *-----*

```

Fig. 4

5/18




Fig.5

6/18

Peps. TLVDEQP----LENYLDMEYFGTIGIGTPAQDFTVVFDTGSSNLWVPSVYCSSL--ACTN  
 Secr. EMVDN-----LRGKSGQGYVEMTVGSPPTLNILVDTGSSNFAVGAAPHPFL-----  
 Cath. PAVTEGP IPEVLKNYMDAQYYGEIGIGTPPQCFTVVFDTGSSNLWVPSIHCKLLDIACWI  
       \* ----1----- \*       \* \* \*       \*\*\*\*\* \*       -----2--

Peps. HNRFPEDSSTYQSTSETVSITYGTGSMTGILGYDTVQV-----G---GISDTN  
 Secr. HRYYQRQLSSTYRDLRKGVVYPYTQGWEGELGTDLVSI-----PHGPNVTVRA  
 Cath. HHKYNSDKSSTYVKNGTSFDIHYGSGSLSGYLSQDTSVPCQSASSASALG---GVKVER  
       -       \*\*\*\*       -----3----- \* \* \*       -----4-----

Peps. QIFGLSETEPGSFLYYAPFDGILGLAYPSIS--SSGATPVFDNIWNQGLVSQDLFSVYLS  
 Secr. NIAAITESDK--FFINGSNWEGILGLAYAEIARPDDSLEPFFDSL VKQTHVP--NLFSLQLC  
 Cath. QVFGKATKQPGITFIAAKFDGILGMAYPRIS--VNNVLPVFDNLMQQLVDQNIFS FYLS  
       -----5-----\*\*\*\* -----6----- \*\*       \* \*       \*\* \*

Peps. ADD-----KS--GSVVIFGGIDSSYYTGSLNWVPVTVEGYWQITVDSITMNGETI  
 Secr. GAGFPLNQSEVLASV--GGSMIIGGIDHSLYTGSLWYTPIRREWYEVIIVRVEINGQDL  
 Cath. RDP-----DAQPGGELMLGGTDSKYYKGSLSYLVNTRKAYWQVHLDQVEVASGLT  
       -----7----- \*\* \*       \* \*\*\*       -----8-----

Peps. A--CAEGC--QAIVDTGTSLLTGPTSPIANIQSDIGASENSD-----GDMVVSCTAI  
 Secr. KMDCKEYNYDKSIVDSGTTNLRPLPKVFEEAAVKSIIKAASSTEKFPDGFWLGEQLV--CWQA  
 Cath. L--CKEGC--EAIVDTGTSLMVGPFVDEVRELQKAIGAVPLIQ-----GEYMIPCEKV  
       \* \*       \*\*\* \*\*       \*       \* \*       -----9----- \*

Peps. SSLPDI VFTI-----NGVQYPVPPSAYILQSEGS----CISGFQGMNVP--TESG  
 Secr. GTPWNIFPVISLYLMGEVTNQSFRTILPQQYLRPVEDV----ATSQDDCYKFAISQSS  
 Cath. STLPAITLKL-----GGKGYKLSPEDYTLKVSQAGKTLCLSGFMGMDIP--PPSG  
       \*       -----10----- \* \*       -----11-----

Peps. ELWILGDVFIRQYFTVFD RANNQVGLAPVA  
 Secr. TGTVMGAVIMEGFYVVFDRARKRIGFAVSA  
 Cath. PLWILGDVFIGRYTTFDRDNNRVGF AEAA  
       --       \* \*       \*\*\*\*       \* \* \*

Fig. 6

7/18




Fig. 7

8/18

01 MLEADDQGCI EEQGVEDSAN EDSVDAKPDR SSFVPSLFSK KKKNVTMRSI KTTRDRVPTY

61 QYNMNFELG KCIIINNKNF DKVTGMGVRN GTDKDAEALF KCFRSLGFDV IVYNDSCAK  
-----1-----

121 MQDLLKKASE EDHTNAACFA CILLSHGEEN VIYKDGVTG IKDLTAHFRG DRSKTILLEK  
-----2-----

181 KLFFIQACRG TELDDGIQAD SGPINDTDAN PRYKIPVEAD FLFAYSTVEG YYSWRSPGRG  
-----3-----4-----

241 SWFVQALCSI LEEHGKDLEI MQILTRVNDR VARHFESQSD DPHFHEKKQI PCVVSMLTKE  
-----5-----

301 LYFSQ

Fig. 8



9/18




Fig. 9

10/18




Fig. 10

11/18




Fig. 11

12/18




Fig. 12

13/18




Fig. 13

14/18




Fig. 14

15/18




Fig. 15

16/18




Fig. 16



17/18




Fig. 17a




Fig. 17b

18/18




Fig. 18

## SEQUENCE LISTING

<110> DIREVO Biotech AG  
 <120> NEW BIOLOGICAL ENTITIES AND THE USE THEREOF  
 <130> 041481wo/JH/ml  
 <160> 126  
 <170> PatentIn version 3.1

<210> 1  
 <211> 224  
 <212> PRT  
 <213> Homo sapiens

<400> 1  
 Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr Gln Val  
 1 5 10 15  
 Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu Ile Asn Glu  
 20 25 30  
 Gln Trp Val Val Ser Ala Gly His Cys Tyr Lys Ser Arg Ile Gln Val  
 35 40 45  
 Arg Leu Gly Glu His Asn Ile Glu Val Leu Glu Gly Asn Glu Gln Phe  
 50 55 60  
 Ile Asn Ala Ala Lys Ile Ile Arg His Pro Gln Tyr Asp Arg Lys Thr  
 65 70 75 80  
 Leu Asn Asn Asp Ile Met Leu Ile Lys Leu Ser Ser Arg Ala Val Ile  
 85 90 95  
 Asn Ala Arg Val Ser Thr Ile Ser Leu Pro Thr Ala Pro Pro Ala Thr  
 100 105 110  
 Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly Asn Thr Ala Ser Ser Gly  
 115 120 125  
 Ala Asp Tyr Pro Asp Glu Leu Gln Cys Leu Asp Ala Pro Val Leu Ser  
 130 135 140  
 Gln Ala Lys Cys Glu Ala Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met  
 145 150 155 160  
 Phe Cys Val Gly Phe Leu Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp  
 165 170 175  
 Ser Gly Gly Pro Val Val Cys Asn Gly Gln Leu Gln Gly Val Val Ser  
 180 185 190  
 Trp Gly Asp Gly Cys Ala Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys  
 195 200 205  
 Val Tyr Asn Tyr Val Lys Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser  
 210 215 220

<210> 2  
 <211> 235  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> trypsin variant 1

<400> 2  
 Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr Gln Val  
 1 5 10 15  
 Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu Ile Asn Glu  
 20 25 30  
 Gln Trp Val Val Ser Ala Gly His Cys Tyr Asp Ala Val Gly Arg Asp  
 35 40 45  
 Lys Ser Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Glu Val Leu  
 50 55 60

```

Glu Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro
65          70          75          80
Gln Tyr Asp Arg Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys Leu
85          90          95
Ser Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser Leu Pro
100         105         110
Thr Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly
115         120         125
Asn Thr Ile Thr Asn Ser Thr Ala Ser Ser Gly Ala Asp Tyr Pro Asp
130         135         140
Glu Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Ala Lys Cys Glu
145         150         155         160
Ala Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly Phe
165         170         175
Leu Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Val
180         185         190
Val Cys Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly Asp Gly Cys
195         200         205
Ala Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val Tyr Asn Tyr Val
210         215         220
Lys Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser
225         230         235

```

```

<210> 3
<211> 235
<212> PRT
<213> artificial sequence
<220>
<223> trypsin variant 2

```

```

<400> 3
Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr Gln Val
1          5          10          15
Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu Ile Asn Glu
20         25         30
Gln Trp Val Val Ser Ala Gly His Cys Tyr Asn Gly Arg Asp Leu Glu
35         40         45
Lys Ser Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Glu Val Leu
50         55         60
Glu Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro
65         70         75         80
Gln Tyr Asp Arg Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys Leu
85         90         95
Ser Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser Leu Pro
100        105        110
Thr Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly
115        120        125
Asn Val Arg Gly Thr Trp Thr Ala Ser Ser Gly Ala Asp Tyr Pro Asp
130        135        140
Glu Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Ala Lys Cys Glu
145        150        155        160
Ala Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly Phe
165        170        175
Leu Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Val
180        185        190
Val Cys Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly Asp Gly Cys
195        200        205
Ala Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val Tyr Asn Tyr Val
210        215        220
Lys Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser
225        230        235

```

<210> 4  
 <211> 235  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> trypsin variant 3

<400> 4  
 Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr Gln Val  
 1 5 10 15  
 Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu Ile Asn Glu  
 20 25 30  
 Gln Trp Val Val Ser Ala Gly His Cys Tyr Ala Ala Thr Asn Gly Asp  
 35 40 45  
 Lys Ser Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Glu Val Leu  
 50 55 60  
 Glu Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro  
 65 70 75 80  
 Gln Tyr Asp Arg Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys Leu  
 85 90 95  
 Ser Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser Leu Pro  
 100 105 110  
 Thr Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly  
 115 120 125  
 Asn Arg Lys Asp Phe Trp Thr Ala Ser Ser Gly Ala Asp Tyr Pro Asp  
 130 135 140  
 Glu Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Ala Lys Cys Glu  
 145 150 155 160  
 Ala Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly Phe  
 165 170 175  
 Leu Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Val  
 180 185 190  
 Val Cys Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly Asp Gly Cys  
 195 200 205  
 Ala Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val Tyr Asn Tyr Val  
 210 215 220  
 Lys Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser  
 225 230 235

<210> 5  
 <211> 259  
 <212> PRT  
 <213> Homo sapiens

<400> 5  
 Ile Val Glu Gly Ser Asp Ala Glu Ile Gly Met Ser Pro Trp Gln Val  
 1 5 10 15  
 Met Leu Phe Arg Lys Ser Pro Gln Glu Leu Leu Cys Gly Ala Ser Leu  
 20 25 30  
 Ile Ser Asp Arg Trp Val Leu Thr Ala Ala His Cys Leu Leu Tyr Pro  
 35 40 45  
 Pro Trp Asp Lys Asn Phe Thr Glu Asn Asp Leu Leu Val Arg Ile Gly  
 50 55 60  
 Lys His Ser Arg Thr Arg Tyr Glu Arg Asn Ile Glu Lys Ile Ser Met  
 65 70 75 80  
 Leu Glu Lys Ile Tyr Ile His Pro Arg Tyr Asn Trp Arg Glu Asn Leu  
 85 90 95  
 Asp Arg Asp Ile Ala Leu Met Lys Leu Lys Lys Pro Val Ala Phe Ser  
 100 105 110  
 Asp Tyr Ile His Pro Val Cys Leu Pro Asp Arg Glu Thr Ala Ala Ser  
 115 120 125

Leu Leu Gln Ala Gly Tyr Lys Gly Arg Val Thr Gly Trp Gly Asn Leu  
 130 135 140  
 Lys Glu Thr Trp Thr Ala Asn Val Gly Lys Gly Gln Pro Ser Val Leu  
 145 150 155 160  
 Gln Val Val Asn Leu Pro Ile Val Glu Arg Pro Val Cys Lys Asp Ser  
 165 170 175  
 Thr Arg Ile Arg Ile Thr Asp Asn Met Phe Cys Ala Gly Tyr Lys Pro  
 180 185 190  
 Asp Glu Gly Lys Arg Gly Asp Ala Cys Glu Gly Asp Ser Gly Gly Pro  
 195 200 205  
 Phe Val Met Lys Ser Pro Phe Asn Asn Arg Trp Tyr Gln Met Gly Ile  
 210 215 220  
 Val Ser Trp Gly Glu Gly Cys Asp Arg Asp Gly Lys Tyr Gly Phe Tyr  
 225 230 235 240  
 Thr His Val Phe Arg Leu Lys Lys Trp Ile Gln Lys Val Ile Asp Gln  
 245 250 255  
 Phe Gly Glu

<210> 6  
 <211> 235  
 <212> PRT  
 <213> Homo sapiens

<400> 6  
 Ile Val Gly Gly Ser Asn Ala Lys Glu Gly Ala Trp Pro Trp Val Val  
 1 5 10 15  
 Gly Leu Tyr Tyr Gly Gly Arg Leu Leu Cys Gly Ala Ser Leu Val Ser  
 20 25 30  
 Ser Asp Trp Leu Val Ser Ala Ala His Cys Val Tyr Gly Arg Asn Leu  
 35 40 45  
 Glu Pro Ser Lys Trp Thr Ala Ile Leu Gly Leu His Met Lys Ser Asn  
 50 55 60  
 Leu Thr Ser Pro Gln Thr Val Pro Arg Leu Ile Asp Glu Ile Val Ile  
 65 70 75 80  
 Asn Pro His Tyr Asn Arg Arg Arg Lys Asp Asn Asp Ile Ala Met Met  
 85 90 95  
 His Leu Glu Phe Lys Val Asn Tyr Thr Asp Tyr Ile Gln Pro Ile Cys  
 100 105 110  
 Leu Pro Glu Glu Asn Gln Val Phe Pro Pro Gly Arg Asn Cys Ser Ile  
 115 120 125  
 Ala Gly Trp Gly Thr Val Val Tyr Gln Gly Thr Thr Ala Asn Ile Leu  
 130 135 140  
 Gln Glu Ala Asp Val Pro Leu Leu Ser Asn Glu Arg Cys Gln Gln Gln  
 145 150 155 160  
 Met Pro Glu Tyr Asn Ile Thr Glu Asn Met Ile Cys Ala Gly Tyr Glu  
 165 170 175  
 Glu Gly Gly Ile Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Met  
 180 185 190  
 Cys Gln Glu Asn Asn Arg Trp Phe Leu Ala Gly Val Thr Ser Phe Gly  
 195 200 205  
 Tyr Lys Cys Ala Leu Pro Asn Arg Pro Gly Val Tyr Ala Arg Val Ser  
 210 215 220  
 Arg Phe Thr Glu Trp Ile Gln Ser Phe Leu His  
 225 230 235

<210> 7  
 <211> 275  
 <212> PRT  
 <213> Bacillus subtilis

&lt;400&gt; 7

```

Ile Ala His Glu Tyr Ala Gln Ser Val Pro Tyr Gly Ile Ser Gln Ile
1          5          10          15
Lys Ala Pro Ala Leu His Ser Gln Gly Tyr Thr Gly Ser Asn Val Lys
          20          25          30
Val Ala Val Ile Asp Ser Gly Ile Asp Ser Ser His Pro Asp Leu Asn
          35          40          45
Val Arg Gly Gly Ala Ser Phe Val Pro Ser Glu Thr Asn Pro Tyr Gln
          50          55          60
Asp Gly Ser Ser His Gly Thr His Val Ala Gly Thr Ile Ala Ala Leu
65          70          75          80
Asn Asn Ser Ile Gly Val Leu Gly Val Ser Pro Ser Ala Ser Leu Tyr
          85          90          95
Ala Val Lys Val Leu Asp Ser Thr Gly Ser Gly Gln Tyr Ser Trp Ile
          100          105          110
Ile Asn Gly Ile Glu Trp Ala Ile Ser Asn Asn Met Asp Val Ile Asn
          115          120          125
Met Ser Leu Gly Gly Pro Thr Gly Ser Thr Ala Leu Lys Thr Val Val
130          135          140
Asp Lys Ala Val Ser Ser Gly Ile Val Val Ala Ala Ala Ala Gly Asn
145          150          155          160
Glu Gly Ser Ser Gly Ser Thr Ser Thr Val Gly Tyr Pro Ala Lys Tyr
          165          170          175
Pro Ser Thr Ile Ala Val Gly Ala Val Asn Ser Ser Asn Gln Arg Ala
          180          185          190
Ser Phe Ser Ser Ala Gly Ser Glu Leu Asp Val Met Ala Pro Gly Val
          195          200          205
Ser Ile Gln Ser Thr Leu Pro Gly Gly Thr Tyr Gly Ala Tyr Asn Gly
210          215          220
Thr Ser Met Ala Thr Pro His Val Ala Gly Ala Ala Ala Leu Ile Leu
225          230          235          240
Ser Lys His Pro Thr Trp Thr Asn Ala Gln Val Arg Asp Arg Leu Glu
          245          250          255
Ser Thr Ala Thr Tyr Leu Gly Asn Ser Phe Tyr Tyr Gly Lys Gly Leu
          260          265          270
Ile Asn Val
          275

```

&lt;210&gt; 8

&lt;211&gt; 320

&lt;212&gt; PRT

&lt;213&gt; Murinae gen. sp.

&lt;400&gt; 8

```

Val Ala Lys Arg Arg Ala Lys Arg Asp Val Tyr Gln Glu Pro Thr Asp
1          5          10          15
Pro Lys Phe Pro Gln Gln Trp Tyr Leu Ser Gly Val Thr Gln Arg Asp
          20          25          30
Leu Asn Val Lys Glu Ala Trp Ala Gln Gly Phe Thr Gly His Gly Ile
          35          40          45
Val Val Ser Ile Leu Asp Asp Gly Ile Glu Lys Asn His Pro Asp Leu
          50          55          60
Ala Gly Asn Tyr Asp Pro Gly Ala Ser Phe Asp Val Asn Asp Gln Asp
65          70          75          80
Pro Asp Pro Gln Pro Arg Tyr Thr Gln Met Asn Asp Asn Arg His Gly
          85          90          95
Thr Arg Cys Ala Gly Glu Val Ala Ala Val Ala Asn Asn Gly Val Cys
          100          105          110
Gly Val Gly Val Ala Tyr Asn Ala Arg Ile Gly Gly Val Arg Met Leu
          115          120          125
Asp Gly Glu Val Thr Asp Ala Val Glu Ala Arg Ser Leu Gly Leu Asn
130          135          140

```

```

Pro Asn His Ile His Ile Tyr Ser Ala Ser Trp Gly Pro Glu Asp Asp
145                               150           155           160
Gly Lys Thr Val Asp Gly Pro Ala Arg Leu Ala Glu Glu Ala Phe Phe
                               165           170           175
Arg Gly Val Ser Gln Gly Arg Gly Gly Leu Gly Ser Ile Phe Val Trp
                               180           185           190
Ala Ser Gly Asn Gly Gly Arg Glu His Asp Ser Cys Asn Cys Asp Gly
                               195           200           205
Tyr Thr Asn Ser Ile Tyr Thr Leu Ser Ile Ser Ser Ala Thr Gln Phe
                               210           215           220
Gly Asn Val Pro Trp Tyr Ser Glu Ala Cys Ser Ser Thr Leu Ala Thr
225                               230           235           240
Thr Tyr Ser Ser Gly Asn Gln Asn Glu Lys Gln Ile Val Thr Thr Asp
                               245           250           255
Leu Arg Gln Lys Cys Thr Glu Ser His Thr Gly Thr Ser Ala Ser Ala
                               260           265           270
Pro Leu Ala Ala Gly Ile Ile Ala Leu Thr Leu Glu Ala Asn Lys Asn
                               275           280           285
Leu Thr Trp Arg Asp Met Gln His Leu Val Val Gln Thr Ser Lys Pro
                               290           295           300
Ala His Leu Asn Ala Asp Asp Trp Ala Thr Asn Gly Val Gly Arg Lys
305                               310           315           320

```

```

<210> 9
<211> 330
<212> PRT
<213> Homo sapiens

```

```

<400> 9
Glu Lys Glu Arg Ser Lys Arg Ser Ala Leu Arg Asp Ser Ala Leu Asn
1                               5           10           15
Leu Phe Asn Asp Pro Met Trp Asn Gln Gln Trp Tyr Leu Gln Asp Thr
                               20           25           30
Arg Met Thr Ala Ala Leu Pro Lys Leu Asp Leu His Val Ile Pro Val
                               35           40           45
Trp Gln Lys Gly Ile Thr Gly Lys Gly Val Val Ile Thr Val Leu Asp
50                               55           60
Asp Gly Leu Glu Trp Asn His Thr Asp Ile Tyr Ala Asn Tyr Asp Pro
65                               70           75           80
Glu Ala Ser Tyr Asp Phe Asn Asp Asn Asp His Asp Pro Phe Pro Arg
                               85           90           95
Tyr Asp Pro Thr Asn Glu Asn Lys His Gly Thr Arg Cys Ala Gly Glu
                               100          105          110
Ile Ala Met Gln Ala Asn Asn His Lys Cys Gly Val Gly Val Ala Tyr
115                               120          125
Asn Ser Lys Val Gly Gly Ile Arg Met Leu Asp Gly Ile Val Thr Asp
130                               135          140
Ala Ile Glu Ala Ser Ser Ile Gly Phe Asn Pro Gly His Val Asp Ile
145                               150          155          160
Tyr Ser Ala Ser Trp Gly Pro Asn Asp Asp Gly Lys Thr Val Glu Gly
                               165          170          175
Pro Gly Arg Leu Ala Gln Lys Ala Phe Glu Tyr Gly Val Lys Gln Gly
                               180          185          190
Arg Gln Gly Lys Gly Ser Ile Phe Val Trp Ala Ser Gly Asn Gly Gly
195                               200          205
Arg Gln Gly Asp Asn Cys Asp Cys Asp Gly Tyr Thr Asp Ser Ile Tyr
210                               215          220
Thr Ile Ser Ile Ser Ser Ala Ser Gln Gln Gly Leu Ser Pro Trp Tyr
225                               230          235          240
Ala Glu Lys Cys Ser Ser Thr Leu Ala Thr Ser Tyr Ser Ser Gly Asp
                               245          250          255
Tyr Thr Asp Gln Arg Ile Thr Ser Ala Asp Leu His Asn Asp Cys Thr

```



				260					265					270		
Glu	Thr	His	Thr	Gly	Thr	Ser	Ala	Ser	Ala	Pro	Leu	Ala	Ala	Gly	Ile	
				275					280				285			
Phe	Ala	Leu	Ala	Leu	Glu	Ala	Asn	Pro	Asn	Leu	Thr	Trp	Arg	Asp	Met	
				290			295				300					
Gln	His	Leu	Val	Val	Trp	Thr	Ser	Glu	Tyr	Asp	Pro	Leu	Ala	Asn	Asn	
						310				315					320	
Pro	Gly	Trp	Lys	Lys	Asn	Gly	Ala	Gly	Leu							
				325					330							

```
<210> 10
<211> 297
<212> PRT
<213> Homo sapiens
```

[illegible]

```
<210> 11
<211> 328
<212> PRT
<213> Homo sapiens
```

<400> 11

Thr Leu Val Asp Glu Gln Pro Leu Glu Asn Tyr Leu Asp Met Glu Tyr  
 1 5 10 15  
 Phe Gly Thr Ile Gly Ile Gly Thr Pro Ala Gln Asp Phe Thr Val Val  
 20 25 30  
 Phe Asp Thr Gly Ser Ser Asn Leu Trp Val Pro Ser Val Tyr Cys Ser  
 35 40 45  
 Ser Leu Ala Cys Thr Asn His Asn Arg Phe Asn Pro Glu Asp Ser Ser  
 50 55 60  
 Thr Tyr Gln Ser Thr Ser Glu Thr Val Ser Ile Thr Tyr Gly Thr Gly  
 65 70 75 80  
 Ser Met Thr Gly Ile Leu Gly Tyr Asp Thr Val Gln Val Gly Gly Ile  
 85 90 95  
 Ser Asp Thr Asn Gln Ile Phe Gly Leu Ser Glu Thr Glu Pro Gly Ser  
 100 105 110  
 Phe Leu Tyr Tyr Ala Pro Phe Asp Gly Ile Leu Gly Leu Ala Tyr Pro  
 115 120 125  
 Ser Ile Ser Ser Ser Gly Ala Thr Pro Val Phe Asp Asn Ile Trp Asn  
 130 135 140  
 Gln Gly Leu Val Ser Gln Asp Leu Phe Ser Val Tyr Leu Ser Ala Asp  
 145 150 155 160  
 Asp Lys Ser Gly Ser Val Val Ile Phe Gly Gly Ile Asp Ser Ser Tyr  
 165 170 175  
 Tyr Thr Gly Ser Leu Asn Trp Val Pro Val Thr Val Glu Gly Tyr Trp  
 180 185 190  
 Gln Ile Thr Val Asp Ser Ile Thr Met Asn Gly Glu Thr Ile Ala Cys  
 195 200 205  
 Ala Glu Gly Cys Gln Ala Ile Val Asp Thr Gly Thr Ser Leu Leu Thr  
 210 215 220  
 Gly Pro Thr Ser Pro Ile Ala Asn Ile Gln Ser Asp Ile Gly Ala Ser  
 225 230 235 240  
 Glu Asn Ser Asp Gly Asp Met Val Val Ser Cys Ser Ala Ile Ser Ser  
 245 250 255  
 Leu Pro Asp Ile Val Phe Thr Ile Asn Gly Val Gln Tyr Pro Val Pro  
 260 265 270  
 Pro Ser Ala Tyr Ile Leu Gln Ser Glu Gly Ser Cys Ile Ser Gly Phe  
 275 280 285  
 Gln Gly Met Asn Val Pro Thr Glu Ser Gly Glu Leu Trp Ile Leu Gly  
 290 295 300  
 Asp Val Phe Ile Arg Gln Tyr Phe Thr Val Phe Asp Arg Ala Asn Asn  
 305 310 315 320  
 Gln Val Gly Leu Ala Pro Val Ala  
 325

<210> 12  
 <211> 358  
 <212> PRT  
 <213> Homo sapiens

<400> 12  
 Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val  
 1 5 10 15  
 Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp  
 20 25 30  
 Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu  
 35 40 45  
 His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg  
 50 55 60  
 Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu  
 65 70 75 80  
 Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg  
 85 90 95  
 Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly

```

      100      105      110
Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg
      115      120      125
Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr
      130      135      140
His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro
      145      150      155
Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile
      165      170      175
Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro
      180      185      190
Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile
      195      200      205
Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys
      210      215      220
Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val
      225      230      235
Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys
      245      250      255
Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala
      260      265      270
Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met
      275      280      285
Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln
      290      295      300
Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr
      305      310      315
Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val
      325      330      335
Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile
      340      345      350
Gly Phe Ala Val Ser Ala
      355

```

```

<210> 13
<211> 351
<212> PRT
<213> Homo sapiens

```

```

<400> 13
Pro Ala Val Thr Glu Gly Pro Ile Pro Glu Val Leu Lys Asn Tyr Met
1      5      10      15
Asp Ala Gln Tyr Tyr Gly Glu Ile Gly Ile Gly Thr Pro Pro Gln Cys
      20      25      30
Phe Thr Val Val Phe Asp Thr Gly Ser Ser Asn Leu Trp Val Pro Ser
      35      40      45
Ile His Cys Lys Leu Leu Asp Ile Ala Cys Trp Ile His His Lys Tyr
      50      55      60
Asn Ser Asp Lys Ser Ser Thr Tyr Val Lys Asn Gly Thr Ser Phe Asp
      65      70      75      80
Ile His Tyr Gly Ser Gly Ser Leu Ser Gly Tyr Leu Ser Gln Asp Thr
      85      90      95
Val Ser Val Pro Cys Gln Ser Ala Ser Ser Ala Ser Ala Leu Gly Gly
      100      105      110
Val Lys Val Glu Arg Gln Val Phe Gly Glu Ala Thr Lys Gln Pro Gly
      115      120      125
Ile Thr Phe Ile Ala Ala Lys Phe Asp Gly Ile Leu Gly Met Ala Tyr
      130      135      140
Pro Arg Ile Ser Val Asn Asn Val Leu Pro Val Phe Asp Asn Leu Met
      145      150      155      160
Gln Gln Lys Leu Val Asp Gln Asn Ile Phe Ser Phe Tyr Leu Ser Arg
      165      170      175

```

```

Asp Pro Asp Ala Gln Pro Gly Gly Glu Leu Met Leu Gly Gly Thr Asp
      180      185      190
Ser Lys Tyr Tyr Lys Gly Ser Leu Ser Tyr Leu Asn Val Thr Arg Lys
      195      200      205
Ala Tyr Trp Gln Val His Leu Asp Gln Val Glu Val Ala Ser Gly Leu
      210      215      220
Thr Leu Cys Lys Glu Gly Cys Glu Ala Ile Val Asp Thr Gly Thr Ser
225      230      235      240
Leu Met Val Gly Pro Val Asp Glu Val Arg Glu Leu Gln Lys Ala Ile
      245      250      255
Gly Ala Val Pro Leu Ile Gln Gly Glu Tyr Met Ile Pro Cys Glu Lys
      260      265      270
Val Ser Thr Leu Pro Ala Ile Thr Leu Lys Leu Gly Gly Lys Gly Tyr
      275      280      285
Lys Leu Ser Pro Glu Asp Tyr Thr Leu Lys Val Ser Gln Ala Gly Lys
      290      295      300
Thr Leu Cys Leu Ser Gly Phe Met Gly Met Asp Ile Pro Pro Pro Ser
305      310      315      320
Gly Pro Leu Trp Ile Leu Gly Asp Val Phe Ile Gly Arg Tyr Tyr Thr
      325      330      335
Val Phe Asp Arg Asp Asn Asn Arg Val Gly Phe Ala Glu Ala Ala
      340      345      350

```

```

<210> 14
<211> 305
<212> PRT
<213> Homo sapiens

```

```

<400> 14
Met Leu Glu Ala Asp Asp Gln Gly Cys Ile Glu Glu Gln Gly Val Glu
1      5      10      15
Asp Ser Ala Asn Glu Asp Ser Val Asp Ala Lys Pro Asp Arg Ser Ser
      20      25      30
Phe Val Pro Ser Leu Phe Ser Lys Lys Lys Lys Asn Val Thr Met Arg
      35      40      45
Ser Ile Lys Thr Thr Arg Asp Arg Val Pro Thr Tyr Gln Tyr Asn Met
      50      55      60
Asn Phe Glu Lys Leu Gly Lys Cys Ile Ile Ile Asn Asn Lys Asn Phe
      65      70      75      80
Asp Lys Val Thr Gly Met Gly Val Arg Asn Gly Thr Asp Lys Asp Ala
      85      90      95
Glu Ala Leu Phe Lys Cys Phe Arg Ser Leu Gly Phe Asp Val Ile Val
      100      105      110
Tyr Asn Asp Cys Ser Cys Ala Lys Met Gln Asp Leu Leu Lys Lys Ala
      115      120      125
Ser Glu Glu Asp His Thr Asn Ala Ala Cys Phe Ala Cys Ile Leu Leu
      130      135      140
Ser His Gly Glu Glu Asn Val Ile Tyr Gly Lys Asp Gly Val Thr Pro
      145      150      155      160
Ile Lys Asp Leu Thr Ala His Phe Arg Gly Asp Arg Ser Lys Thr Leu
      165      170      175
Leu Glu Lys Pro Lys Leu Phe Phe Ile Gln Ala Cys Arg Gly Thr Glu
      180      185      190
Leu Asp Asp Gly Ile Gln Ala Asp Ser Gly Pro Ile Asn Asp Thr Asp
      195      200      205
Ala Asn Pro Arg Tyr Lys Ile Pro Val Glu Ala Asp Phe Leu Phe Ala
      210      215      220
Tyr Ser Thr Val Pro Gly Tyr Tyr Ser Trp Arg Ser Pro Gly Arg Gly
      225      230      235      240
Ser Trp Phe Val Gln Ala Leu Cys Ser Ile Leu Glu Glu His Gly Lys
      245      250      255
Asp Leu Glu Ile Met Gln Ile Leu Thr Arg Val Asn Asp Arg Val Ala

```

[illegible]

```
<210> 15
<211> 262
<212> PRT
<213> Streptomyces sp. K15
```

[illegible]

```
<210> 16
<211> 256
<212> PRT
<213> Human cytomegalovirus
```

```

<400> 16
Met Thr Met Asp Glu Gln Gln Ser Gln Ala Val Ala Pro Val Tyr Val
1          5          10          15
Gly Gly Phe Leu Ala Arg Tyr Asp Gln Ser Pro Asp Glu Ala Glu Leu
          20          25          30
Leu Leu Pro Arg Asp Val Val Glu His Trp Leu His Ala Gln Gly Gln
          35          40          45

```

Gly Gln Pro Ser Leu Ser Val Ala Leu Pro Leu Asn Ile Asn His Asp  
 50 55 60  
 Asp Thr Ala Val Val Gly His Val Ala Ala Met Gln Ser Val Arg Asp  
 65 70 75 80  
 Gly Leu Phe Cys Leu Gly Cys Val Thr Ser Pro Arg Phe Leu Glu Ile  
 85 90 95  
 Val Arg Arg Ala Ser Glu Lys Ser Glu Leu Val Ser Arg Gly Pro Val  
 100 105 110  
 Ser Pro Leu Gln Pro Asp Lys Val Val Glu Phe Leu Ser Gly Ser Tyr  
 115 120 125  
 Ala Gly Leu Ser Leu Ser Ser Arg Arg Cys Asp Asp Val Glu Gln Ala  
 130 135 140  
 Thr Ser Leu Ser Gly Ser Glu Thr Thr Pro Phe Lys His Val Ala Leu  
 145 150 155 160  
 Cys Ser Val Gly Arg Arg Arg Gly Thr Leu Ala Val Tyr Gly Arg Asp  
 165 170 175  
 Pro Glu Trp Val Thr Gln Arg Phe Pro Asp Leu Thr Ala Ala Asp Arg  
 180 185 190  
 Asp Gly Leu Arg Ala Gln Trp Gln Arg Cys Gly Ser Thr Ala Val Asp  
 195 200 205  
 Ala Ser Gly Asp Pro Phe Arg Ser Asp Ser Tyr Gly Leu Leu Gly Asn  
 210 215 220  
 Ser Val Asp Ala Leu Tyr Ile Arg Glu Arg Leu Pro Lys Leu Arg Tyr  
 225 230 235 240  
 Asp Lys Gln Leu Val Gly Val Thr Glu Arg Glu Ser Tyr Val Lys Ala  
 245 250 255

<210> 17  
 <211> 248  
 <212> PRT  
 <213> Escherichia coli

<400> 17  
 Val Arg Ser Phe Ile Tyr Glu Pro Phe Gln Ile Pro Ser Gly Ser Met  
 1 5 10 15  
 Met Pro Thr Leu Leu Ile Gly Asp Phe Ile Leu Val Glu Lys Phe Ala  
 20 25 30  
 Tyr Gly Ile Lys Asp Pro Ile Tyr Gln Lys Thr Leu Ile Glu Thr Gly  
 35 40 45  
 His Pro Lys Arg Gly Asp Ile Val Val Phe Lys Tyr Pro Glu Asp Pro  
 50 55 60  
 Lys Leu Asp Tyr Ile Lys Arg Ala Val Gly Leu Pro Gly Asp Lys Val  
 65 70 75 80  
 Thr Tyr Asp Pro Val Ser Lys Glu Leu Thr Ile Gln Pro Gly Cys Ser  
 85 90 95  
 Ser Gly Gln Ala Cys Glu Asn Ala Leu Pro Val Thr Tyr Ser Asn Val  
 100 105 110  
 Glu Pro Ser Asp Phe Val Gln Thr Phe Ser Arg Arg Asn Gly Gly Glu  
 115 120 125  
 Ala Thr Ser Gly Phe Phe Glu Val Pro Lys Asn Glu Thr Lys Glu Asn  
 130 135 140  
 Gly Ile Arg Leu Ser Glu Arg Lys Glu Thr Leu Gly Asp Val Thr His  
 145 150 155 160  
 Arg Ile Leu Thr Val Pro Ile Ala Gln Asp Gln Val Gly Met Tyr Tyr  
 165 170 175  
 Gln Gln Pro Gly Gln Gln Leu Ala Thr Trp Ile Val Pro Pro Gly Gln  
 180 185 190  
 Tyr Phe Met Met Gly Asp Asn Arg Asp Asn Ser Ala Asp Ser Arg Tyr  
 195 200 205  
 Trp Gly Phe Val Pro Glu Ala Asn Leu Val Gly Arg Ala Thr Ala Ile  
 210 215 220  
 Trp Met Ser Phe Asp Lys Gln Glu Gly Glu Trp Pro Thr Gly Leu Arg

225                                  230  
Leu Ser Arg Ile Gly Gly Ile His  
                                245

```
<210> 18
<211> 317
<212> PRT
<213> Serratia marcescens
```

<400>	18															
Met	Glu	Gln	Leu	Arg	Gly	Leu	Tyr	Pro	Pro	Leu	Ala	Ala	Tyr	Asp	Ser	
1				5					10					15		
Gly	Trp	Leu	Asp	Thr	Gly	Asp	Gly	His	Arg	Ile	Tyr	Trp	Glu	Leu	Ser	
			20					25					30			
Gly	Asn	Pro	Asn	Gly	Lys	Pro	Ala	Val	Phe	Ile	His	Gly	Gly	Pro	Gly	
		35					40					45				
Gly	Gly	Ile	Ser	Pro	His	His	Arg	Gln	Leu	Phe	Asp	Pro	Glu	Arg	Tyr	
	50					55					60					
Lys	Val	Leu	Leu	Phe	Asp	Gln	Arg	Gly	Cys	Gly	Arg	Ser	Arg	Pro	His	
65					70					75					80	
Ala	Ser	Leu	Asp	Asn	Asn	Thr	Thr	Trp	His	Leu	Val	Ala	Asp	Ile	Glu	
				85					90					95		
Arg	Leu	Arg	Glu	Met	Ala	Gly	Val	Glu	Gln	Trp	Leu	Val	Phe	Gly	Gly	
			100					105					110			
Ser	Trp	Gly	Ser	Thr	Leu	Ala	Leu	Ala	Tyr	Ala	Gln	Thr	His	Pro	Glu	
		115					120				125					
Arg	Val	Ser	Glu	Met	Val	Leu	Arg	Gly	Ile	Phe	Thr	Leu	Arg	Lys	Gln	
	130					135					140					
Arg	Leu	His	Trp	Tyr	Tyr	Gln	Asp	Gly	Ala	Ser	Arg	Phe	Phe	Pro	Glu	
145					150					155					160	
Lys	Trp	Glu	Arg	Val	Leu	Ser	Ile	Leu	Ser	Asp	Asp	Glu	Arg	Lys	Asp	
				165					170					175		
Val	Ile	Ala	Ala	Tyr	Arg	Gln	Arg	Leu	Thr	Ser	Ala	Asp	Pro	Gln	Val	
			180					185					190			
Gln	Leu	Glu	Ala	Ala	Lys	Leu	Trp	Ser	Val	Trp	Glu	Gly	Glu	Thr	Val	
		195					200					205				
Thr	Leu	Leu	Pro	Ser	Arg	Glu	Ser	Ala	Ser	Phe	Gly	Glu	Asp	Asp	Phe	
	210					215					220					
Ala	Leu	Ala	Phe	Ala	Arg	Ile	Glu	Asn	His	Tyr	Phe	Thr	His	Leu	Gly	
225					230					235					240	
Phe	Leu	Glu	Ser	Asp	Asp	Gln	Leu	Leu	Arg	Asn	Val	Pro	Leu	Ile	Arg	
				245					250					255		
His	Ile	Pro	Ala	Val	Ile	Val	His	Gly	Arg	Tyr	Asp	Met	Ala	Cys	Gln	
			260					265				270				
Val	Gln	Asn	Ala	Trp	Asp	Leu	Ala	Lys	Ala	Trp	Pro	Glu	Ala	Glu	Leu	
		275					280					285				
His	Ile	Val	Glu	Gly	Ala	Gly	His	Ser	Tyr	Asp	Glu	Pro	Gly	Ile	Leu	
	290					295					300					
His	Gln	Leu	Met	Ile	Ala	Thr	Asp	Arg	Phe	Ala	Gly	Lys				
305					310					315						

```
<210> 19
<211> 229
<212> PRT
<213> Escherichia coli
```

```

<400> 19
Met Glu Leu Leu Leu Leu Ser Asn Ser Thr Leu Pro Gly Lys Ala Trp
1          5          10          15
Leu Glu His Ala Leu Pro Leu Ile Ala Asn Gln Leu Asn Gly Arg Arg
          20          25          30

```

Ser Ala Val Phe Ile Pro Phe Ala Gly Val Thr Gln Thr Trp Asp Glu  
 35 40 45  
 Tyr Thr Asp Lys Thr Ala Glu Val Leu Ala Pro Leu Gly Val Asn Val  
 50 55 60  
 Thr Gly Ile His Arg Val Ala Asp Pro Leu Ala Ala Ile Glu Lys Ala  
 65 70 75 80  
 Glu Ile Ile Ile Val Gly Gly Gly Asn Thr Phe Gln Leu Leu Lys Glu  
 85 90 95  
 Ser Arg Glu Arg Gly Leu Leu Ala Pro Met Ala Asp Arg Val Lys Arg  
 100 105 110  
 Gly Ala Leu Tyr Ile Gly Trp Ser Ala Gly Ala Asn Leu Ala Cys Pro  
 115 120 125  
 Thr Ile Arg Thr Thr Asn Asp Met Pro Ile Val Asp Pro Asn Gly Phe  
 130 135 140  
 Asp Ala Leu Asp Leu Phe Pro Leu Gln Ile Asn Pro His Phe Thr Asn  
 145 150 155 160  
 Ala Leu Pro Glu Gly His Lys Gly Glu Thr Arg Glu Gln Arg Ile Arg  
 165 170 175  
 Glu Leu Leu Val Val Ala Pro Glu Leu Thr Val Ile Gly Leu Pro Glu  
 180 185 190  
 Gly Asn Trp Ile Gln Val Ser Asn Gly Gln Ala Val Leu Gly Gly Pro  
 195 200 205  
 Asn Thr Thr Trp Val Phe Lys Ala Gly Glu Glu Ala Val Ala Leu Glu  
 210 215 220  
 Ala Gly His Arg Phe  
 225

<210> 20  
 <211> 99  
 <212> PRT  
 <213> Human immunodeficiency virus

<400> 20  
 Pro Gln Ile Thr Leu Trp Gln Arg Pro Leu Val Thr Val Lys Ile Gly  
 1 5 10 15  
 Gly Gln Leu Arg Glu Ala Leu Leu Asp Thr Gly Ala Asp Asp Thr Val  
 20 25 30  
 Leu Glu Asp Ile Asn Leu Pro Gly Lys Trp Lys Pro Lys Met Ile Gly  
 35 40 45  
 Gly Ile Gly Gly Phe Ile Lys Val Arg Gln Tyr Asp Gln Ile Leu Ile  
 50 55 60  
 Glu Ile Cys Gly Lys Lys Ala Ile Gly Thr Val Leu Val Gly Pro Thr  
 65 70 75 80  
 Pro Val Asn Ile Ile Gly Arg Asn Met Leu Thr Gln Ile Gly Cys Thr  
 85 90 95  
 Leu Asn Phe

<210> 21  
 <211> 297  
 <212> PRT  
 <213> Escherichia coli

<400> 21  
 Ser Thr Glu Thr Leu Ser Phe Thr Pro Asp Asn Ile Asn Ala Asp Ile  
 1 5 10 15  
 Ser Leu Gly Thr Leu Ser Gly Lys Thr Lys Glu Arg Val Tyr Leu Ala  
 20 25 30  
 Glu Glu Gly Gly Arg Lys Val Ser Gln Leu Asp Trp Lys Phe Asn Asn  
 35 40 45  
 Ala Ala Ile Ile Lys Gly Ala Ile Asn Trp Asp Leu Met Pro Gln Ile



50	55	60
Ser Ile Gly Ala Ala Gly	Trp Thr Thr Leu Gly	Ser Arg Gly Gly Asn
65	70	75
Met Val Asp Gln Asp Trp	Met Asp Ser Ser Asn	Pro Gly Thr Trp Thr
85	90	95
Asp Glu Ala Arg His Pro	Asp Thr Gln Leu Asn	Tyr Ala Asn Glu Phe
100	105	110
Asp Leu Asn Ile Lys Gly	Trp Leu Leu Asn Glu	Pro Asn Tyr Arg Leu
115	120	125
Gly Leu Met Ala Gly Tyr	Gln Glu Ser Arg Tyr	Ser Phe Thr Ala Arg
130	135	140
Gly Gly Ser Tyr Ile Tyr	Ser Ser Glu Glu Gly	Phe Arg Asp Asp Ile
145	150	155
Gly Ser Phe Pro Asn Gly	Glu Arg Ala Ile Gly	Tyr Lys Gln Arg Phe
165	170	175
Lys Met Pro Tyr Ile Gly	Leu Thr Gly Ser Tyr	Arg Tyr Glu Asp Phe
180	185	190
Glu Leu Gly Gly Thr Phe	Lys Tyr Ser Gly Trp	Val Glu Ser Ser Asp
195	200	205
Asn Asp Glu His Tyr Asp	Pro Lys Gly Arg Ile	Thr Tyr Arg Ser Lys
210	215	220
Val Lys Asp Gln Asn Tyr	Tyr Ser Val Ala Val	Asn Ala Gly Tyr Tyr
225	230	235
Val Thr Pro Asn Ala Lys	Val Tyr Val Glu Gly	Ala Trp Asn Arg Val
245	250	255
Thr Asn Lys Lys Gly Asn	Thr Ser Leu Tyr Asp	His Asn Asn Asn Thr
260	265	270
Ser Asp Tyr Ser Lys Asn	Gly Ala Gly Ile Glu	Asn Tyr Asn Phe Ile
275	280	285
Thr Thr Ala Gly Leu Lys	Tyr Thr Phe	
290	295	

<210> 22  
 <211> 212  
 <212> PRT  
 <213> Carica papaya

<400> 22

Ile Pro Glu Tyr Val Asp	Trp Arg Gln Lys Gly	Ala Val Thr Pro Val
1	5	10
Lys Asn Gln Gly Ser Cys	Gly Ser Cys Trp Ala	Phe Ser Ala Val Val
20	25	30
Thr Ile Glu Gly Ile Ile	Lys Ile Arg Thr Gly	Asn Leu Asn Gln Tyr
35	40	45
Ser Glu Gln Glu Leu Leu	Asp Cys Asp Arg Arg	Ser Tyr Gly Cys Asn
50	55	60
Gly Gly Tyr Pro Trp Ser	Ala Leu Gln Leu Val	Ala Gln Tyr Gly Ile
65	70	75
His Tyr Arg Asn Thr Tyr	Pro Tyr Glu Gly Val	Gln Arg Tyr Cys Arg
85	90	95
Ser Arg Glu Lys Gly Pro	Tyr Ala Ala Lys Thr	Asp Gly Val Arg Gln
100	105	110
Val Gln Pro Tyr Asn Gln	Gly Ala Leu Leu Tyr	Ser Ile Ala Asn Gln
115	120	125
Pro Val Ser Val Val Leu	Gln Ala Ala Gly Lys	Asp Phe Gln Leu Tyr
130	135	140
Arg Gly Gly Ile Phe Val	Gly Pro Cys Gly Asn	Lys Val Asp His Ala
145	150	155
Val Ala Ala Val Gly Tyr	Gly Pro Asn Tyr Ile	Leu Ile Lys Asn Ser
165	170	175
Trp Gly Thr Gly Trp Gly	Glu Asn Gly Tyr Ile	Arg Ile Lys Arg Gly
180	185	190

Thr Gly Asn Ser Tyr Gly Val Cys Gly Leu Tyr Thr Ser Ser Phe Tyr  
 195 200 205  
 Pro Val Lys Asn  
 210

<210> 23  
 <211> 699  
 <212> PRT  
 <213> Homo sapiens

<400> 23  
 Ala Gly Ile Ala Ala Lys Leu Ala Lys Asp Arg Glu Ala Ala Glu Gly  
 1 5 10 15  
 Leu Gly Ser His Glu Arg Ala Ile Lys Tyr Leu Asn Gln Asp Tyr Glu  
 20 25 30  
 Ala Leu Arg Asn Glu Cys Leu Glu Ala Gly Thr Leu Phe Gln Asp Pro  
 35 40 45  
 Ser Phe Pro Ala Ile Pro Ser Ala Leu Gly Phe Lys Glu Leu Gly Pro  
 50 55 60  
 Tyr Ser Ser Lys Thr Arg Gly Met Arg Trp Lys Arg Pro Thr Glu Ile  
 65 70 75 80  
 Cys Ala Asp Pro Gln Phe Ile Ile Gly Gly Ala Thr Arg Thr Asp Ile  
 85 90 95  
 Cys Gln Gly Ala Leu Gly Asp Cys Trp Leu Leu Ala Ala Ile Ala Ser  
 100 105 110  
 Leu Thr Leu Asn Glu Glu Ile Leu Ala Arg Val Val Pro Leu Asn Gln  
 115 120 125  
 Ser Phe Gln Glu Asn Tyr Ala Gly Ile Phe His Phe Gln Phe Trp Gln  
 130 135 140  
 Tyr Gly Glu Trp Val Glu Val Val Val Asp Asp Arg Leu Pro Thr Lys  
 145 150 155 160  
 Asp Gly Glu Leu Leu Phe Val His Ser Ala Glu Gly Ser Glu Phe Trp  
 165 170 175  
 Ser Ala Leu Leu Glu Lys Ala Tyr Ala Lys Ile Asn Gly Cys Tyr Glu  
 180 185 190  
 Ala Leu Ser Gly Gly Ala Thr Thr Glu Gly Phe Glu Asp Phe Thr Gly  
 195 200 205  
 Gly Ile Ala Glu Trp Tyr Glu Leu Lys Lys Pro Pro Pro Asn Leu Phe  
 210 215 220  
 Lys Ile Ile Gln Lys Ala Leu Gln Lys Gly Ser Leu Leu Gly Cys Ser  
 225 230 235 240  
 Ile Asp Ile Thr Ser Ala Ala Asp Ser Glu Ala Ile Thr Phe Gln Lys  
 245 250 255  
 Leu Val Lys Gly His Ala Tyr Ser Val Thr Gly Ala Glu Glu Val Glu  
 260 265 270  
 Ser Asn Gly Ser Leu Gln Lys Leu Ile Arg Ile Arg Asn Pro Trp Gly  
 275 280 285  
 Glu Val Glu Trp Thr Gly Arg Trp Asn Asp Asn Cys Pro Ser Trp Asn  
 290 295 300  
 Thr Ile Asp Pro Glu Glu Arg Glu Arg Leu Thr Arg Arg His Glu Asp  
 305 310 315 320  
 Gly Glu Phe Trp Met Ser Phe Ser Asp Phe Leu Arg His Tyr Ser Arg  
 325 330 335  
 Leu Glu Ile Cys Asn Leu Thr Pro Asp Thr Leu Thr Ser Asp Thr Tyr  
 340 345 350  
 Lys Lys Trp Lys Leu Thr Lys Met Asp Gly Asn Trp Arg Arg Gly Ser  
 355 360 365  
 Thr Ala Gly Gly Cys Arg Asn Tyr Pro Asn Thr Phe Trp Met Asn Pro  
 370 375 380  
 Gln Tyr Leu Ile Lys Leu Glu Glu Glu Asp Glu Asp Glu Glu Asp Gly  
 385 390 395 400  
 Glu Ser Gly Cys Thr Phe Leu Val Gly Leu Ile Gln Lys His Arg Arg

				405					410					415	
Arg	Gln	Arg	Lys	Met	Gly	Glu	Asp	Met	His	Thr	Ile	Gly	Phe	Gly	Ile
			420					425					430		
Tyr	Glu	Val	Pro	Glu	Glu	Leu	Ser	Gly	Gln	Thr	Asn	Ile	His	Leu	Ser
		435					440				445				
Lys	Asn	Phe	Phe	Leu	Thr	Asn	Arg	Ala	Arg	Glu	Arg	Ser	Asp	Thr	Phe
	450					455				460					
Ile	Asn	Leu	Arg	Glu	Val	Leu	Asn	Arg	Phe	Lys	Leu	Pro	Pro	Gly	Glu
465					470					475					480
Tyr	Ile	Leu	Val	Pro	Ser	Thr	Phe	Glu	Pro	Asn	Lys	Asp	Gly	Asp	Phe
			485						490					495	
Cys	Ile	Arg	Val	Phe	Ser	Glu	Lys	Lys	Ala	Asp	Tyr	Gln	Ala	Val	Asp
			500					505					510		
Asp	Glu	Ile	Glu	Ala	Asn	Leu	Glu	Glu	Phe	Asp	Ile	Ser	Glu	Asp	Asp
		515					520					525			
Ile	Asp	Asp	Gly	Val	Arg	Arg	Leu	Phe	Ala	Gln	Leu	Ala	Gly	Glu	Asp
	530					535					540				
Ala	Glu	Ile	Ser	Ala	Phe	Glu	Leu	Gln	Thr	Ile	Leu	Arg	Arg	Val	Leu
545					550					555					560
Ala	Lys	Arg	Gln	Asp	Ile	Lys	Ser	Asp	Gly	Phe	Ser	Ile	Glu	Thr	Cys
			565					570					575		
Lys	Ile	Met	Val	Asp	Met	Leu	Asp	Ser	Asp	Gly	Ser	Gly	Lys	Leu	Gly
		580					585					590			
Leu	Lys	Glu	Phe	Tyr	Ile	Leu	Trp	Thr	Lys	Ile	Gln	Lys	Tyr	Gln	Lys
		595				600						605			
Ile	Tyr	Arg	Glu	Ile	Asp	Val	Asp	Arg	Ser	Gly	Thr	Met	Asn	Ser	Tyr
	610				615					620					
Glu	Met	Arg	Lys	Ala	Leu	Glu	Glu	Ala	Gly	Phe	Lys	Met	Pro	Cys	Gln
625					630					635					640
Leu	His	Gln	Val	Ile	Val	Ala	Arg	Phe	Ala	Asp	Asp	Gln	Leu	Ile	Ile
			645					650					655		
Asp	Phe	Asp	Asn	Phe	Val	Arg	Cys	Leu	Val	Arg	Leu	Glu	Thr	Leu	Phe
			660				665					670			
Lys	Ile	Phe	Lys	Gln	Leu	Asp	Pro	Glu	Asn	Thr	Gly	Thr	Ile	Glu	Leu
	675					680						685			
Asp	Leu	Ile	Ser	Trp	Leu	Cys	Phe	Ser	Val	Leu					
	690				695										

<210> 24  
 <211> 221  
 <212> PRT  
 <213> Tobacco etch virus

<400> 24

Gly	Glu	Ser	Leu	Phe	Lys	Gly	Pro	Arg	Asp	Tyr	Asn	Pro	Ile	Ser	Ser
1			5						10					15	
Thr	Ile	Cys	His	Leu	Thr	Asn	Glu	Ser	Asp	Gly	His	Thr	Thr	Ser	Leu
			20					25					30		
Tyr	Gly	Ile	Gly	Phe	Gly	Pro	Phe	Ile	Ile	Thr	Asn	Lys	His	Leu	Phe
	35					40					45				
Arg	Arg	Asn	Asn	Gly	Thr	Leu	Leu	Val	Gln	Ser	Leu	His	Gly	Val	Phe
	50				55					60					
Lys	Val	Lys	Asn	Thr	Thr	Leu	Gln	Gln	His	Leu	Ile	Asp	Gly	Arg	
65				70					75					80	
Asp	Met	Ile	Ile	Ile	Arg	Met	Pro	Lys	Asp	Phe	Pro	Pro	Phe	Pro	Gln
			85					90					95		
Lys	Leu	Lys	Phe	Arg	Glu	Pro	Gln	Arg	Glu	Glu	Arg	Ile	Cys	Leu	Val
			100				105					110			
Thr	Thr	Asn	Phe	Gln	Thr	Lys	Ser	Met	Ser	Ser	Met	Val	Ser	Asp	Thr
		115				120						125			
Ser	Cys	Thr	Phe	Pro	Ser	Ser	Asp	Gly	Ile	Phe	Trp	Lys	His	Trp	Ile
	130					135					140				

Gln Thr Lys Asp Gly Gln Cys Gly Ser Pro Leu Val Ser Thr Arg Asp  
 145 150 155 160  
 Gly Phe Ile Val Gly Ile His Ser Ala Ser Asn Phe Thr Asn Thr Asn  
 165 170 175  
 Asn Tyr Phe Thr Ser Val Pro Lys Asn Phe Met Glu Leu Leu Thr Asn  
 180 185 190  
 Gln Glu Ala Gln Gln Trp Val Ser Gly Trp Arg Leu Asn Ala Asp Ser  
 195 200 205  
 Val Leu Trp Gly Gly His Lys Val Phe Met Asp Lys Pro  
 210 215 220

<210> 25  
 <211> 371  
 <212> PRT  
 <213> Streptococcus pyogenes

<400> 25  
 Asp Gln Asn Phe Ala Arg Asn Glu Lys Glu Ala Lys Asp Ser Ala Ile  
 1 5 10 15  
 Thr Phe Ile Gln Lys Ser Ala Ala Ile Lys Ala Gly Ala Arg Ser Ala  
 20 25 30  
 Glu Asp Ile Lys Leu Asp Lys Val Asn Leu Gly Gly Glu Leu Ser Gly  
 35 40 45  
 Ser Asn Met Tyr Val Tyr Asn Ile Ser Thr Gly Gly Phe Val Ile Val  
 50 55 60  
 Ser Gly Asp Lys Arg Ser Pro Glu Ile Leu Gly Tyr Ser Thr Ser Gly  
 65 70 75 80  
 Ser Phe Asp Val Asn Gly Lys Glu Asn Ile Ala Ser Phe Met Glu Ser  
 85 90 95  
 Tyr Val Glu Gln Ile Lys Glu Asn Lys Lys Leu Asp Ser Thr Tyr Ala  
 100 105 110  
 Gly Thr Ala Glu Ile Lys Gln Pro Val Val Lys Ser Leu Leu Asp Ser  
 115 120 125  
 Lys Gly Ile His Tyr Asn Gln Gly Asn Pro Tyr Asn Leu Leu Thr Pro  
 130 135 140  
 Val Ile Glu Lys Val Lys Pro Gly Glu Gln Ser Phe Val Gly Gln His  
 145 150 155 160  
 Ala Ala Thr Gly Ser Val Ala Thr Ala Thr Ala Gln Ile Met Lys Tyr  
 165 170 175  
 His Asn Tyr Pro Asn Lys Gly Leu Lys Asp Tyr Thr Tyr Thr Leu Ser  
 180 185 190  
 Ser Asn Asn Pro Tyr Phe Asn His Pro Lys Asn Leu Phe Ala Ala Ile  
 195 200 205  
 Ser Thr Arg Gln Tyr Asn Trp Asn Asn Ile Leu Pro Thr Tyr Ser Gly  
 210 215 220  
 Arg Glu Ser Asn Val Gln Lys Met Ala Ile Ser Glu Leu Met Ala Asp  
 225 230 235 240  
 Val Gly Ile Ser Val Asp Met Asp Tyr Gly Pro Ser Ser Gly Ser Ala  
 245 250 255  
 Gly Ser Ser Arg Val Gln Arg Ala Leu Lys Glu Asn Phe Gly Tyr Asn  
 260 265 270  
 Gln Ser Val His Gln Ile Asn Arg Gly Asp Phe Ser Lys Gln Asp Trp  
 275 280 285  
 Glu Ala Gln Ile Asp Lys Glu Leu Ser Gln Asn Gln Pro Val Tyr Tyr  
 290 295 300  
 Gln Gly Val Gly Lys Val Gly Gly His Ala Phe Val Ile Asp Gly Ala  
 305 310 315 320  
 Asp Gly Arg Asn Phe Tyr His Val Asn Trp Gly Trp Gly Gly Val Ser  
 325 330 335  
 Asp Gly Phe Phe Arg Leu Asp Ala Leu Asn Pro Ser Ala Leu Gly Thr  
 340 345 350  
 Gly Gly Gly Ala Gly Gly Phe Asn Gly Tyr Gln Ser Ala Val Val Gly  
 355 360 365

Ile Lys Pro  
370

<210> 26  
<211> 353  
<212> PRT  
<213> Homo sapiens

<400> 26  
Lys Lys His Thr Gly Tyr Val Gly Leu Lys Asn Gln Gly Ala Thr Cys  
1 5 10 15  
Tyr Met Asn Ser Leu Leu Gln Thr Leu Phe Phe Thr Asn Gln Leu Arg  
20 25 30  
Lys Ala Val Tyr Met Met Pro Thr Glu Gly Asp Asp Ser Ser Lys Ser  
35 40 45  
Val Pro Leu Ala Leu Gln Arg Val Phe Tyr Glu Leu Gln His Ser Asp  
50 55 60  
Lys Pro Val Gly Thr Lys Lys Leu Thr Lys Ser Phe Gly Trp Glu Thr  
65 70 75 80  
Leu Asp Ser Phe Met Gln His Asp Val Gln Glu Leu Cys Arg Val Leu  
85 90 95  
Leu Asp Asn Val Glu Asn Lys Met Lys Gly Thr Cys Val Glu Gly Thr  
100 105 110  
Ile Pro Lys Leu Phe Arg Gly Lys Met Val Ser Tyr Ile Gln Cys Lys  
115 120 125  
Glu Val Asp Tyr Arg Ser Asp Arg Arg Glu Asp Tyr Tyr Asp Ile Gln  
130 135 140  
Leu Ser Ile Lys Gly Lys Lys Asn Ile Phe Glu Ser Phe Val Asp Tyr  
145 150 155 160  
Val Ala Val Glu Gln Leu Asp Gly Asp Asn Lys Tyr Asp Ala Gly Glu  
165 170 175  
His Gly Leu Gln Glu Ala Glu Lys Gly Val Lys Phe Leu Thr Leu Pro  
180 185 190  
Pro Val Leu His Leu Gln Leu Met Arg Phe Met Tyr Asp Pro Gln Thr  
195 200 205  
Asp Gln Asn Ile Lys Ile Asn Asp Arg Phe Glu Phe Pro Glu Gln Leu  
210 215 220  
Pro Leu Asp Glu Phe Leu Gln Lys Thr Asp Pro Lys Asp Pro Ala Asn  
225 230 235 240  
Tyr Ile Leu His Ala Val Leu Val His Ser Gly Asp Asn His Gly Gly  
245 250 255  
His Tyr Val Val Tyr Leu Asn Pro Lys Gly Asp Gly Lys Trp Cys Lys  
260 265 270  
Phe Asp Asp Asp Val Val Ser Arg Cys Thr Lys Glu Glu Ala Ile Glu  
275 280 285  
His Asn Tyr Gly Gly His Asp Asp Asp Leu Ser Val Arg His Cys Thr  
290 295 300  
Asn Ala Tyr Met Leu Val Tyr Ile Arg Glu Ser Lys Leu Ser Glu Val  
305 310 315 320  
Leu Gln Ala Val Thr Asp His Asp Ile Pro Gln Gln Leu Val Glu Arg  
325 330 335  
Leu Gln Glu Glu Lys Arg Ile Glu Ala Gln Lys Arg Lys Glu Arg Gln  
340 345 350  
Glu

<210> 27  
<211> 174  
<212> PRT  
<213> Staphylococcus aureus

&lt;400&gt; 27

```

Tyr Asn Glu Gln Tyr Val Asn Lys Leu Glu Asn Phe Lys Ile Arg Glu
1      5      10      15
Thr Gln Gly Asn Asn Gly Trp Cys Ala Gly Tyr Thr Met Ser Ala Leu
      20      25      30
Leu Asn Ala Thr Tyr Asn Thr Asn Lys Tyr His Ala Glu Ala Val Met
      35      40      45
Arg Phe Leu His Pro Asn Leu Gln Gly Gln Gln Phe Gln Phe Thr Gly
      50      55      60
Leu Thr Pro Arg Glu Met Ile Tyr Phe Gly Gln Thr Gln Gly Arg Ser
65      70      75      80
Pro Gln Leu Leu Asn Arg Met Thr Thr Tyr Asn Glu Val Asp Asn Leu
      85      90      95
Thr Lys Asn Asn Lys Gly Ile Ala Ile Leu Gly Ser Arg Val Glu Ser
      100     105     110
Arg Asn Gly Met His Ala Gly His Ala Met Ala Val Val Gly Asn Ala
      115     120     125
Lys Leu Asn Asn Gly Gln Glu Val Ile Ile Ile Trp Asn Pro Trp Asp
      130     135     140
Asn Gly Phe Met Thr Gln Asp Ala Lys Asn Asn Val Ile Pro Val Ser
145     150     155     160
Asn Gly Asp His Tyr Gln Trp Tyr Ser Ser Ile Tyr Gly Tyr
      165     170

```

&lt;210&gt; 28

&lt;211&gt; 221

&lt;212&gt; PRT

<213> *Saccharomyces cerevisiae*

&lt;400&gt; 28

```

Gly Ser Leu Val Pro Glu Leu Asn Glu Lys Asp Asp Asp Gln Val Gln
1      5      10      15
Lys Ala Leu Ala Ser Arg Glu Asn Thr Gln Leu Met Asn Arg Asp Asn
      20      25      30
Ile Glu Ile Thr Val Arg Asp Phe Lys Thr Leu Ala Pro Arg Arg Trp
      35      40      45
Leu Asn Asp Thr Ile Ile Glu Phe Phe Met Lys Tyr Ile Glu Lys Ser
      50      55      60
Thr Pro Asn Thr Val Ala Phe Asn Ser Phe Phe Tyr Thr Asn Leu Ser
65      70      75      80
Glu Arg Gly Tyr Gln Gly Val Arg Arg Trp Met Lys Arg Lys Lys Thr
      85      90      95
Gln Ile Asp Lys Leu Asp Lys Ile Phe Thr Pro Ile Asn Leu Asn Gln
      100     105     110
Ser His Trp Ala Leu Gly Ile Ile Asp Leu Lys Lys Lys Thr Ile Gly
      115     120     125
Tyr Val Asp Ser Leu Ser Asn Gly Pro Asn Ala Met Ser Phe Ala Ile
      130     135     140
Leu Thr Asp Leu Gln Lys Tyr Val Met Glu Glu Ser Lys His Thr Ile
145     150     155     160
Gly Glu Asp Phe Asp Leu Ile His Leu Asp Cys Pro Gln Gln Pro Asn
      165     170     175
Gly Tyr Asp Cys Gly Ile Tyr Val Cys Met Asn Thr Leu Tyr Gly Ser
      180     185     190
Ala Asp Ala Pro Leu Asp Phe Asp Tyr Lys Asp Ala Ile Arg Met Arg
      195     200     205
Arg Phe Ile Ala His Leu Ile Leu Thr Asp Ala Leu Lys
      210     215     220

```

&lt;210&gt; 29

&lt;211&gt; 166

&lt;212&gt; PRT

<213> *Pyrococcus horikoshii*

&lt;400&gt; 29

```

Met Lys Val Leu Phe Leu Thr Ala Asn Glu Phe Glu Asp Val Glu Leu
1      5      10
Ile Tyr Pro Tyr His Arg Leu Lys Glu Gly His Glu Val Tyr Ile
      20      25      30
Ala Ser Phe Glu Arg Gly Thr Ile Thr Gly Lys His Gly Tyr Ser Val
      35      40      45
Lys Val Asp Leu Thr Phe Asp Lys Val Asn Pro Glu Glu Phe Asp Ala
50      55      60
Leu Val Leu Pro Gly Gly Arg Ala Pro Glu Arg Val Arg Leu Asn Glu
65      70      75      80
Lys Ala Val Ser Ile Ala Arg Lys Met Phe Ser Glu Gly Lys Pro Val
      85      90      95
Ala Ser Ile Cys His Gly Pro Gln Ile Leu Ile Ser Ala Gly Val Leu
      100      105      110
Arg Gly Arg Lys Gly Thr Ser Tyr Pro Gly Ile Lys Asp Asp Met Ile
      115      120      125
Asn Ala Gly Val Glu Trp Val Asp Ala Glu Val Val Val Asp Gly Asn
      130      135      140
Trp Val Ser Ser Arg Val Pro Ala Asp Leu Tyr Ala Trp Met Arg Glu
145      150      155      160
Phe Val Lys Leu Leu Lys
      165

```

&lt;210&gt; 30

&lt;211&gt; 316

&lt;212&gt; PRT

<213> *Bacillus thermoproteolyticus*

&lt;400&gt; 30

```

Ile Thr Gly Thr Ser Thr Val Gly Val Gly Arg Gly Val Leu Gly Asp
1      5      10
Gln Lys Asn Ile Asn Thr Thr Tyr Ser Thr Tyr Tyr Tyr Leu Gln Asp
      20      25      30
Asn Thr Arg Gly Asp Gly Ile Phe Thr Tyr Asp Ala Lys Tyr Arg Thr
      35      40      45
Thr Leu Pro Gly Ser Leu Trp Ala Asp Ala Asp Asn Gln Phe Phe Ala
50      55      60
Ser Tyr Asp Ala Pro Ala Val Asp Ala His Tyr Tyr Ala Gly Val Thr
65      70      75      80
Tyr Asp Tyr Tyr Lys Asn Val His Asn Arg Leu Ser Tyr Asp Gly Asn
      85      90      95
Asn Ala Ala Ile Arg Ser Ser Val His Tyr Ser Gln Gly Tyr Asn Asn
      100      105      110
Ala Phe Trp Asn Gly Ser Glu Met Val Tyr Gly Asp Gly Asp Gly Gln
      115      120      125
Thr Phe Ile Pro Leu Ser Gly Gly Ile Asp Val Val Ala His Glu Leu
      130      135      140
Thr His Ala Val Thr Asp Tyr Thr Ala Gly Leu Ile Tyr Gln Asn Glu
145      150      155      160
Ser Gly Ala Ile Asn Glu Ala Ile Ser Asp Ile Phe Gly Thr Leu Val
      165      170      175
Glu Phe Tyr Ala Asn Lys Asn Pro Asp Trp Glu Ile Gly Glu Asp Val
      180      185      190
Tyr Thr Pro Gly Ile Ser Gly Asp Ser Leu Arg Ser Met Ser Asp Pro
      195      200      205
Ala Lys Tyr Gly Asp Pro Asp His Tyr Ser Lys Arg Tyr Thr Gly Thr
      210      215      220
Gln Asp Asn Gly Gly Val His Ile Asn Ser Gly Ile Ile Asn Lys Ala

```

```

225          230          235          240
Ala Tyr Leu Ile Ser Gln Gly Gly Thr His Tyr Gly Val Ser Val Val
          245          250          255
Gly Ile Gly Arg Asp Lys Leu Gly Lys Ile Phe Tyr Arg Ala Leu Thr
          260          265          270
Gln Tyr Leu Thr Pro Thr Ser Asn Phe Ser Gln Leu Arg Ala Ala Ala
          275          280          285
Val Gln Ser Ala Thr Asp Leu Tyr Gly Ser Thr Ser Gln Glu Val Ala
          290          295          300
Ser Val Lys Gln Ala Phe Asp Ala Val Gly Val Lys
305          310          315

```

```

<210> 31
<211> 169
<212> PRT
<213> Homo sapiens

```

```

<400> 31
Val Leu Thr Glu Gly Asn Pro Arg Trp Glu Gln Thr His Leu Thr Tyr
1          5          10          15
Arg Ile Glu Asn Tyr Thr Pro Asp Leu Pro Arg Ala Asp Val Asp His
          20          25          30
Ala Ile Glu Lys Ala Phe Gln Leu Trp Ser Asn Val Thr Pro Leu Thr
          35          40          45
Phe Thr Lys Val Ser Glu Gly Gln Ala Asp Ile Met Ile Ser Phe Val
          50          55          60
Arg Gly Asp His Arg Asp Asn Ser Pro Phe Asp Gly Pro Gly Gly Asn
65          70          75          80
Leu Ala His Ala Phe Gln Pro Gly Pro Gly Ile Gly Gly Asp Ala His
          85          90          95
Phe Asp Glu Asp Glu Arg Trp Thr Asn Asn Phe Arg Glu Tyr Asn Leu
          100          105          110
His Arg Val Ala Ala His Glu Leu Gly His Ser Leu Gly Leu Ser His
          115          120          125
Ser Thr Asp Ile Gly Ala Leu Met Tyr Pro Ser Tyr Thr Phe Ser Gly
          130          135          140
Asp Val Gln Leu Ala Gln Asp Asp Ile Asp Gly Ile Gln Ala Ile Tyr
145          150          155          160
Gly Arg Ser Gln Asn Pro Val Gln Pro
          165

```

```

<210> 32
<211> 496
<212> PRT
<213> Homo sapiens

```

```

<400> 32
Gln Tyr Ser Pro Asn Thr Gln Gln Gly Arg Thr Ser Ile Val His Leu
1          5          10          15
Phe Glu Trp Arg Trp Val Asp Ile Ala Leu Glu Cys Glu Arg Tyr Leu
          20          25          30
Ala Pro Lys Gly Phe Gly Gly Val Gln Val Ser Pro Pro Asn Glu Asn
          35          40          45
Val Ala Ile Tyr Asn Pro Phe Arg Pro Trp Trp Glu Arg Tyr Gln Pro
          50          55          60
Val Ser Tyr Lys Leu Cys Thr Arg Ser Gly Asn Glu Asp Glu Phe Arg
65          70          75          80
Asn Met Val Thr Arg Cys Asn Asn Val Gly Val Arg Ile Tyr Val Asp
          85          90          95
Ala Val Ile Asn His Met Cys Gly Asn Ala Val Ser Ala Gly Thr Ser
          100          105          110

```



Ser Thr Cys Gly Ser Tyr Phe Asn Pro Gly Ser Arg Asp Phe Pro Ala  
 115 120 125  
 Val Pro Tyr Ser Gly Trp Asp Phe Asn Asp Gly Lys Cys Lys Thr Gly  
 130 135 140  
 Ser Gly Asp Ile Glu Asn Tyr Asn Asp Ala Thr Gln Val Arg Asp Cys  
 145 150 155 160  
 Arg Leu Thr Gly Leu Leu Asp Leu Ala Leu Glu Lys Asp Tyr Val Arg  
 165 170 175  
 Ser Lys Ile Ala Glu Tyr Met Asn His Leu Ile Asp Ile Gly Val Ala  
 180 185 190  
 Gly Phe Arg Leu Asp Ala Ser Lys His Met Trp Pro Gly Asp Ile Lys  
 195 200 205  
 Ala Ile Leu Asp Lys Leu His Asn Leu Asn Ser Asn Trp Phe Pro Ala  
 210 215 220  
 Gly Ser Lys Pro Phe Ile Tyr Gln Glu Val Ile Asp Leu Gly Gly Glu  
 225 230 235 240  
 Pro Ile Lys Ser Ser Asp Tyr Phe Gly Asn Gly Arg Val Thr Glu Phe  
 245 250 255  
 Lys Tyr Gly Ala Lys Leu Gly Thr Val Ile Arg Lys Trp Asn Gly Glu  
 260 265 270  
 Lys Met Ser Tyr Leu Lys Asn Trp Gly Glu Gly Trp Gly Phe Val Pro  
 275 280 285  
 Ser Asp Arg Ala Leu Val Phe Val Asp Asn His Asp Asn Gln Arg Gly  
 290 295 300  
 His Gly Ala Gly Gly Ala Ser Ile Leu Thr Phe Trp Asp Ala Arg Leu  
 305 310 315 320  
 Tyr Lys Met Ala Val Gly Phe Met Leu Ala His Pro Tyr Gly Phe Thr  
 325 330 335  
 Arg Val Met Ser Ser Tyr Arg Trp Pro Arg Gln Phe Gln Asn Gly Asn  
 340 345 350  
 Asp Val Asn Asp Trp Val Gly Pro Pro Asn Asn Asn Gly Val Ile Lys  
 355 360 365  
 Glu Val Thr Ile Asn Pro Asp Thr Thr Cys Gly Asn Asp Trp Val Cys  
 370 375 380  
 Glu His Arg Trp Arg Gln Ile Arg Asn Met Val Ile Phe Arg Asn Val  
 385 390 395 400  
 Val Asp Gly Gln Pro Phe Thr Asn Trp Tyr Asp Asn Gly Ser Asn Gln  
 405 410 415  
 Val Ala Phe Gly Arg Gly Asn Arg Gly Phe Ile Val Phe Asn Asn Asp  
 420 425 430  
 Asp Trp Ser Phe Ser Leu Thr Leu Gln Thr Gly Leu Pro Ala Gly Thr  
 435 440 445  
 Tyr Cys Asp Val Ile Ser Gly Asp Lys Ile Asn Gly Asn Cys Thr Gly  
 450 455 460  
 Ile Lys Ile Tyr Val Ser Asp Asp Gly Lys Ala His Phe Ser Ile Ser  
 465 470 475 480  
 Asn Ser Ala Glu Asp Pro Phe Ile Ala Ile His Ala Glu Ser Lys Leu  
 485 490 495

<210> 33  
 <211> 370  
 <212> PRT  
 <213> Trichoderma reesei

<400> 33  
 Gln Pro Gly Thr Ser Thr Pro Glu Val His Pro Lys Leu Thr Thr Tyr  
 1 5 10 15  
 Lys Cys Thr Lys Ser Gly Gly Cys Val Ala Gln Asp Thr Ser Val Val  
 20 25 30  
 Leu Asp Trp Asn Tyr Arg Trp Met His Asp Ala Asn Tyr Asn Ser Cys  
 35 40 45  
 Thr Val Asn Gly Gly Val Asn Thr Thr Leu Cys Pro Asp Glu Ala Thr

50		55		60
Cys Gly Lys Asn Cys Phe Ile Glu Gly Val Asp Tyr Ala Ala Ser Gly				
65		70		75
Val Thr Thr Ser Gly Ser Ser Leu Thr Met Asn Gln Tyr Met Pro Ser				80
	85		90	95
Ser Ser Gly Gly Tyr Ser Ser Val Ser Pro Arg Leu Tyr Leu Leu Asp				
	100		105	110
Ser Asp Gly Glu Tyr Val Met Leu Lys Leu Asn Gly Gln Glu Leu Ser				
	115		120	125
Phe Asp Val Asp Leu Ser Ala Leu Pro Cys Gly Glu Asn Gly Ser Leu				
	130		135	140
Tyr Leu Ser Gln Met Asp Glu Asn Gly Gly Ala Asn Gln Tyr Asn Thr				
	145		150	155
Ala Gly Ala Asn Tyr Gly Ser Gly Tyr Cys Asp Ala Gln Cys Pro Val				
	165		170	175
Gln Thr Trp Arg Asn Gly Thr Leu Asn Thr Ser His Gln Gly Phe Cys				
	180		185	190
Cys Asn Glu Met Asp Ile Leu Glu Gly Asn Ser Arg Ala Asn Ala Leu				
	195		200	205
Thr Pro His Ser Cys Thr Ala Thr Ala Cys Asp Ser Ala Gly Cys Gly				
	210		215	220
Phe Asn Pro Tyr Gly Ser Gly Tyr Lys Ser Tyr Tyr Gly Pro Gly Asp				
	225		230	235
Thr Val Asp Thr Ser Lys Thr Phe Thr Ile Ile Thr Gln Phe Asn Thr				
	245		250	255
Asp Asn Gly Ser Pro Ser Gly Asn Leu Val Ser Ile Thr Arg Lys Tyr				
	260		265	270
Gln Gln Asn Gly Val Asp Ile Pro Ser Ala Gln Pro Gly Gly Asp Thr				
	275		280	285
Ile Ser Ser Cys Pro Ser Ala Ser Ala Tyr Gly Gly Leu Ala Thr Met				
	290		295	300
Gly Lys Ala Leu Ser Ser Gly Met Val Leu Val Phe Ser Ile Trp Asn				
	305		310	315
Asp Asn Ser Gln Tyr Met Asn Trp Leu Asp Ser Gly Asn Ala Gly Pro				
	325		330	335
Cys Ser Ser Thr Glu Gly Asn Pro Ser Asn Ile Leu Ala Asn Asn Pro				
	340		345	350
Asn Thr His Val Val Phe Ser Asn Ile Arg Trp Gly Asp Ile Gly Ser				
	355		360	365
Thr Thr				
370				

<210> 34  
 <211> 223  
 <212> PRT  
 <213> Aspergillus niger

<400> 34  
 Gln Thr Met Cys Ser Gln Tyr Asp Ser Ala Ser Ser Pro Pro Tyr Ser  
 1 5 10 15  
 Val Asn Gln Asn Leu Trp Gly Glu Tyr Gln Gly Thr Gly Ser Gln Cys  
 20 25 30  
 Val Tyr Val Asp Lys Leu Ser Ser Gly Ala Ser Trp His Thr Glu  
 35 40 45  
 Trp Thr Trp Ser Gly Gly Glu Gly Thr Val Lys Ser Tyr Ser Asn Ser  
 50 55 60  
 Gly Val Thr Phe Asn Lys Lys Leu Val Ser Asp Val Ser Ser Ile Pro  
 65 70 75 80  
 Thr Ser Val Glu Trp Lys Gln Asp Asn Thr Asn Val Asn Ala Asp Val  
 85 90 95  
 Ala Tyr Asp Leu Phe Thr Ala Ala Asn Val Asp His Ala Thr Ser Ser  
 100 105 110

Gly Asp Tyr Glu Leu Met Ile Trp Leu Ala Arg Tyr Gly Asn Ile Gln  
           115                          120                          125  
 Pro Ile Gly Lys Gln Ile Ala Thr Ala Thr Val Gly Gly Lys Ser Trp  
           130                          135                          140  
 Glu Val Trp Tyr Gly Ser Thr Thr Gln Ala Gly Ala Glu Gln Arg Thr  
 145                          150                          155                          160  
 Tyr Ser Phe Val Ser Glu Ser Pro Ile Asn Ser Tyr Ser Gly Asp Ile  
                           165                          170                          175  
 Asn Ala Phe Phe Ser Tyr Leu Thr Gln Asn Gln Gly Phe Pro Ala Ser  
                           180                          185                          190  
 Ser Gln Tyr Leu Ile Asn Leu Gln Phe Gly Thr Glu Ala Phe Thr Gly  
           195                          200                          205  
 Gly Pro Ala Thr Phe Thr Val Asp Asn Trp Thr Ala Ser Val Asn  
           210                          215                          220

<210> 35  
 <211> 184  
 <212> PRT  
 <213> *Aspergillus niger*

<400> 35  
 Ser Ala Gly Ile Asn Tyr Val Gln Asn Tyr Asn Gly Asn Leu Gly Asp  
 1                          5                          10                          15  
 Phe Thr Tyr Asp Glu Ser Ala Gly Thr Phe Ser Met Tyr Trp Glu Asp  
                           20                          25                          30  
 Gly Val Ser Ser Asp Phe Val Val Gly Leu Gly Trp Thr Thr Gly Ser  
           35                          40                          45  
 Ser Asn Ala Ile Thr Tyr Ser Ala Glu Tyr Ser Ala Ser Gly Ser Ala  
 50                          55                          60  
 Ser Tyr Leu Ala Val Tyr Gly Trp Val Asn Tyr Pro Gln Ala Glu Tyr  
 65                          70                          75                          80  
 Tyr Ile Val Glu Asp Tyr Gly Asp Tyr Asn Pro Cys Ser Ser Ala Thr  
                           85                          90                          95  
 Ser Leu Gly Thr Val Tyr Ser Asp Gly Ser Thr Tyr Gln Val Cys Thr  
           100                          105                          110  
 Asp Thr Arg Thr Asn Glu Pro Ser Ile Thr Gly Thr Ser Thr Phe Thr  
           115                          120                          125  
 Gln Tyr Phe Ser Val Arg Glu Ser Thr Arg Thr Ser Gly Thr Val Thr  
           130                          135                          140  
 Val Ala Asn His Phe Asn Phe Trp Ala His His Gly Phe Gly Asn Ser  
 145                          150                          155                          160  
 Asp Phe Asn Tyr Gln Val Val Ala Val Glu Ala Trp Ser Gly Ala Gly  
                           165                          170                          175  
 Ser Ala Ser Val Thr Ile Ser Ser  
           180

<210> 36  
 <211> 313  
 <212> PRT  
 <213> *Streptomyces lividans*

<400> 36  
 Ala Glu Ser Thr Leu Gly Ala Ala Ala Ala Gln Ser Gly Arg Tyr Phe  
 1                          5                          10                          15  
 Gly Thr Ala Ile Ala Ser Gly Arg Leu Ser Asp Ser Thr Tyr Thr Ser  
           20                          25                          30  
 Ile Ala Gly Arg Glu Phe Asn Met Val Thr Ala Glu Asn Glu Met Lys  
           35                          40                          45  
 Ile Asp Ala Thr Glu Pro Gln Arg Gly Gln Phe Asn Phe Ser Ser Ala  
 50                          55                          60  
 Asp Arg Val Tyr Asn Trp Ala Val Gln Asn Gly Lys Gln Val Arg Gly

65					70					75					80
His	Thr	Leu	Ala	Trp	His	Ser	Gln	Gln	Pro	Gly	Trp	Met	Gln	Ser	Leu
				85					90					95	
Ser	Gly	Ser	Ala	Leu	Arg	Gln	Ala	Met	Ile	Asp	His	Ile	Asn	Gly	Val
			100					105					110		
Met	Ala	His	Tyr	Lys	Gly	Lys	Ile	Val	Gln	Trp	Asp	Val	Val	Asn	Glu
			115				120					125			
Ala	Phe	Ala	Asp	Gly	Ser	Ser	Gly	Ala	Arg	Arg	Asp	Ser	Asn	Leu	Gln
			130				135				140				
Arg	Ser	Gly	Asn	Asp	Trp	Ile	Glu	Val	Ala	Phe	Arg	Thr	Ala	Arg	Ala
			145				150				155				160
Ala	Asp	Pro	Ser	Ala	Lys	Leu	Cys	Tyr	Asn	Asp	Tyr	Asn	Val	Glu	Asn
				165					170					175	
Trp	Thr	Trp	Ala	Lys	Thr	Gln	Ala	Met	Tyr	Asn	Met	Val	Arg	Asp	Phe
			180					185					190		
Lys	Gln	Arg	Gly	Val	Pro	Ile	Asp	Cys	Val	Gly	Phe	Gln	Ser	His	Phe
			195				200					205			
Asn	Ser	Gly	Ser	Pro	Tyr	Asn	Ser	Asn	Phe	Arg	Thr	Thr	Leu	Gln	Asn
			210				215				220				
Phe	Ala	Ala	Leu	Gly	Val	Asp	Val	Ala	Ile	Thr	Glu	Leu	Asp	Ile	Gln
			225				230				235				240
Gly	Ala	Pro	Ala	Ser	Thr	Tyr	Ala	Asn	Val	Thr	Asn	Asp	Cys	Leu	Ala
				245				250					255		
Val	Ser	Arg	Cys	Leu	Gly	Ile	Thr	Val	Trp	Gly	Val	Arg	Asp	Ser	Asp
			260					265					270		
Ser	Trp	Arg	Ser	Glu	Gln	Thr	Pro	Leu	Leu	Phe	Asn	Asn	Asp	Gly	Ser
			275				280					285			
Lys	Lys	Ala	Ala	Tyr	Thr	Ala	Val	Leu	Asp	Ala	Leu	Asn	Gly	Gly	Ala
			290				295				300				
Ser	Ser	Glu	Pro	Pro	Ala	Asp	Gly	Gly							
			305				310								

<210> 37  
 <211> 362  
 <212> PRT  
 <213> *Aspergillus niger*

<400> 37  
 Met His Ser Phe Ala Ser Leu Leu Ala Tyr Gly Leu Val Ala Gly Ala  
 1 5 10 15  
 Thr Phe Ala Ser Ala Ser Pro Ile Glu Ala Arg Asp Ser Cys Thr Phe  
 20 25 30  
 Thr Thr Ala Ala Ala Ala Lys Ala Gly Lys Ala Lys Cys Ser Thr Ile  
 35 40 45  
 Thr Leu Asn Asn Ile Glu Val Pro Ala Gly Thr Thr Leu Asp Leu Thr  
 50 55 60  
 Gly Leu Thr Ser Gly Thr Lys Val Ile Phe Glu Gly Thr Thr Thr Phe  
 65 70 75 80  
 Gln Tyr Glu Glu Trp Ala Gly Pro Leu Ile Ser Met Ser Gly Glu His  
 85 90 95  
 Ile Thr Val Thr Gly Ala Ser Gly His Leu Ile Asn Cys Asp Gly Ala  
 100 105 110  
 Arg Trp Trp Asp Gly Lys Gly Thr Ser Gly Lys Lys Lys Pro Lys Phe  
 115 120 125  
 Phe Tyr Ala His Gly Leu Asp Ser Ser Ser Ile Thr Gly Leu Asn Ile  
 130 135 140  
 Lys Asn Thr Pro Leu Met Ala Phe Ser Val Gln Ala Asn Asp Ile Thr  
 145 150 155 160  
 Phe Thr Asp Val Thr Ile Asn Asn Ala Asp Gly Asp Thr Gln Gly Gly  
 165 170 175  
 His Asn Thr Asp Ala Phe Asp Val Gly Asn Ser Val Gly Val Asn Ile  
 180 185 190

```

Ile Lys Pro Trp Val His Asn Gln Asp Asp Cys Leu Ala Val Asn Ser
    195                200                205
Gly Glu Asn Ile Trp Phe Thr Gly Gly Thr Cys Ile Gly Gly His Gly
    210                215                220
Leu Ser Ile Gly Ser Val Gly Asp Arg Ser Asn Asn Val Val Lys Asn
    225                230                235                240
Val Thr Ile Glu His Ser Thr Val Ser Asn Ser Glu Asn Ala Val Arg
    245                250                255
Ile Lys Thr Ile Ser Gly Ala Thr Gly Ser Val Ser Glu Ile Thr Tyr
    260                265                270
Ser Asn Ile Val Met Ser Gly Ile Ser Asp Tyr Gly Val Val Ile Gln
    275                280                285
Gln Asp Tyr Glu Asp Gly Lys Pro Thr Gly Lys Pro Thr Asn Gly Val
    290                295                300
Thr Ile Gln Asp Val Lys Leu Glu Ser Val Thr Gly Ser Val Asp Ser
    305                310                315                320
Gly Ala Thr Glu Ile Tyr Leu Leu Cys Gly Ser Gly Ser Cys Ser Asp
    325                330                335
Trp Thr Trp Asp Val Lys Val Thr Gly Gly Lys Lys Ser Thr Ala
    340                345                350
Cys Lys Asn Phe Pro Ser Val Ala Ser Cys
    355                360

```

```

<210> 38
<211> 383
<212> PRT
<213> Pseudomonas cellulosa

```

```

<400> 38
Arg Ala Asp Val Lys Pro Val Thr Val Lys Leu Val Asp Ser Gln Ala
1      5      10
Thr Met Glu Thr Arg Ser Leu Phe Ala Phe Met Gln Glu Gln Arg Arg
    20      25      30
His Ser Ile Met Phe Gly His Gln His Glu Thr Thr Gln Gly Leu Thr
    35      40      45
Ile Thr Arg Thr Asp Gly Thr Gln Ser Asp Thr Phe Asn Ala Val Gly
    50      55      60
Asp Phe Ala Ala Val Tyr Gly Trp Asp Thr Leu Ser Ile Val Ala Pro
    65      70      75      80
Lys Ala Glu Gly Asp Ile Val Ala Gln Val Lys Lys Ala Tyr Ala Arg
    85      90      95
Gly Gly Ile Ile Thr Val Ser Ser His Phe Asp Asn Pro Lys Thr Asp
    100     105     110
Thr Gln Lys Gly Val Trp Pro Val Gly Thr Ser Trp Asp Gln Thr Pro
    115     120     125
Ala Val Val Asp Ser Leu Pro Gly Gly Ala Tyr Asn Asn Pro Val Leu Asn
    130     135     140
Gly Tyr Leu Asp Gln Val Ala Glu Trp Ala Asn Asn Leu Lys Asp Glu
    145     150     155     160
Gln Gly Arg Leu Ile Pro Val Ile Phe Arg Leu Tyr His Ala Asn Thr
    165     170     175
Gly Ser Trp Phe Trp Trp Gly Asp Lys Gln Ser Thr Pro Glu Gln Tyr
    180     185     190
Lys Gln Leu Phe Arg Tyr Ser Val Glu Tyr Leu Arg Asp Val Lys Gly
    195     200     205
Val Arg Asn Phe Leu Tyr Ala Tyr Ser Pro Asn Asn Phe Trp Asp Val
    210     215     220
Thr Glu Ala Asn Tyr Leu Glu Arg Tyr Pro Gly Asp Glu Trp Val Asp
    225     230     235     240
Val Leu Gly Phe Asp Thr Tyr Gly Pro Val Ala Asp Asn Ala Asp Trp
    245     250     255
Phe Arg Asn Val Val Ala Asn Ala Ala Leu Val Ala Arg Met Ala Glu

```

Ala	Arg	Gly	Lys	Ile	Pro	Val	Ile	Ser	Glu	Ile	Gly	Ile	Arg	Ala	Pro
		275					280					285			
Asp	Ile	Glu	Ala	Gly	Leu	Tyr	Asp	Asn	Gln	Trp	Tyr	Arg	Lys	Leu	Ile
	290					295					300				
Ser	Gly	Leu	Lys	Ala	Asp	Pro	Asp	Ala	Arg	Glu	Ile	Ala	Phe	Leu	Leu
305					310					315					320
Val	Trp	Arg	Asn	Ala	Pro	Gln	Gly	Val	Pro	Gly	Pro	Asn	Gly	Thr	Gln
			325						330					335	
Val	Pro	His	Tyr	Trp	Val	Pro	Ala	Asn	Arg	Pro	Glu	Asn	Ile	Asn	Asn
			340					345					350		
Gly	Thr	Leu	Glu	Asp	Phe	Gln	Ala	Phe	Tyr	Ala	Asp	Glu	Phe	Thr	Ala
		355					360				365				
Phe	Asn	Arg	Asp	Ile	Glu	Gln	Val	Tyr	Gln	Arg	Pro	Thr	Leu	Ile	
	370					375					380				

```
<210> 39
<211> 419
<212> PRT
<213> Bacillus circulans
```

<400>	39														
Leu	Gln	Pro	Ala	Thr	Ala	Glu	Ala	Ala	Asp	Ser	Tyr	Lys	Ile	Val	Gly
1			5						10					15	
Tyr	Tyr	Pro	Ser	Trp	Ala	Ala	Tyr	Gly	Arg	Asn	Tyr	Asn	Val	Ala	Asp
			20					25					30		
Ile	Asp	Pro	Thr	Lys	Val	Thr	His	Ile	Asn	Tyr	Ala	Phe	Ala	Asp	Ile
		35					40				45				
Cys	Trp	Asn	Gly	Ile	His	Gly	Asn	Pro	Asp	Pro	Ser	Gly	Pro	Asn	Pro
	50					55					60				
Val	Thr	Trp	Thr	Cys	Gln	Asn	Glu	Lys	Ser	Gln	Thr	Ile	Asn	Val	Pro
65				70					75					80	
Asn	Gly	Thr	Ile	Val	Leu	Gly	Asp	Pro	Trp	Ile	Asp	Thr	Gly	Lys	Thr
			85						90					95	
Phe	Ala	Gly	Asp	Thr	Trp	Asp	Gln	Pro	Ile	Ala	Gly	Asn	Ile	Asn	Gln
			100					105					110		
Leu	Asn	Lys	Leu	Lys	Gln	Thr	Asn	Pro	Asn	Leu	Lys	Thr	Ile	Ile	Ser
		115					120					125			
Val	Gly	Gly	Trp	Thr	Trp	Ser	Asn	Arg	Phe	Ser	Asp	Val	Ala	Ala	Thr
	130					135					140				
Ala	Ala	Thr	Arg	Glu	Val	Phe	Ala	Asn	Ser	Ala	Val	Asp	Phe	Leu	Arg
145				150					155					160	
Lys	Tyr	Asn	Phe	Asp	Gly	Val	Asp	Leu	Asp	Trp	Glu	Tyr	Pro	Val	Ser
			165					170					175		
Gly	Gly	Leu	Asp	Gly	Asn	Ser	Lys	Arg	Pro	Glu	Asp	Lys	Gln	Asn	Tyr
		180						185					190		
Thr	Leu	Leu	Leu	Ser	Lys	Ile	Arg	Glu	Lys	Leu	Asp	Ala	Ala	Gly	Ala
		195					200					205			
Val	Asp	Gly	Lys	Lys	Tyr	Leu	Leu	Thr	Ile	Ala	Ser	Gly	Ala	Ser	Ala
	210					215					220				
Thr	Tyr	Ala	Ala	Asn	Thr	Glu	Leu	Ala	Lys	Ile	Ala	Ala	Ile	Val	Asp
225				230					235					240	
Trp	Ile	Asn	Ile	Met	Thr	Tyr	Asp	Phe	Asn	Gly	Ala	Trp	Gln	Lys	Ile
			245					250					255		
Ser	Ala	His	Asn	Ala	Pro	Leu	Asn	Tyr	Asp	Pro	Ala	Ala	Ser	Ala	Ala
		260						265					270		
Gly	Val	Pro	Asp	Ala	Asn	Thr	Phe	Asn	Val	Ala	Ala	Gly	Ala	Gln	Gly
		275					280					285			
His	Leu	Asp	Ala	Gly	Val	Pro	Ala	Ala	Lys	Leu	Val	Leu	Gly	Val	Pro
	290					295					300				
Phe	Tyr	Gly	Arg	Gly	Trp	Asp	Gly	Cys	Ala	Gln	Ala	Gly	Asn	Gly	Gln
305				310					315					320	

[illegible]

```
<210> 40
<211> 317
<212> PRT
<213> Candida antarctica
```

<400>	40														
Leu	Pro	Ser	Gly	Ser	Asp	Pro	Ala	Phe	Ser	Gln	Pro	Lys	Ser	Val	Leu
1				5					10					15	
Asp	Ala	Gly	Leu	Thr	Cys	Gln	Gly	Ala	Ser	Pro	Ser	Ser	Val	Ser	Lys
			20					25					30		
Pro	Ile	Leu	Leu	Val	Pro	Gly	Thr	Gly	Thr	Thr	Gly	Pro	Gln	Ser	Phe
		35					40					45			
Asp	Ser	Asn	Trp	Ile	Pro	Leu	Ser	Thr	Gln	Leu	Gly	Tyr	Thr	Pro	Cys
	50					55					60				
Trp	Ile	Ser	Pro	Pro	Pro	Phe	Met	Leu	Asn	Asp	Thr	Gln	Val	Asn	Thr
65					70					75					80
Glu	Tyr	Met	Val	Asn	Ala	Ile	Thr	Ala	Leu	Tyr	Ala	Gly	Ser	Gly	Asn
				85					90					95	
Asn	Lys	Leu	Pro	Val	Leu	Thr	Trp	Ser	Gln	Gly	Gly	Leu	Val	Ala	Gln
			100					105					110		
Trp	Gly	Leu	Thr	Phe	Phe	Pro	Ser	Ile	Arg	Ser	Lys	Val	Asp	Arg	Leu
		115					120					125			
Met	Ala	Phe	Ala	Pro	Asp	Tyr	Lys	Gly	Thr	Val	Leu	Ala	Gly	Pro	Leu
	130					135					140				
Asp	Ala	Leu	Ala	Val	Ser	Ala	Pro	Ser	Val	Trp	Gln	Gln	Thr	Thr	Gly
145					150					155					160
Ser	Ala	Leu	Thr	Thr	Ala	Leu	Arg	Asn	Ala	Gly	Gly	Leu	Thr	Gln	Ile
				165					170					175	
Val	Pro	Thr	Thr	Asn	Leu	Tyr	Ser	Ala	Thr	Asp	Glu	Ile	Val	Gln	Pro
				180					185				190		
Gln	Val	Ser	Asn	Ser	Pro	Leu	Asp	Ser	Ser	Tyr	Leu	Phe	Asn	Gly	Lys
		195					200					205			
Asn	Val	Gln	Ala	Gln	Ala	Val	Cys	Gly	Pro	Leu	Phe	Val	Ile	Asp	His
	210					215					220				
Ala	Gly	Ser	Leu	Thr	Ser	Gln	Phe	Ser	Tyr	Val	Val	Gly	Arg	Ser	Ala
225					230					235					240
Leu	Arg	Ser	Thr	Thr	Gly	Gln	Ala	Arg	Ser	Ala	Asp	Tyr	Gly	Ile	Thr
				245					250					255	
Asp	Cys	Asn	Pro	Leu	Pro	Ala	Asn	Asp	Leu	Thr	Pro	Glu	Gln	Lys	Val
		260					265						270		
Ala	Ala	Ala	Ala	Leu	Leu	Ala	Pro	Ala	Ala	Ala	Ala	Ile	Val	Ala	Gly
		275					280					285			
Pro	Lys	Gln	Asn	Cys	Glu	Pro	Asp	Leu	Met	Pro	Tyr	Ala	Arg	Pro	Phe
	290					295					300				
Ala	Val	Gly	Lys	Arg	Thr	Cys	Ser	Gly	Ile	Val	Thr	Pro			
305					310					315					

<210> 41  
 <211> 434  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> chimera of guinea pig and homo sapiens (human= approx. last 30 amino acids)

<400> 41  
 Ala Glu Val Cys Tyr Ser His Leu Gly Cys Phe Ser Asp Glu Lys Pro  
 1 5 10 15  
 Trp Ala Gly Thr Ser Gln Arg Pro Ile Lys Ser Leu Pro Ser Asp Pro  
 20 25 30  
 Lys Lys Ile Asn Thr Arg Phe Leu Leu Tyr Thr Asn Glu Asn Gln Asn  
 35 40 45  
 Ser Tyr Gln Leu Ile Thr Ala Thr Asp Ile Ala Thr Ile Lys Ala Ser  
 50 55 60  
 Asn Phe Asn Leu Asn Arg Lys Thr Arg Phe Ile Ile His Gly Phe Thr  
 65 70 75 80  
 Asp Ser Gly Glu Asn Ser Trp Leu Ser Asp Met Cys Lys Asn Met Phe  
 85 90 95  
 Gln Val Glu Lys Val Asn Cys Ile Cys Val Asp Trp Lys Gly Gly Ser  
 100 105 110  
 Lys Ala Gln Tyr Ser Gln Ala Ser Gln Asn Ile Arg Val Val Gly Ala  
 115 120 125  
 Glu Val Ala Tyr Leu Val Gln Val Leu Ser Thr Ser Leu Asn Tyr Ala  
 130 135 140  
 Pro Glu Asn Val His Ile Ile Gly His Ser Leu Gly Ala His Thr Ala  
 145 150 155 160  
 Gly Glu Ala Gly Lys Arg Leu Asn Gly Leu Val Gly Arg Ile Thr Gly  
 165 170 175  
 Leu Asp Pro Ala Glu Pro Tyr Phe Gln Asp Thr Pro Glu Glu Val Arg  
 180 185 190  
 Leu Asp Pro Ser Asp Ala Lys Phe Val Asp Val Ile His Thr Asp Ile  
 195 200 205  
 Ser Pro Ile Leu Pro Ser Leu Gly Phe Gly Met Ser Gln Lys Val Gly  
 210 215 220  
 His Met Asp Phe Phe Pro Asn Gly Gly Lys Asp Met Pro Gly Cys Lys  
 225 230 235 240  
 Thr Gly Ile Ser Cys Asn His His Arg Ser Ile Glu Tyr Tyr His Ser  
 245 250 255  
 Ser Ile Leu Asn Pro Glu Gly Phe Leu Gly Tyr Pro Cys Ala Ser Tyr  
 260 265 270  
 Asp Glu Phe Gln Glu Ser Gly Cys Phe Pro Cys Pro Ala Lys Gly Cys  
 275 280 285  
 Pro Lys Met Gly His Phe Ala Asp Gln Tyr Pro Gly Lys Thr Asn Ala  
 290 295 300  
 Val Glu Gln Thr Phe Phe Leu Asn Thr Gly Ala Ser Asp Asn Phe Thr  
 305 310 315 320  
 Arg Trp Arg Tyr Lys Val Thr Val Thr Leu Ser Gly Glu Lys Asp Pro  
 325 330 335  
 Ser Gly Asn Ile Asn Val Ala Leu Leu Gly Lys Asn Gly Asn Ser Ala  
 340 345 350  
 Gln Tyr Gln Val Phe Lys Gly Thr Leu Lys Pro Asp Ala Ser Tyr Thr  
 355 360 365  
 Asn Ser Ile Asp Val Glu Leu Asn Val Gly Thr Ile Gln Lys Val Thr  
 370 375 380  
 Phe Leu Trp Lys Arg Ser Gly Ile Ser Val Ser Lys Pro Lys Met Gly  
 385 390 395 400  
 Ala Ser Arg Ile Thr Val Gln Ser Gly Lys Asp Gly Thr Lys Tyr Asn  
 405 410 415  
 Phe Cys Ser Ser Asp Ile Val Gln Glu Asn Val Glu Gln Thr Leu Ser



420                                      425                                      430  
 Pro Cys  
  
 <210> 42  
 <211> 471  
 <212> PRT  
 <213> Escherichia coli  
  
 <400> 42  
 Met Lys Gln Ser Thr Ile Ala Leu Ala Leu Leu Pro Leu Leu Phe Thr  
 1                                      5                                      10                                      15  
 Pro Val Thr Lys Ala Arg Thr Pro Glu Met Pro Val Leu Glu Asn Arg  
                                     20                                      25                                      30  
 Ala Ala Gln Gly Asp Ile Thr Ala Pro Gly Gly Ala Arg Arg Leu Thr  
                                     35                                      40                                      45  
 Gly Asp Gln Thr Ala Ala Leu Arg Asp Ser Leu Ser Asp Lys Pro Ala  
 50                                      55                                      60  
 Lys Asn Ile Ile Leu Leu Ile Gly Asp Gly Met Gly Asp Ser Glu Ile  
 65                                      70                                      75                                      80  
 Thr Ala Ala Arg Asn Tyr Ala Glu Gly Ala Gly Gly Phe Phe Lys Gly  
                                     85                                      90                                      95  
 Ile Asp Ala Leu Pro Leu Thr Gly Gln Tyr Thr His Tyr Ala Leu Asn  
                                     100                                      105                                      110  
 Lys Lys Thr Gly Lys Pro Asp Tyr Val Thr Asp Ser Ala Ala Ser Ala  
                                     115                                      120                                      125  
 Thr Ala Trp Ser Thr Gly Val Lys Thr Tyr Asn Gly Ala Leu Gly Val  
 130                                      135                                      140  
 Asp Ile His Glu Lys Asp His Pro Thr Ile Leu Glu Met Ala Lys Ala  
 145                                      150                                      155                                      160  
 Ala Gly Leu Ala Thr Gly Asn Val Ser Thr Ala Glu Leu Gln Asp Ala  
                                     165                                      170                                      175  
 Thr Pro Ala Ala Leu Val Ala His Val Thr Ser Arg Lys Cys Tyr Gly  
                                     180                                      185                                      190  
 Pro Ser Ala Thr Ser Glu Lys Cys Pro Gly Asn Ala Leu Glu Lys Gly  
                                     195                                      200                                      205  
 Gly Lys Gly Ser Ile Thr Glu Gln Leu Leu Asn Ala Arg Ala Asp Val  
                                     210                                      215                                      220  
 Thr Leu Gly Gly Gly Ala Lys Thr Phe Ala Glu Thr Ala Thr Ala Gly  
 225                                      230                                      235                                      240  
 Glu Trp Gln Gly Lys Thr Leu Arg Glu Gln Ala Gln Ala Arg Gly Tyr  
                                     245                                      250                                      255  
 Gln Leu Val Ser Asp Ala Ala Ser Leu Asn Ser Val Thr Glu Ala Asn  
                                     260                                      265                                      270  
 Gln Gln Lys Pro Leu Leu Gly Leu Phe Ala Asp Gly Asn Met Pro Val  
                                     275                                      280                                      285  
 Arg Trp Leu Gly Pro Lys Ala Thr Tyr His Gly Asn Ile Asp Lys Pro  
 290                                      295                                      300  
 Ala Val Thr Cys Thr Pro Asn Pro Gln Arg Asn Asp Ser Val Pro Thr  
 305                                      310                                      315                                      320  
 Leu Ala Gln Met Thr Asp Lys Ala Ile Glu Leu Leu Ser Lys Asn Glu  
                                     325                                      330                                      335  
 Lys Gly Phe Phe Leu Gln Val Glu Gly Ala Ser Ile Asp Lys Gln Asp  
                                     340                                      345                                      350  
 His Ala Ala Asn Pro Cys Gly Gln Ile Gly Glu Thr Val Asp Leu Asp  
                                     355                                      360                                      365  
 Glu Ala Val Gln Arg Ala Leu Glu Phe Ala Lys Lys Glu Gly Asn Thr  
 370                                      375                                      380  
 Leu Val Ile Val Thr Ala Asp His Ala His Ala Ser Gln Ile Val Ala  
 385                                      390                                      395                                      400  
 Pro Asp Thr Lys Ala Pro Gly Leu Thr Gln Ala Leu Asn Thr Lys Asp  
                                     405                                      410                                      415

Gly Ala Val Met Val Met Ser Tyr Gly Asn Ser Glu Glu Asp Ser Gln  
                           420                          425                          430  
 Glu His Thr Gly Ser Gln Leu Arg Ile Ala Ala Tyr Gly Pro His Ala  
                           435                          440                          445  
 Ala Asn Val Val Gly Leu Thr Asp Gln Thr Asp Leu Phe Tyr Thr Met  
                           450                          455                          460  
 Lys Ala Ala Leu Gly Leu Lys  
 465                          470

<210> 43  
 <211> 260  
 <212> PRT  
 <213> Bovine

<400> 43  
 Leu Lys Ile Ala Ala Phe Asn Ile Arg Thr Phe Gly Glu Thr Lys Met  
 1                          5                          10                          15  
 Ser Asn Ala Thr Leu Ala Ser Tyr Ile Val Arg Ile Val Arg Tyr  
                           20                          25                          30  
 Asp Ile Val Leu Ile Gln Glu Val Arg Asp Ser His Leu Val Ala Val  
                           35                          40                          45  
 Gly Lys Leu Leu Asp Tyr Leu Asn Gln Asp Asp Pro Asn Thr Tyr His  
                           50                          55                          60  
 Tyr Val Val Ser Glu Pro Leu Gly Arg Asn Ser Tyr Lys Glu Arg Tyr  
 65                          70                          75                          80  
 Leu Phe Leu Phe Arg Pro Asn Lys Val Ser Val Leu Asp Thr Tyr Gln  
                           85                          90                          95  
 Tyr Asp Asp Gly Cys Glu Ser Cys Gly Asn Asp Ser Phe Ser Arg Glu  
                           100                          105                          110  
 Pro Ala Val Val Lys Phe Ser Ser His Ser Thr Lys Val Lys Glu Phe  
                           115                          120                          125  
 Ala Ile Val Ala Leu His Ser Ala Pro Ser Asp Ala Val Ala Glu Ile  
                           130                          135                          140  
 Asn Ser Leu Tyr Asp Val Tyr Leu Asp Val Gln Gln Lys Trp His Leu  
 145                          150                          155                          160  
 Asn Asp Val Met Leu Met Gly Asp Phe Asn Ala Asp Cys Ser Tyr Val  
                           165                          170                          175  
 Thr Ser Ser Gln Trp Ser Ser Ile Arg Leu Arg Thr Ser Ser Thr Phe  
                           180                          185                          190  
 Gln Trp Leu Ile Pro Asp Ser Ala Asp Thr Thr Ala Thr Ser Thr Asn  
                           195                          200                          205  
 Cys Ala Tyr Asp Arg Ile Val Val Ala Gly Ser Leu Leu Gln Ser Ser  
                           210                          215                          220  
 Val Val Pro Gly Ser Ala Ala Pro Phe Asp Phe Gln Ala Ala Tyr Gly  
 225                          230                          235                          240  
 Leu Ser Asn Glu Met Ala Leu Ala Ile Ser Asp His Tyr Pro Val Glu  
                           245                          250                          255  
 Val Thr Leu Thr  
                           260

<210> 44  
 <211> 686  
 <212> PRT  
 <213> Bacillus circulans

<400> 44  
 Ala Pro Asp Thr Ser Val Ser Asn Lys Gln Asn Phe Ser Thr Asp Val  
 1                          5                          10                          15  
 Ile Tyr Gln Ile Phe Thr Asp Arg Phe Ser Asp Gly Asn Pro Ala Asn  
                           20                          25                          30  
 Asn Pro Thr Gly Ala Ala Phe Asp Gly Thr Cys Thr Asn Leu Arg Leu

		35					40				45				
Tyr	Cys	Gly	Gly	Asp	Trp	Gln	Gly	Ile	Ile	Asn	Lys	Ile	Asn	Asp	Gly
	50					55					60				
Tyr	Leu	Thr	Gly	Met	Gly	Val	Thr	Ala	Ile	Trp	Ile	Ser	Gln	Pro	Val
65					70					75					80
Glu	Asn	Ile	Tyr	Ser	Ile	Ile	Asn	Tyr	Ser	Gly	Val	Asn	Asn	Thr	Ala
				85					90					95	
Tyr	His	Gly	Tyr	Trp	Ala	Arg	Asp	Phe	Lys	Lys	Thr	Asn	Pro	Ala	Tyr
		100						105					110		
Gly	Thr	Ile	Ala	Asp	Phe	Gln	Asn	Leu	Ile	Ala	Ala	Ala	His	Ala	Lys
		115					120					125			
Asn	Ile	Lys	Val	Ile	Ile	Asp	Phe	Ala	Pro	Asn	His	Thr	Ser	Pro	Ala
	130					135					140				
Ser	Ser	Asp	Gln	Pro	Ser	Phe	Ala	Glu	Asn	Gly	Arg	Leu	Tyr	Asp	Asn
145					150					155					160
Gly	Thr	Leu	Leu	Gly	Gly	Tyr	Thr	Asn	Asp	Thr	Gln	Asn	Leu	Phe	His
				165					170					175	
His	Asn	Gly	Gly	Thr	Asp	Phe	Ser	Thr	Thr	Glu	Asn	Gly	Ile	Tyr	Lys
		180					185						190		
Asn	Leu	Tyr	Asp	Leu	Ala	Asp	Leu	Asn	His	Asn	Asn	Ser	Thr	Val	Asp
	195						200					205			
Val	Tyr	Leu	Lys	Asp	Ala	Ile	Lys	Met	Trp	Leu	Asp	Leu	Gly	Ile	Asp
	210					215					220				
Gly	Ile	Arg	Met	Asp	Ala	Val	Lys	His	Met	Pro	Phe	Gly	Trp	Gln	Lys
225					230					235					240
Ser	Phe	Met	Ala	Ala	Val	Asn	Asn	Tyr	Lys	Pro	Val	Phe	Thr	Phe	Gly
				245					250					255	
Glu	Trp	Phe	Leu	Gly	Val	Asn	Glu	Val	Ser	Pro	Glu	Asn	His	Lys	Phe
		260					265						270		
Ala	Asn	Glu	Ser	Gly	Met	Ser	Leu	Leu	Asp	Phe	Arg	Phe	Ala	Gln	Lys
		275					280					285			
Val	Arg	Gln	Val	Phe	Arg	Asp	Asn	Thr	Asp	Asn	Met	Tyr	Gly	Leu	Lys
	290					295					300				
Ala	Met	Leu	Glu	Gly	Ser	Ala	Ala	Asp	Tyr	Ala	Gln	Val	Asp	Asp	Gln
305					310					315					320
Val	Thr	Phe	Ile	Asp	Asn	His	Asp	Met	Glu	Arg	Phe	His	Ala	Ser	Asn
				325					330					335	
Ala	Asn	Arg	Arg	Lys	Leu	Glu	Gln	Ala	Leu	Ala	Phe	Thr	Leu	Thr	Ser
				340				345					350		
Arg	Gly	Val	Pro	Ala	Ile	Tyr	Tyr	Gly	Thr	Glu	Gln	Tyr	Met	Ser	Gly
		355					360					365			
Gly	Thr	Asp	Pro	Asp	Asn	Arg	Ala	Arg	Ile	Pro	Ser	Phe	Ser	Thr	Ser
	370					375					380				
Thr	Thr	Ala	Tyr	Gln	Val	Ile	Gln	Lys	Leu	Ala	Pro	Leu	Arg	Lys	Cys
385					390					395					400
Asn	Pro	Ala	Ile	Ala	Tyr	Gly	Ser	Thr	Gln	Glu	Arg	Trp	Ile	Asn	Asn
				405					410					415	
Asp	Val	Leu	Ile	Tyr	Glu	Arg	Lys	Phe	Gly	Ser	Asn	Val	Ala	Val	Val
		420						425					430		
Ala	Val	Asn	Arg	Asn	Leu	Asn	Ala	Pro	Ala	Ser	Ile	Ser	Gly	Leu	Val
		435					440					445			
Thr	Ser	Leu	Pro	Gln	Gly	Ser	Tyr	Asn	Asp	Val	Leu	Gly	Gly	Leu	Leu
	450					455					460				
Asn	Gly	Asn	Thr	Leu	Ser	Val	Gly	Ser	Gly	Gly	Ala	Ala	Ser	Asn	Phe
465					470					475					480
Thr	Leu	Ala	Ala	Gly	Gly	Thr	Ala	Val	Trp	Gln	Tyr	Thr	Ala	Ala	Thr
				485					490					495	
Ala	Thr	Pro	Thr	Ile	Gly	His	Val	Gly	Pro	Met	Met	Ala	Lys	Pro	Gly
		500						505					510		
Val	Thr	Ile	Thr	Ile	Asp	Gly	Arg	Gly	Phe	Gly	Ser	Ser	Lys	Gly	Thr
		515					520					525			
Val	Tyr	Phe	Gly	Thr	Thr	Ala	Val	Ser	Gly	Ala	Asp	Ile	Thr	Ser	Trp
	530					535					540				

Glu Asp Thr Gln Ile Lys Val Lys Ile Pro Ala Val Ala Gly Gly Asn  
 545 550 555 560  
 Tyr Asn Ile Lys Val Ala Asn Ala Ala Gly Thr Ala Ser Asn Val Tyr  
 565 570 575  
 Asp Asn Phe Glu Val Leu Ser Gly Asp Gln Val Ser Val Arg Phe Val  
 580 585 590  
 Val Asn Asn Ala Thr Thr Ala Leu Gly Gln Asn Val Tyr Leu Thr Gly  
 595 600 605  
 Ser Val Ser Glu Leu Gly Asn Trp Asp Pro Ala Lys Ala Ile Gly Pro  
 610 615 620  
 Met Tyr Asn Gln Val Val Tyr Gln Tyr Pro Asn Trp Tyr Tyr Asp Val  
 625 630 635 640  
 Ser Val Pro Ala Gly Lys Thr Ile Glu Phe Lys Phe Leu Lys Lys Gln  
 645 650 655  
 Gly Ser Thr Val Thr Trp Glu Gly Gly Ser Asn His Thr Phe Thr Ala  
 660 665 670  
 Pro Ser Ser Gly Thr Ala Thr Ile Asn Val Asn Trp Gln Pro  
 675 680 685

<210> 45  
 <211> 404  
 <212> PRT  
 <213> Amycolatopsis orientalis

<400> 45  
 Met Arg Val Leu Ile Thr Gly Cys Gly Ser Arg Gly Asp Thr Glu Pro  
 1 5 10 15  
 Leu Val Ala Leu Ala Ala Arg Leu Arg Glu Leu Gly Ala Asp Ala Arg  
 20 25 30  
 Met Cys Leu Pro Pro Asp Tyr Val Glu Arg Cys Ala Glu Val Gly Val  
 35 40 45  
 Pro Met Val Pro Val Gly Arg Ala Val Arg Ala Gly Ala Arg Glu Pro  
 50 55 60  
 Gly Glu Leu Pro Pro Gly Ala Ala Glu Val Val Thr Glu Val Val Ala  
 65 70 75 80  
 Glu Trp Phe Asp Lys Val Pro Ala Ala Ile Glu Gly Cys Asp Ala Val  
 85 90 95  
 Val Thr Thr Gly Leu Leu Pro Ala Ala Val Ala Val Arg Ser Met Ala  
 100 105 110  
 Glu Lys Leu Gly Ile Pro Tyr Arg Tyr Thr Val Leu Ser Pro Asp His  
 115 120 125  
 Leu Pro Ser Glu Gln Ser Gln Ala Glu Arg Asp Met Tyr Asn Gln Gly  
 130 135 140  
 Ala Asp Arg Leu Phe Gly Asp Ala Val Asn Ser His Arg Ala Ser Ile  
 145 150 155 160  
 Gly Leu Pro Pro Val Glu His Leu Tyr Asp Tyr Gly Tyr Thr Asp Gln  
 165 170 175  
 Pro Trp Leu Ala Ala Asp Pro Val Leu Ser Pro Leu Arg Pro Thr Asp  
 180 185 190  
 Leu Gly Thr Val Gln Thr Gly Ala Trp Ile Leu Pro Asp Glu Arg Pro  
 195 200 205  
 Leu Ser Ala Glu Leu Glu Ala Phe Leu Ala Ala Gly Ser Thr Pro Val  
 210 215 220  
 Tyr Val Gly Phe Gly Ser Ser Ser Arg Pro Ala Thr Ala Asp Ala Ala  
 225 230 235 240  
 Lys Met Ala Ile Lys Ala Val Arg Ala Ser Gly Arg Arg Ile Val Leu  
 245 250 255  
 Ser Arg Gly Trp Ala Asp Leu Val Leu Pro Asp Asp Gly Ala Asp Cys  
 260 265 270  
 Phe Val Val Gly Glu Val Asn Leu Gln Glu Leu Phe Gly Arg Val Ala  
 275 280 285  
 Ala Ala Ile His His Asp Ser Ala Gly Thr Thr Leu Leu Ala Met Arg

```

      290                      295                      300
Ala Gly Ile Pro Gln Ile Val Val Arg Arg Val Val Asp Asn Val Val
305                      310                      315                      320
Glu Gln Ala Tyr His Ala Asp Arg Val Ala Glu Leu Gly Val Gly Val
      325                      330                      335
Ala Val Asp Gly Pro Val Pro Thr Ile Asp Ser Leu Ser Ala Ala Leu
      340                      345                      350
Asp Thr Ala Leu Ala Pro Glu Ile Arg Ala Arg Ala Thr Thr Val Ala
      355                      360                      365
Asp Thr Ile Arg Ala Asp Gly Thr Thr Val Ala Ala Gln Leu Leu Phe
      370                      375                      380
Asp Ala Val Ser Leu Glu Lys Pro Thr Val Pro Ala Leu Glu His His
385                      390                      395                      400
His His His His

```

```

<210> 46
<211> 292
<212> PRT
<213> Pseudomonas sp.

```

```

<400> 46
Ser Ile Glu Arg Leu Gly Tyr Leu Gly Phe Ala Val Lys Asp Val Pro
1                      5                      10                      15
Ala Trp Asp His Phe Leu Thr Lys Ser Val Gly Leu Met Ala Ala Gly
      20                      25                      30
Ser Ala Gly Asp Ala Ala Leu Tyr Arg Ala Asp Gln Arg Ala Trp Arg
      35                      40                      45
Ile Ala Val Gln Pro Gly Glu Leu Asp Asp Leu Ala Tyr Ala Gly Leu
      50                      55                      60
Glu Val Asp Asp Ala Ala Ala Leu Glu Arg Met Ala Asp Lys Leu Arg
65                      70                      75                      80
Gln Ala Gly Val Ala Phe Thr Arg Gly Asp Glu Ala Leu Met Gln Gln
      85                      90                      95
Arg Lys Val Met Gly Leu Leu Cys Leu Gln Asp Pro Phe Gly Leu Pro
      100                     105                     110
Leu Glu Ile Tyr Tyr Gly Pro Ala Glu Ile Phe His Glu Pro Phe Leu
      115                     120                     125
Pro Ser Ala Pro Val Ser Gly Phe Val Thr Gly Asp Gln Gly Ile Gly
      130                     135                     140
His Phe Val Arg Cys Val Pro Asp Thr Ala Lys Ala Met Ala Phe Tyr
145                     150                     155                     160
Thr Glu Val Leu Gly Phe Val Leu Ser Asp Ile Ile Asp Ile Gln Met
      165                     170                     175
Gly Pro Glu Thr Ser Val Pro Ala His Phe Leu His Cys Asn Gly Arg
      180                     185                     190
His His Thr Ile Ala Leu Ala Ala Phe Pro Ile Pro Lys Arg Ile His
      195                     200                     205
His Phe Met Leu Gln Ala Asn Thr Ile Asp Asp Val Gly Tyr Ala Phe
      210                     215                     220
Asp Arg Leu Asp Ala Ala Gly Arg Ile Thr Ser Leu Leu Gly Arg His
225                     230                     235                     240
Thr Asn Asp Gln Thr Leu Ser Phe Tyr Ala Asp Thr Pro Ser Pro Met
      245                     250                     255
Ile Glu Val Glu Phe Gly Trp Gly Pro Arg Thr Val Asp Ser Ser Trp
      260                     265                     270
Thr Val Ala Arg His Ser Arg Thr Ala Met Trp Gly His Lys Ser Val
      275                     280                     285
Arg Gly Gln Arg
290

```

<210> 47  
 <211> 311  
 <212> PRT  
 <213> Acitenobacter sp.

<400> 47  
 Met Glu Val Lys Ile Phe Asn Thr Gln Asp Val Gln Asp Phe Leu Arg  
 1 5 10 15  
 Val Ala Ser Gly Leu Glu Gln Glu Gly Gly Asn Pro Arg Val Lys Gln  
 20 25 30  
 Ile Ile His Arg Val Leu Ser Asp Leu Tyr Lys Ala Ile Glu Asp Leu  
 35 40 45  
 Asn Ile Thr Ser Asp Glu Tyr Trp Ala Gly Val Ala Tyr Leu Asn Gln  
 50 55 60  
 Leu Gly Ala Asn Gln Glu Ala Gly Leu Leu Ser Pro Gly Leu Gly Phe  
 65 70 75 80  
 Asp His Tyr Leu Asp Met Arg Met Asp Ala Glu Asp Ala Ala Leu Gly  
 85 90 95  
 Ile Glu Asn Ala Thr Pro Arg Thr Ile Glu Gly Pro Leu Tyr Val Ala  
 100 105 110  
 Gly Ala Pro Glu Ser Val Gly Tyr Ala Arg Met Asp Asp Gly Ser Asp  
 115 120 125  
 Pro Asn Gly His Thr Leu Ile Leu His Gly Thr Ile Phe Asp Ala Asp  
 130 135 140  
 Gly Lys Pro Leu Pro Asn Ala Lys Val Glu Ile Trp His Ala Asn Thr  
 145 150 155 160  
 Lys Gly Phe Tyr Ser His Phe Asp Pro Thr Gly Glu Gln Gln Ala Phe  
 165 170 175  
 Asn Met Arg Arg Ser Ile Ile Thr Asp Glu Asn Gly Gln Tyr Arg Val  
 180 185 190  
 Arg Thr Ile Leu Pro Ala Gly Tyr Gly Cys Pro Pro Glu Gly Pro Thr  
 195 200 205  
 Gln Gln Leu Leu Asn Gln Leu Gly Arg His Gly Asn Arg Pro Ala His  
 210 215 220  
 Ile His Tyr Phe Val Ser Ala Asp Gly His Arg Lys Leu Thr Thr Gln  
 225 230 235 240  
 Ile Asn Val Ala Gly Asp Pro Tyr Thr Tyr Asp Asp Phe Ala Tyr Ala  
 245 250 255  
 Thr Arg Glu Gly Leu Val Val Asp Ala Val Glu His Thr Asp Pro Glu  
 260 265 270  
 Ala Ile Lys Ala Asn Asp Val Glu Gly Pro Phe Ala Glu Met Val Phe  
 275 280 285  
 Asp Leu Lys Leu Thr Arg Leu Val Asp Gly Val Asp Asn Gln Val Val  
 290 295 300  
 Asp Arg Pro Arg Leu Ala Val  
 305 310

<210> 48  
 <211> 414  
 <212> PRT  
 <213> Pseudomonas putida

<400> 48  
 Thr Thr Glu Thr Ile Gln Ser Asn Ala Asn Leu Ala Pro Leu Pro Pro  
 1 5 10 15  
 His Val Pro Glu His Leu Val Phe Asp Phe Asp Met Tyr Asn Pro Ser  
 20 25 30  
 Asn Leu Ser Ala Gly Val Gln Glu Ala Trp Ala Val Leu Gln Glu Ser  
 35 40 45  
 Asn Val Pro Asp Leu Val Trp Thr Arg Cys Asn Gly Gly His Trp Ile  
 50 55 60  
 Ala Thr Arg Gly Gln Leu Ile Arg Glu Ala Tyr Glu Asp Tyr Arg His

65					70					75					80
Phe	Ser	Ser	Glu	Cys	Pro	Phe	Ile	Pro	Arg	Glu	Ala	Gly	Glu	Ala	Tyr
				85					90					95	
Asp	Phe	Ile	Pro	Thr	Ser	Met	Asp	Pro	Pro	Glu	Gln	Arg	Gln	Phe	Arg
			100					105					110		
Ala	Leu	Ala	Asn	Gln	Val	Val	Gly	Met	Pro	Val	Val	Asp	Lys	Leu	Glu
		115					120					125			
Asn	Arg	Ile	Gln	Glu	Leu	Ala	Cys	Ser	Leu	Ile	Glu	Ser	Leu	Arg	Pro
	130					135					140				
Gln	Gly	Gln	Cys	Asn	Phe	Thr	Glu	Asp	Tyr	Ala	Glu	Pro	Phe	Pro	Ile
145					150					155					160
Arg	Ile	Phe	Met	Leu	Leu	Ala	Gly	Leu	Pro	Glu	Glu	Asp	Ile	Pro	His
			165						170					175	
Leu	Lys	Tyr	Leu	Thr	Asp	Gln	Met	Thr	Arg	Pro	Asp	Gly	Ser	Met	Thr
			180					185					190		
Phe	Ala	Glu	Ala	Lys	Glu	Ala	Leu	Tyr	Asp	Tyr	Leu	Ile	Pro	Ile	Ile
		195					200					205			
Glu	Gln	Arg	Arg	Gln	Lys	Pro	Gly	Thr	Asp	Ala	Ile	Ser	Ile	Val	Ala
	210					215					220				
Asn	Gly	Gln	Val	Asn	Gly	Arg	Pro	Ile	Thr	Ser	Asp	Glu	Ala	Lys	Arg
225					230					235					240
Met	Cys	Gly	Leu	Leu	Leu	Val	Gly	Gly	Leu	Asp	Thr	Val	Val	Asn	Phe
			245						250					255	
Leu	Ser	Phe	Ser	Met	Glu	Phe	Leu	Ala	Lys	Ser	Pro	Glu	His	Arg	Gln
		260					265						270		
Glu	Leu	Ile	Gln	Arg	Pro	Glu	Arg	Ile	Pro	Ala	Ala	Cys	Glu	Glu	Leu
		275					280					285			
Leu	Arg	Arg	Phe	Ser	Leu	Val	Ala	Asp	Gly	Arg	Ile	Leu	Thr	Ser	Asp
	290					295					300				
Tyr	Glu	Phe	His	Gly	Val	Gln	Leu	Lys	Lys	Gly	Asp	Gln	Ile	Leu	Leu
305					310					315					320
Pro	Gln	Met	Leu	Ser	Gly	Leu	Asp	Glu	Arg	Glu	Asn	Ala	Cys	Pro	Met
			325						330					335	
His	Val	Asp	Phe	Ser	Arg	Gln	Lys	Val	Ser	His	Thr	Thr	Phe	Gly	His
		340						345					350		
Gly	Ser	His	Leu	Cys	Leu	Gly	Gln	His	Leu	Ala	Arg	Arg	Glu	Ile	Ile
		355					360					365			
Val	Thr	Leu	Lys	Glu	Trp	Leu	Thr	Arg	Ile	Pro	Asp	Phe	Ser	Ile	Ala
	370					375					380				
Pro	Gly	Ala	Gln	Ile	Gln	His	Lys	Ser	Gly	Ile	Val	Ser	Gly	Val	Gln
385					390					395					400
Ala	Leu	Pro	Leu	Val	Trp	Asp	Pro	Ala	Thr	Thr	Lys	Ala	Val		
			405						410						

<210> 49  
 <211> 374  
 <212> PRT  
 <213> Equus caballus

<400> 49

Ser	Thr	Ala	Gly	Lys	Val	Ile	Lys	Cys	Lys	Ala	Ala	Val	Leu	Trp	Glu
1				5					10					15	
Glu	Lys	Lys	Pro	Phe	Ser	Ile	Glu	Glu	Val	Glu	Val	Ala	Pro	Pro	Lys
			20					25				30			
Ala	His	Glu	Val	Arg	Ile	Lys	Met	Val	Ala	Thr	Gly	Ile	Cys	Arg	Ser
		35					40					45			
Asp	Asp	His	Val	Val	Ser	Gly	Thr	Leu	Val	Thr	Pro	Leu	Pro	Val	Ile
		50				55					60				
Ala	Gly	His	Glu	Ala	Ala	Gly	Ile	Val	Glu	Ser	Ile	Gly	Glu	Gly	Val
65					70					75					80
Thr	Thr	Val	Arg	Pro	Gly	Asp	Lys	Val	Ile	Pro	Leu	Phe	Thr	Pro	Gln
				85					90					95	

Cys Gly Lys Cys Arg Val Cys Lys His Pro Glu Gly Asn Phe Cys Leu  
 100 105 110  
 Lys Asn Asp Leu Ser Met Pro Arg Gly Thr Met Gln Asp Gly Thr Ser  
 115 120 125  
 Arg Phe Thr Cys Arg Gly Lys Pro Ile His His Phe Leu Gly Thr Ser  
 130 135 140  
 Thr Phe Ser Gln Tyr Thr Val Val Asp Glu Ile Ser Val Ala Lys Ile  
 145 150 155 160  
 Asp Ala Ala Ser Pro Leu Glu Lys Val Cys Leu Ile Gly Cys Gly Phe  
 165 170 175  
 Ser Thr Gly Tyr Gly Ser Ala Val Lys Val Ala Lys Val Thr Gln Gly  
 180 185 190  
 Ser Thr Cys Ala Val Phe Gly Leu Gly Gly Val Gly Leu Ser Val Ile  
 195 200 205  
 Met Gly Cys Lys Ala Ala Gly Ala Ala Arg Ile Ile Gly Val Asp Ile  
 210 215 220  
 Asn Lys Asp Lys Phe Ala Lys Ala Lys Glu Val Gly Ala Thr Glu Cys  
 225 230 235 240  
 Val Asn Pro Gln Asp Tyr Lys Lys Pro Ile Gln Glu Val Leu Thr Glu  
 245 250 255  
 Met Ser Asn Gly Gly Val Asp Phe Ser Phe Glu Val Ile Gly Arg Leu  
 260 265 270  
 Asp Thr Met Val Thr Ala Leu Ser Cys Cys Gln Glu Ala Tyr Gly Val  
 275 280 285  
 Ser Val Ile Val Gly Val Pro Asp Ser Gln Asn Leu Ser Met Asn  
 290 295 300  
 Pro Met Leu Leu Leu Ser Gly Arg Thr Trp Lys Gly Ala Ile Phe Gly  
 305 310 315 320  
 Gly Phe Lys Ser Lys Asp Ser Val Pro Lys Leu Val Ala Asp Phe Met  
 325 330 335  
 Ala Lys Lys Phe Ala Leu Asp Pro Leu Ile Thr His Val Leu Pro Phe  
 340 345 350  
 Glu Lys Ile Asn Glu Gly Phe Asp Leu Leu Arg Ser Gly Glu Ser Ile  
 355 360 365  
 Arg Thr Ile Leu Thr Phe  
 370

<210> 50  
 <211> 297  
 <212> PRT  
 <213> Escherichia coli

<400> 50  
 Met Ala Thr Asn Leu Arg Gly Val Met Ala Ala Leu Leu Thr Pro Phe  
 1 5 10 15  
 Asp Gln Gln Gln Ala Leu Asp Lys Ala Ser Leu Arg Arg Leu Val Gln  
 20 25 30  
 Phe Asn Ile Gln Gln Gly Ile Asp Gly Leu Tyr Val Gly Gly Ser Thr  
 35 40 45  
 Gly Glu Ala Phe Val Gln Ser Leu Ser Glu Arg Glu Gln Val Leu Glu  
 50 55 60  
 Ile Val Ala Glu Glu Gly Lys Gly Lys Ile Lys Leu Ile Ala His Val  
 65 70 75 80  
 Gly Cys Val Thr Thr Ala Glu Ser Gln Gln Leu Ala Ala Ser Ala Lys  
 85 90 95  
 Arg Tyr Gly Phe Asp Ala Val Ser Ala Val Thr Pro Phe Tyr Tyr Pro  
 100 105 110  
 Phe Ser Phe Glu Glu His Cys Asp His Tyr Arg Ala Ile Ile Asp Ser  
 115 120 125  
 Ala Asp Gly Leu Pro Met Val Val Tyr Asn Ile Pro Ala Leu Ser Gly  
 130 135 140  
 Val Lys Leu Thr Leu Asp Gln Ile Asn Thr Leu Val Thr Leu Pro Gly



```

145          150          155          160
Val Gly Ala Leu Lys Gln Thr Ser Gly Asp Leu Tyr Gln Met Glu Gln
          165          170          175
Ile Arg Arg Glu His Pro Asp Leu Val Leu Tyr Asn Gly Tyr Asp Glu
          180          185          190
Ile Phe Ala Ser Gly Leu Leu Ala Gly Ala Asp Gly Gly Ile Gly Ser
          195          200          205
Thr Tyr Asn Ile Met Gly Trp Arg Tyr Gln Gly Ile Val Lys Ala Leu
          210          215          220
Lys Glu Gly Asp Ile Gln Thr Ala Gln Lys Leu Gln Thr Glu Cys Asn
225          230          235          240
Lys Val Ile Asp Leu Leu Ile Lys Thr Gly Val Phe Arg Gly Leu Lys
          245          250          255
Thr Val Leu His Tyr Met Asp Val Val Ser Val Pro Leu Cys Arg Lys
          260          265          270
Pro Phe Gly Pro Val Asp Glu Lys Tyr Leu Pro Glu Leu Lys Ala Leu
          275          280          285
Ala Gln Gln Leu Met Gln Glu Arg Gly
          290          295

```

```

<210> 51
<211> 268
<212> PRT
<213> Salmonella typhimurium

```

```

<400> 51
Met Glu Arg Tyr Glu Asn Leu Phe Ala Gln Leu Asn Asp Arg Arg Glu
1          5          10          15
Gly Ala Phe Val Pro Phe Val Thr Leu Gly Asp Pro Gly Ile Glu Gln
          20          25          30
Ser Leu Lys Ile Ile Asp Thr Leu Ile Asp Ala Gly Ala Asp Ala Leu
          35          40          45
Glu Leu Gly Val Pro Phe Ser Asp Pro Leu Ala Asp Gly Pro Thr Ile
50          55          60
Gln Asn Ala Asn Leu Arg Ala Phe Ala Ala Gly Val Thr Pro Ala Gln
65          70          75          80
Cys Phe Glu Met Leu Ala Leu Ile Arg Glu Lys His Pro Thr Ile Pro
          85          90          95
Ile Gly Leu Leu Met Tyr Ala Asn Leu Val Phe Asn Asn Gly Ile Asp
          100          105          110
Ala Phe Tyr Ala Arg Cys Glu Gln Val Gly Val Asp Ser Val Leu Val
          115          120          125
Ala Asp Val Pro Val Glu Glu Ser Ala Pro Phe Arg Gln Ala Ala Leu
          130          135          140
Arg His Asn Ile Ala Pro Ile Phe Ile Cys Pro Pro Asn Ala Asp Asp
145          150          155          160
Asp Leu Leu Arg Gln Val Ala Ser Tyr Gly Arg Gly Tyr Thr Tyr Leu
          165          170          175
Leu Ser Arg Ser Gly Val Thr Gly Ala Glu Asn Arg Gly Ala Leu Pro
          180          185          190
Leu His His Leu Ile Glu Lys Leu Lys Glu Tyr His Ala Ala Pro Ala
          195          200          205
Leu Gln Gly Phe Gly Ile Ser Ser Pro Glu Gln Val Ser Ala Ala Val
          210          215          220
Arg Ala Gly Ala Ala Gly Ala Ile Ser Gly Ser Ala Ile Val Lys Ile
225          230          235          240
Ile Glu Lys Asn Leu Ala Ser Pro Lys Gln Met Leu Ala Glu Leu Arg
          245          250          255
Ser Phe Val Ser Ala Met Lys Ala Ala Ser Arg Ala
          260          265

```

<210> 52  
 <211> 393  
 <212> PRT  
 <213> Actinoplanes missouriensis

<400> 52  
 Ser Val Gln Ala Thr Arg Glu Asp Lys Phe Ser Phe Gly Leu Trp Thr  
 1 5 10 15  
 Val Gly Trp Gln Ala Arg Asp Ala Phe Gly Asp Ala Thr Arg Thr Ala  
 20 25 30  
 Leu Asp Pro Val Glu Ala Val His Lys Leu Ala Glu Ile Gly Ala Tyr  
 35 40 45  
 Gly Ile Thr Phe His Asp Asp Leu Val Pro Phe Gly Ser Asp Ala  
 50 55 60  
 Gln Thr Arg Asp Gly Ile Ile Ala Gly Phe Lys Lys Ala Leu Asp Glu  
 65 70 75 80  
 Thr Gly Leu Ile Val Pro Met Val Thr Thr Asn Leu Phe Thr His Pro  
 85 90 95  
 Val Phe Lys Asp Gly Gly Phe Thr Ser Asn Asp Arg Ser Val Arg Arg  
 100 105 110  
 Tyr Ala Ile Arg Lys Val Leu Arg Gln Met Asp Leu Gly Ala Glu Leu  
 115 120 125  
 Gly Ala Lys Thr Leu Val Leu Trp Gly Gly Arg Glu Gly Ala Glu Tyr  
 130 135 140  
 Asp Ser Ala Lys Asp Val Ser Ala Ala Leu Asp Arg Tyr Arg Glu Ala  
 145 150 155 160  
 Leu Asn Leu Leu Ala Gln Tyr Ser Glu Asp Arg Gly Tyr Gly Leu Arg  
 165 170 175  
 Phe Ala Ile Glu Pro Lys Pro Asn Glu Pro Arg Gly Asp Ile Leu Leu  
 180 185 190  
 Pro Thr Ala Gly His Ala Ile Ala Phe Val Gln Glu Leu Glu Arg Pro  
 195 200 205  
 Glu Leu Phe Gly Ile Asn Pro Glu Thr Gly Asn Glu Gln Met Ser Asn  
 210 215 220  
 Leu Asn Phe Thr Gln Gly Ile Ala Gln Ala Leu Trp His Lys Lys Leu  
 225 230 235 240  
 Phe His Ile Asp Leu Asn Gly Gln His Gly Pro Lys Phe Asp Gln Asp  
 245 250 255  
 Leu Val Phe Gly His Gly Asp Leu Leu Asn Ala Phe Ser Leu Val Asp  
 260 265 270  
 Leu Leu Glu Asn Gly Pro Asp Gly Ala Pro Ala Tyr Asp Gly Pro Arg  
 275 280 285  
 His Phe Asp Tyr Lys Pro Ser Arg Thr Glu Asp Tyr Asp Gly Val Trp  
 290 295 300  
 Glu Ser Ala Lys Ala Asn Ile Arg Met Tyr Leu Leu Leu Lys Glu Arg  
 305 310 315 320  
 Ala Lys Ala Phe Arg Ala Asp Pro Glu Val Gln Glu Ala Leu Ala Ala  
 325 330 335  
 Ser Lys Val Ala Glu Leu Lys Thr Pro Thr Leu Asn Pro Gly Glu Gly  
 340 345 350  
 Tyr Ala Glu Leu Leu Ala Asp Arg Ser Ala Phe Glu Asp Tyr Asp Ala  
 355 360 365  
 Asp Ala Val Gly Ala Lys Gly Phe Gly Phe Val Lys Leu Asn Gln Leu  
 370 375 380  
 Ala Ile Glu His Leu Leu Gly Ala Arg  
 385 390

<210> 53  
 <211> 348  
 <212> PRT  
 <213> Bacteriophage T7

&lt;400&gt; 53

```

Val Asn Ile Lys Thr Asn Pro Phe Lys Ala Val Ser Phe Val Glu Ser
1      5      10      15
Ala Ile Lys Lys Ala Leu Asp Asn Ala Gly Tyr Leu Ile Ala Glu Ile
20      25      30
Lys Tyr Asp Gly Val Arg Gly Asn Ile Cys Val Asp Asn Thr Ala Asn
35      40      45
Ser Tyr Trp Leu Ser Arg Val Ser Lys Thr Ile Pro Ala Leu Glu His
50      55      60
Leu Asn Gly Phe Asp Val Arg Trp Lys Arg Leu Leu Asn Asp Asp Arg
65      70      75      80
Cys Phe Tyr Lys Asp Gly Phe Met Leu Asp Gly Glu Leu Met Val Lys
85      90      95
Gly Val Asp Phe Asn Thr Gly Ser Gly Leu Leu Arg Thr Lys Trp Thr
100     105     110
Asp Thr Lys Asn Gln Glu Phe His Glu Glu Leu Phe Val Glu Pro Ile
115     120     125
Arg Lys Lys Asp Lys Val Pro Phe Lys Leu His Thr Gly His Leu His
130     135     140
Ile Lys Leu Tyr Ala Ile Leu Pro Leu His Ile Val Glu Ser Gly Glu
145     150     155     160
Asp Cys Asp Val Met Thr Leu Leu Met Gln Glu His Val Lys Asn Met
165     170     175
Leu Pro Leu Leu Gln Glu Tyr Phe Pro Glu Ile Glu Trp Gln Ala Ala
180     185     190
Glu Ser Tyr Glu Val Tyr Asp Met Val Glu Leu Gln Gln Leu Tyr Glu
195     200     205
Gln Lys Arg Ala Glu Gly His Glu Gly Leu Ile Val Lys Asp Pro Met
210     215     220
Cys Ile Tyr Lys Arg Gly Lys Lys Ser Gly Trp Trp Lys Met Lys Pro
225     230     235     240
Glu Asn Glu Ala Asp Gly Ile Ile Gln Gly Leu Val Trp Gly Thr Lys
245     250     255
Gly Leu Ala Asn Glu Gly Lys Val Ile Gly Phe Glu Val Leu Leu Glu
260     265     270
Ser Gly Arg Leu Val Asn Ala Thr Asn Ile Ser Arg Ala Leu Met Asp
275     280     285
Glu Phe Thr Glu Thr Val Lys Glu Ala Thr Leu Ser Gln Trp Gly Phe
290     295     300
Phe Ser Pro Tyr Gly Ile Gly Asp Asn Asp Ala Cys Thr Ile Asn Pro
305     310     315     320
Tyr Asp Gly Trp Ala Cys Gln Ile Ser Tyr Met Glu Glu Thr Pro Asp
325     330     335
Gly Ser Leu Arg His Pro Ser Phe Val Met Phe Arg
340     345

```

&lt;210&gt; 54

&lt;211&gt; 42

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; binding site for restr1 and restr2

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (2)..(40)

&lt;223&gt;

&lt;400&gt; 54

```

g gtg gta tca gca ggc cac tgc tac aag tcc cgc atc cag gt
Val Val Ser Ala Gly His Cys Tyr Lys Ser Arg Ile Gln
1      5      10

```

42

<210> 55  
 <211> 13  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> binding site for restr1 and restr2  
  
 <400> 55  
 Val Val Ser Ala Gly His Cys Tyr Lys Ser Arg Ile Gln  
 1 5 10

<210> 56  
 <211> 42  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> forward primer restr1

<400> 56  
 ggtggatatcc gcggggccact gctacaagtc ccggatccag gt

42

<210> 57  
 <211> 42  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> reverse primer restr2

<400> 57  
 acctggatcc gggacttgta gcagtggccc gcggatacca cc

42

<210> 58  
 <211> 50  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> binding site for restr3 and restr4  
 <220>  
 <221> CDS  
 <222> (3)..(50)  
 <223>

<400> 58  
 cc act ggc acg aag tgc ctc atc tct ggc tgg ggc aac act gcg agc  
 Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly Asn Thr Ala Ser  
 1 5 10 15  
 tct  
 Ser

47

50

<210> 59  
 <211> 16  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> binding site for restr3 and restr4

<400> 59

Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly Asn Thr Ala Ser Ser  
 1 5 10 15

<210> 60  
 <211> 50  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> forward primer restr3

<400> 60  
 ccactggcac gaagtgcctc atctctggct ggggcaacac tgcgagctct 50

<210> 61  
 <211> 50  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> reverse primer restr4

<400> 61  
 agagctagca gtgttgcccc agccagagat gaggcacttg gtaccagtgg 50

<210> 62  
 <211> 30  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> primer puc-forward

<400> 62  
 ggggtacccc accaccatga atccactcct 30

<210> 63  
 <211> 30  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> primer puc-reverse

<400> 63  
 cgggatccgg tatagagact gaagagatac 30

<210> 64  
 <211> 39  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> oligox-SDR1f  
 <220>  
 <221> misc\_feature  
 <222> (14)..(31)  
 <223> any nucleotide  
 <220>  
 <221> misc\_feature  
 <222> (14)..(31)  
 <223> any nucleotide or amino acid residue  
 <220>

```

<221> CDS
<222> (2)..(37)
<223>

<400> 64
g ggc cac tgc tac nnn nnn nnn nnn nnn nnn aag tcc cg          39
  Gly His Cys Tyr Xaa Xaa Xaa Xaa Xaa Xaa Lys Ser
    1                      5                      10

<210> 65
<211> 12
<212> PRT
<213> artificial sequence
<220>
<221> misc_feature
<222> (5)..(5)
<223> The 'Xaa' at location 5 stands for Lys, Asn, Arg, Ser, Thr, Ile,
      Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, T
      yr, Trp, Cys, or Phe.

<220>
<221> misc_feature
<222> (6)..(6)
<223> The 'Xaa' at location 6 stands for Lys, Asn, Arg, Ser, Thr, Ile,
      Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, T
      yr, Trp, Cys, or Phe.

<220>
<221> misc_feature
<222> (7)..(7)
<223> The 'Xaa' at location 7 stands for Lys, Asn, Arg, Ser, Thr, Ile,
      Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, T
      yr, Trp, Cys, or Phe.

<220>
<221> misc_feature
<222> (8)..(8)
<223> The 'Xaa' at location 8 stands for Lys, Asn, Arg, Ser, Thr, Ile,
      Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, T
      yr, Trp, Cys, or Phe.

<220>
<221> misc_feature
<222> (9)..(9)
<223> The 'Xaa' at location 9 stands for Lys, Asn, Arg, Ser, Thr, Ile,
      Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, T
      yr, Trp, Cys, or Phe.

<220>
<221> misc_feature
<222> (10)..(10)
<223> The 'Xaa' at location 10 stands for Lys, Asn, Arg, Ser, Thr, Ile,
      Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon,
      Tyr, Trp, Cys, or Phe.

<220>
<223> oligox-SDR1f
<220>
<221> misc_feature
<222> (14)..(31)
<223> any nucleotide

```

<220>  
 <221> misc\_feature  
 <222> (14)..(31)  
 <223> any nucleotide or amino acid residue

<400> 65  
 Gly His Cys Tyr Xaa Xaa Xaa Xaa Xaa Xaa Lys Ser  
 1 5 10

<210> 66  
 <211> 45  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> oligox-SDR1r  
 <220>  
 <221> misc\_feature  
 <222> (16)..(33)  
 <223> any nucleotide

<400> 66  
 cgcccggtga cgatgnnnnn nnnnnnnnnn nnnttcaggg cctag

45

<210> 67  
 <211> 47  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> oligox-SDR2f  
 <220>  
 <221> CDS  
 <222> (2)..(46)  
 <223>  
 <220>  
 <221> misc\_feature  
 <222> (29)..(43)  
 <223> any nucleotide or amino acid residue

<400> 67  
 c aag tgc ctc atc tct ggc tgg ggc aac nnn nnn nnn nnn nnn act g  
 Lys Cys Leu Ile Ser Gly Trp Gly Asn Xaa Xaa Xaa Xaa Xaa Thr  
 1 5 10 15

47

<210> 68  
 <211> 15  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <221> misc\_feature  
 <222> (10)..(10)  
 <223> The 'Xaa' at location 10 stands for Lys, Asn, Arg, Ser, Thr, Ile,  
 Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon,  
 Tyr, Trp, Cys, or Phe.

<220>  
 <221> misc\_feature  
 <222> (11)..(11)  
 <223> The 'Xaa' at location 11 stands for Lys, Asn, Arg, Ser, Thr, Ile,  
 Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon,  
 Tyr, Trp, Cys, or Phe.

<220>  
 <221> misc\_feature  
 <222> (12)..(12)  
 <223> The 'Xaa' at location 12 stands for Lys, Asn, Arg, Ser, Thr, Ile, Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, Tyr, Trp, Cys, or Phe.

<220>  
 <221> misc\_feature  
 <222> (13)..(13)  
 <223> The 'Xaa' at location 13 stands for Lys, Asn, Arg, Ser, Thr, Ile, Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, Tyr, Trp, Cys, or Phe.

<220>  
 <221> misc\_feature  
 <222> (14)..(14)  
 <223> The 'Xaa' at location 14 stands for Lys, Asn, Arg, Ser, Thr, Ile, Met, Glu, Asp, Gly, Ala, Val, Gln, His, Pro, Leu, a stop codon, Tyr, Trp, Cys, or Phe.

<220>  
 <223> oligox-SDR2f  
 <220>  
 <221> misc\_feature  
 <222> (29)..(43)  
 <223> any nucleotide or amino acid residue

<400> 68  
 Lys Cys Leu Ile Ser Gly Trp Gly Asn Xaa Xaa Xaa Xaa Xaa Thr  
 1 5 10 15

<210> 69  
 <211> 55  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> oligox-SDR2r  
 <220>  
 <221> misc\_feature  
 <222> (33)..(47)  
 <223> any base  
 <220>  
 <221> misc\_feature  
 <222> (33)..(47)  
 <223> any nucleotide

<400> 69  
 catggttcac ggagtagaga ccgaccccggt tgnnnnnnnnn nnnnnnnntga cgatc 55

<210> 70  
 <211> 59  
 <212> DNA  
 <213> artificial sequence  
 <220>  
 <223> primer SDR1-mutnnb-forward  
 <220>  
 <221> misc\_feature  
 <222> (24)..(40)  
 <223> N=A, C, G, T; B=C, G, T; V=A, C, G



<400> 70  
tggtatccgc gggccactgc tacnnbnnbn nbnnbnnbnn baagtcccg atccaggtg 59

<210> 71  
<211> 52  
<212> DNA  
<213> artificial sequence  
<220>  
<223> primer SDR2-mutnnb-reverse  
<220>  
<221> misc\_feature  
<222> (20)..(33)  
<223> N=A, C, G, T; B=C, G, T; V=A, C, G

<400> 71  
ggcgccagag ctagcagtvn nvnnvnnvnn vnngttgccc cagccagaga tg 52

<210> 72  
<211> 6  
<212> PRT  
<213> artificial sequence

<220>  
<223> variant g SDR1  
<400> 72  
Ala Phe Phe Asn Gly Asp  
1 5

<210> 73  
<211> 5  
<212> PRT  
<213> artificial sequence

<220>  
<223> variant g SDR2  
<400> 73  
Arg Lys Asp Pro Trp  
1 5

<210> 74  
<211> 234  
<212> PRT  
<213> artificial sequence

<220>  
<223> artificial sequence

&lt;400&gt; 74

```

Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr Gln Val
1           5           10           15
Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu Ile Asn Glu
          20           25           30
Gln Trp Val Val Ser Ala Gly His Cys Tyr Ala Ala Phe Asn Gly Lys
          35           40           45
Ser Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Glu Val Leu Glu
          50           55           60
Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro Gln
65           70           75           80
Tyr Asp Arg Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys Leu Ser
          85           90           95
Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser Leu Pro Thr
          100          105          110
Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly Asn
          115          120          125
Arg Lys Asp Phe Trp Thr Ala Ser Ser Gly Ala Asp Tyr Pro Asp Glu
130          135          140
Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Ala Lys Cys Glu Ala
145          150          155          160
Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly Phe Leu
          165          170          175
Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Val Val
          180          185          190
Cys Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly Asp Gly Cys Ala
          195          200          205
Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val Tyr Asn Tyr Val Lys
          210          215          220
Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser
225          230

```

&lt;210&gt; 75

&lt;211&gt; 234

&lt;212&gt; PRT

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; artificial sequence

&lt;400&gt; 75

```

Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr Gln Val

```

```

1             5             10             15
Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu Ile Asn Glu
             20             25             30
Gln Trp Val Val Ser Ala Gly His Cys Tyr Ala Ala Phe Asn Gly Lys
             35             40             45
Ser Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Gly Val Leu Glu
             50             55             60
Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro Gln
65             70             75             80
Tyr Asp Trp Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys Leu Ser
             85             90             95
Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser Leu Pro Thr
             100             105             110
Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser Gly Trp Gly Asn
             115             120             125
Arg Lys Asp Phe Trp Thr Ala Ser Ser Gly Ala Asp Phe Pro Asp Glu
             130             135             140
Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Thr Lys Cys Glu Ala
145             150             155             160
Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly Phe Leu
             165             170             175
Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Val Val
             180             185             190
Arg Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly Asp Gly Cys Ala
             195             200             205
Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val Tyr Asn Tyr Val Lys
             210             215             220
Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser
225             230

```

```

<210> 76
<211> 12
<212> PRT
<213> artificial sequence
<220>
<223> substrate A

```

```

<400> 76
Leu Leu Trp Leu Gly Arg Val Val Gly Gly Pro Val
1             5             10

```

```

<210> 77
<211> 12
<212> PRT
<213> artificial sequence
<220>

```

<223> substrate B

<400> 77

Lys Lys Trp Leu Gly Arg Val Pro Gly Gly Pro Val  
1 5 10

<210> 78

<211> 6

<212> PRT

<213> artificial sequence

<220>

<223> variant1 SDR1

<400> 78

Asp Ala Val Gly Arg Asp  
1 5

<210> 79

<211> 6

<212> PRT

<213> artificial sequence

<220>

<223> variant2 SDR1

<400> 79

Asn Gly Arg Asp Leu Glu  
1 5

<210> 80

<211> 6

<212> PRT

<213> artificial sequence

<220>

<223> variant3 SDR1

<400> 80

Gly Phe Val Met Phe Asn  
1 5

<210> 81

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> variant1 SDR2

<400> 81

Arg Val His Pro Ser  
1 5

<210> 82

<211> 5

<212> PRT

<213> artificial sequence

<220>

<223> variant2 SDR2

<400> 82  
Val Arg Gly Thr Trp  
1 5

<210> 83  
<211> 5  
<212> PRT  
<213> artificial sequence  
<220>

<223> variant3 SDR2

<400> 83  
Arg Ser Pro Leu Thr  
1 5

<210> 84  
<211> 6  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant a SDR1

<400> 84  
Arg Pro Trp Asp Pro Ser  
1 5

<210> 85  
<211> 6  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant b SDR1

<400> 85  
Gly Phe Val Met Phe Asn  
1 5

<210> 86  
<211> 6  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant c SDR1

<400> 86  
Glu Ile Ala Asn Arg Glu  
1 5

<210> 87  
<211> 6  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant d SDR1

<400> 87  
Lys Ala Val Val Gly Thr

1 5

<210> 88  
<211> 6  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant e SDR1

<400> 88  
Val Asn Ile Met Ala Ala  
1 5

<210> 89  
<211> 6  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant f SDR1

<400> 89  
Ala Ala Phe Asn Gly Asp  
1 5

<210> 90  
<211> 5  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant a SDR2  
<400> 90

Val His Pro Thr Ser  
1 5

<210> 91  
<211> 5  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant b SDR2

<400> 91  
Arg Ser Pro Leu Thr  
1 5

<210> 92  
<211> 5  
<212> PRT  
<213> artificial sequence  
<220>  
<223> variant c SDR2

<400> 92  
Arg Gly Ala Arg Thr  
1 5

<210> 93  
 <211> 5  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> variant d SDR2

<400> 93  
 Arg Thr Pro Ile Ser  
 1 5

<210> 94  
 <211> 5  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> variant e SDR2

<400> 94  
 Thr Thr Ala Arg Lys  
 1 5

<210> 95  
 <211> 5  
 <212> PRT  
 <213> artificial sequence  
 <220>  
 <223> variant f SDR2

<400> 95  
 Arg Lys Asp Phe Trp  
 1 5

<210> 96  
 <211> 157  
 <212> PRT  
 <213> Homo sapiens

<400> 96  
 Val Arg Ser Ser Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val  
 1 5 10 15  
 Val Ala Asn Pro Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg  
 20 25 30  
 Ala Asn Ala Leu Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu  
 35 40 45  
 Val Val Pro Ser Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe  
 50 55 60  
 Lys Gly Gln Gly Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile  
 65 70 75 80  
 Ser Arg Ile Ala Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala  
 85 90 95  
 Ile Lys Ser Pro Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys  
 100 105 110  
 Pro Trp Tyr Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys  
 115 120 125  
 Gly Asp Arg Leu Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Leu Phe  
 130 135 140  
 Ala Glu Ser Gly Gln Val Tyr Phe Gly Ile Ile Ala Leu  
 145 150 155

<210> 97  
 <211> 306  
 <212> PRT  
 <213> Homo sapiens

<400> 97  
 Ile Trp Glu Leu Lys Lys Asp Val Tyr Val Val Glu Leu Asp Trp Tyr  
 1 5 10 15  
 Pro Asp Ala Pro Gly Glu Met Val Val Leu Thr Cys Asp Thr Pro Glu  
 20 25 30  
 Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln Ser Ser Glu Val Leu Gly  
 35 40 45  
 Ser Gly Lys Thr Leu Thr Ile Gln Val Lys Glu Phe Gly Asp Ala Gly  
 50 55 60  
 Gln Tyr Thr Cys His Lys Lys Gly Gly Glu Val Leu Ser His Ser Leu Leu  
 65 70 75 80  
 Leu Leu His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys  
 85 90 95  
 Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe Leu Arg Cys Glu Ala Lys  
 100 105 110  
 Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Thr Ile Ser Thr  
 115 120 125  
 Asp Leu Thr Phe Ser Val Lys Ser Ser Arg Gly Ser Ser Asp Pro Gln  
 130 135 140  
 Gly Val Thr Cys Gly Ala Ala Thr Leu Ser Ala Glu Arg Val Arg Gly  
 145 150 155 160  
 Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu Cys Gln Glu Asp Ser Ala  
 165 170 175  
 Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile Glu Val Met Val Asp Ala  
 180 185 190  
 Val His Lys Leu Lys Tyr Glu Asn Tyr Thr Ser Ser Phe Phe Ile Arg  
 195 200 205  
 Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu Gln Leu Lys Pro Leu  
 210 215 220  
 Lys Asn Ser Arg Gln Val Glu Val Ser Trp Glu Tyr Pro Asp Thr Trp  
 225 230 235 240  
 Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Val Gln Val Gln  
 245 250 255  
 Gly Lys Ser Lys Arg Glu Lys Lys Asp Arg Val Phe Thr Asp Lys Thr  
 260 265 270  
 Ser Ala Thr Val Ile Cys Arg Lys Asn Ala Ser Ile Ser Val Arg Ala  
 275 280 285  
 Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser Glu Trp Ala Ser Val Pro  
 290 295 300  
 Cys Ser  
 305

<210> 98  
 <211> 157  
 <212> PRT  
 <213> Homo sapiens

<400> 98  
 Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn  
 1 5 10 15  
 Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp  
 20 25 30  
 Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile  
 35 40 45  
 Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile  
 50 55 60



```

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile
65          70          75          80
Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys
      85          90          95
Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys
      100        105        110
Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu
      115        120        125
Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu
      130        135        140
Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp
145          150          155

```

```

<210> 99
<211> 133
<212> PRT
<213> Homo sapiens

```

```

<400> 99
Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1          5          10          15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
      20          25          30
Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
      35          40          45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
      50          55          60
Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His Leu
65          70          75          80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
      85          90          95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
      100        105        110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
      115        120        125
Ile Ser Thr Leu Thr
130

```

```

<210> 100
<211> 72
<212> PRT
<213> Homo sapiens

```

```

<400> 100
Ser Ala Lys Glu Leu Arg Cys Gln Cys Ile Lys Thr Tyr Ser Lys Pro
1          5          10          15
Phe His Pro Lys Phe Ile Lys Glu Leu Arg Val Ile Glu Ser Gly Pro
      20          25          30
His Cys Ala Asn Thr Glu Ile Ile Val Lys Leu Ser Asp Gly Arg Glu
      35          40          45
Leu Cys Leu Asp Pro Lys Glu Asn Trp Val Gln Arg Val Val Glu Lys
      50          55          60
Phe Leu Lys Arg Ala Glu Asn Ser
65          70

```

```

<210> 101
<211> 74
<212> PRT
<213> Homo sapiens

```

&lt;400&gt; 101

Gly Pro Ala Ser Val Pro Thr Thr Cys Cys Phe Asn Leu Ala Asn Arg  
 1 5 10 15  
 Lys Ile Pro Leu Gln Arg Leu Glu Ser Tyr Arg Arg Ile Thr Ser Gly  
 20 25 30  
 Lys Cys Pro Gln Lys Ala Val Ile Phe Lys Thr Lys Leu Ala Lys Asp  
 35 40 45  
 Ile Cys Ala Asp Pro Lys Lys Lys Trp Val Gln Asp Ser Met Lys Tyr  
 50 55 60  
 Leu Asp Gln Lys Ser Pro Thr Pro Lys Pro  
 65 70

&lt;210&gt; 102

&lt;211&gt; 76

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 102

Gln Pro Asp Ala Ile Asn Ala Pro Val Thr Cys Cys Tyr Asn Phe Thr  
 1 5 10 15  
 Asn Arg Lys Ile Ser Val Gln Arg Leu Ala Ser Tyr Arg Arg Ile Thr  
 20 25 30  
 Ser Ser Lys Cys Pro Lys Glu Ala Val Ile Phe Lys Thr Ile Val Ala  
 35 40 45  
 Lys Glu Ile Cys Ala Asp Pro Lys Gln Lys Trp Val Gln Asp Ser Met  
 50 55 60  
 Asp His Leu Asp Lys Gln Thr Gln Thr Pro Lys Thr  
 65 70 75

&lt;210&gt; 103

&lt;211&gt; 206

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 103

Ala Pro Met Ala Glu Gly Gly Gly Gln Asn His His Glu Val Val Lys  
 1 5 10 15  
 Phe Met Asp Val Tyr Gln Arg Ser Tyr Cys His Pro Ile Glu Thr Leu  
 20 25 30  
 Val Asp Ile Phe Gln Glu Tyr Pro Asp Glu Ile Glu Tyr Ile Phe Lys  
 35 40 45  
 Pro Ser Cys Val Pro Leu Met Arg Cys Gly Gly Cys Cys Asn Asp Glu  
 50 55 60  
 Gly Leu Glu Cys Val Pro Thr Glu Glu Ser Asn Ile Thr Met Gln Ile  
 65 70 75 80  
 Met Arg Ile Lys Pro His Gln Gly Gln His Ile Gly Glu Met Ser Phe  
 85 90 95  
 Leu Gln His Asn Lys Cys Glu Cys Arg Pro Lys Lys Asp Arg Ala Arg  
 100 105 110  
 Gln Glu Lys Lys Ser Val Arg Gly Lys Gly Lys Gly Gln Lys Arg Lys  
 115 120 125  
 Arg Lys Lys Ser Arg Tyr Lys Ser Trp Ser Val Tyr Val Gly Ala Arg  
 130 135 140  
 Cys Cys Leu Met Pro Trp Ser Leu Pro Gly Pro His Pro Cys Gly Pro  
 145 150 155 160  
 Cys Ser Glu Arg Arg Lys His Leu Phe Val Gln Asp Pro Gln Thr Cys  
 165 170 175  
 Lys Cys Ser Cys Lys Asn Thr Asp Ser Arg Cys Lys Ala Arg Gln Leu  
 180 185 190  
 Glu Leu Asn Glu Arg Thr Cys Arg Cys Asp Lys Pro Arg Arg  
 195 200 205

<210> 104  
 <211> 112  
 <212> PRT  
 <213> Homo sapiens

<400> 104

```

Ala Leu Asp Thr Asn Tyr Cys Phe Ser Ser Thr Glu Lys Asn Cys Cys
1          5          10          15
Val Arg Gln Leu Tyr Ile Asp Phe Arg Lys Asp Leu Gly Trp Lys Trp
          20          25          30
Ile His Glu Pro Lys Gly Tyr His Ala Asn Phe Cys Leu Gly Pro Cys
          35          40          45
Pro Tyr Ile Trp Ser Leu Asp Thr Gln Tyr Ser Lys Val Leu Ala Leu
          50          55          60
Tyr Asn Gln His Asn Pro Gly Ala Ser Ala Ala Pro Cys Cys Val Pro
65          70          75          80
Gln Ala Leu Glu Pro Leu Pro Ile Val Tyr Tyr Val Gly Arg Lys Pro
          85          90          95
Lys Val Glu Gln Leu Ser Asn Met Ile Val Arg Ser Cys Lys Cys Ser
          100          105          110

```

<210> 105  
 <211> 30  
 <212> PRT  
 <213> Homo sapiens

<400> 105

```

Phe Val Asn Gln His Leu Cys Gly Ser His Leu Val Glu Ala Leu Tyr
1          5          10          15
Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr Pro Lys Thr
          20          25          30

```

<210> 106  
 <211> 21  
 <212> PRT  
 <213> Homo sapiens

<400> 106

```

Gly Ile Val Glu Gln Cys Cys Thr Ser Ile Cys Ser Leu Tyr Gln Leu
1          5          10          15
Glu Asn Tyr Cys Asn
          20

```

<210> 107  
 <211> 28  
 <212> PRT  
 <213> Homo sapiens

<400> 107

```

Gly Ser Ser Phe Leu Ser Pro Glu His Gln Arg Val Gln Gln Arg Lys
1          5          10          15
Glu Ser Lys Lys Pro Pro Ala Lys Leu Gln Pro Arg
          20          25

```

<210> 108  
 <211> 9

<212> PRT  
 <213> Homo sapiens

<400> 108  
 Arg Val Tyr Ile His Pro Phe His Leu  
 1 5

<210> 109  
 <211> 114  
 <212> PRT  
 <213> Homo sapiens

<400> 109  
 Pro Met Phe Ile Val Asn Thr Asn Val Pro Arg Ala Ser Val Pro Asp  
 1 5 10 15  
 Gly Phe Leu Ser Glu Leu Thr Gln Gln Leu Ala Gln Ala Thr Gly Lys  
 20 25 30  
 Pro Pro Gln Tyr Ile Ala Val His Val Val Pro Asp Gln Leu Met Ala  
 35 40 45  
 Phe Gly Gly Ser Ser Glu Pro Cys Ala Leu Cys Ser Leu His Ser Ile  
 50 55 60  
 Gly Lys Ile Gly Gly Ala Gln Asn Arg Ser Tyr Ser Lys Leu Leu Cys  
 65 70 75 80  
 Gly Leu Leu Ala Glu Arg Leu Arg Ile Ser Pro Asp Arg Val Tyr Ile  
 85 90 95  
 Asn Tyr Tyr Asp Met Asn Ala Ala Asn Val Gly Trp Asn Asn Ser Thr  
 100 105 110  
 Phe Ala

<210> 110  
 <211> 425  
 <212> PRT  
 <213> Homo sapiens

<400> 110  
 Met Gly Pro Arg Arg Leu Leu Leu Val Ala Ala Cys Phe Ser Leu Cys  
 1 5 10 15  
 Gly Pro Leu Leu Ser Ala Arg Thr Arg Ala Arg Arg Pro Glu Ser Lys  
 20 25 30  
 Ala Thr Asn Ala Thr Leu Asp Pro Arg Ser Phe Leu Leu Arg Asn Pro  
 35 40 45  
 Asn Asp Lys Tyr Glu Pro Phe Trp Glu Asp Glu Glu Lys Asn Glu Ser  
 50 55 60  
 Gly Leu Thr Glu Tyr Arg Leu Val Ser Ile Asn Lys Ser Ser Pro Leu  
 65 70 75 80  
 Gln Lys Gln Leu Pro Ala Phe Ile Ser Glu Asp Ala Ser Gly Tyr Leu  
 85 90 95  
 Thr Ser Ser Trp Leu Thr Leu Phe Val Pro Ser Val Tyr Thr Gly Val  
 100 105 110  
 Phe Val Val Ser Leu Pro Leu Asn Ile Met Ala Ile Val Val Phe Ile  
 115 120 125  
 Leu Lys Met Lys Val Lys Lys Pro Ala Val Val Tyr Met Leu His Leu  
 130 135 140  
 Ala Thr Ala Asp Val Leu Phe Val Ser Val Leu Pro Phe Lys Ile Ser  
 145 150 155 160  
 Tyr Tyr Phe Ser Gly Ser Asp Trp Gln Phe Gly Ser Glu Leu Cys Arg  
 165 170 175  
 Phe Val Thr Ala Ala Phe Tyr Cys Asn Met Tyr Ala Ser Ile Leu Leu  
 180 185 190  
 Met Thr Val Ile Ser Ile Asp Arg Phe Leu Ala Val Val Tyr Pro Met

		195					200					205						
Gln	Ser	Leu	Ser	Trp	Arg	Thr	Leu	Gly	Arg	Ala	Ser	Phe	Thr	Cys	Leu			
	210					215					220							
Ala	Ile	Trp	Ala	Leu	Ala	Ile	Ala	Gly	Val	Val	Pro	Leu	Leu	Leu	Lys			
225					230					235					240			
Glu	Gln	Thr	Ile	Gln	Val	Pro	Gly	Leu	Asn	Ile	Thr	Thr	Cys	His	Asp			
				245					250					255				
Val	Leu	Asn	Glu	Thr	Leu	Leu	Glu	Gly	Tyr	Tyr	Ala	Tyr	Tyr	Phe	Ser			
		260					265					270						
Ala	Phe	Ser	Ala	Val	Phe	Phe	Phe	Val	Pro	Leu	Ile	Ile	Ser	Thr	Val			
	275						280				285							
Cys	Tyr	Val	Ser	Ile	Ile	Arg	Cys	Leu	Ser	Ser	Ser	Ala	Val	Ala	Asn			
	290					295				300								
Arg	Ser	Lys	Lys	Ser	Arg	Ala	Leu	Phe	Leu	Ser	Ala	Ala	Val	Phe	Cys			
305					310					315					320			
Ile	Phe	Ile	Ile	Cys	Phe	Gly	Pro	Thr	Asn	Val	Leu	Leu	Ile	Ala	His			
				325					330					335				
Tyr	Ser	Phe	Leu	Ser	His	Thr	Ser	Thr	Thr	Glu	Ala	Ala	Tyr	Phe	Ala			
		340					345					350						
Tyr	Leu	Leu	Cys	Val	Cys	Val	Ser	Ser	Ile	Ser	Cys	Cys	Ile	Asp	Pro			
	355						360				365							
Leu	Ile	Tyr	Tyr	Tyr	Ala	Ser	Ser	Glu	Cys	Gln	Arg	Tyr	Val	Tyr	Ser			
	370					375				380								
Ile	Leu	Cys	Cys	Lys	Glu	Ser	Ser	Asp	Pro	Ser	Ser	Tyr	Asn	Ser	Ser			
385				390					395					400				
Gly	Gln	Leu	Met	Ala	Ser	Lys	Met	Asp	Thr	Cys	Ser	Ser	Asn	Leu	Asn			
			405						410					415				
Asn	Ser	Ile	Tyr	Lys	Lys	Leu	Leu	Thr										
	420						425											

```
<210> 111
<211> 397
<212> PRT
<213> Homo sapiens
```

<400>	111														
Met	Arg	Ser	Pro	Ser	Ala	Ala	Trp	Leu	Leu	Gly	Ala	Ala	Ile	Leu	Leu
1				5					10					15	
Ala	Ala	Ser	Leu	Ser	Cys	Ser	Gly	Thr	Ile	Gln	Gly	Thr	Asn	Arg	Ser
			20					25					30		
Ser	Lys	Gly	Arg	Ser	Leu	Ile	Gly	Lys	Val	Asp	Gly	Thr	Ser	His	Val
		35					40					45			
Thr	Gly	Lys	Gly	Val	Thr	Val	Glu	Thr	Val	Phe	Ser	Val	Asp	Glu	Phe
	50					55				60					
Ser	Ala	Ser	Val	Leu	Thr	Gly	Lys	Leu	Thr	Thr	Val	Phe	Leu	Pro	Ile
65				70						75					80
Val	Tyr	Thr	Ile	Val	Phe	Val	Val	Gly	Leu	Pro	Ser	Asn	Gly	Met	Ala
			85					90					95		
Leu	Trp	Val	Phe	Leu	Phe	Arg	Thr	Lys	Lys	Lys	His	Pro	Ala	Val	Ile
			100					105					110		
Tyr	Met	Ala	Asn	Leu	Ala	Leu	Ala	Asp	Leu	Leu	Ser	Val	Ile	Trp	Phe
		115					120					125			
Pro	Leu	Lys	Ile	Ala	Tyr	His	Ile	His	Gly	Asn	Asn	Trp	Ile	Tyr	Gly
	130					135				140					
Glu	Ala	Leu	Cys	Asn	Val	Leu	Ile	Gly	Phe	Phe	Tyr	Gly	Asn	Met	Tyr
145				150						155					160
Cys	Ser	Ile	Leu	Phe	Met	Thr	Cys	Leu	Ser	Val	Gln	Arg	Tyr	Trp	Val
			165					170					175		
Ile	Val	Asn	Pro	Met	Gly	His	Ser	Arg	Lys	Lys	Ala	Asn	Ile	Ala	Ile
		180					185					190			
Gly	Ile	Ser	Leu	Ala	Ile	Trp	Leu	Leu	Ile	Leu	Leu	Val	Thr	Ile	Pro
	195						200					205			

```

Leu Tyr Val Val Lys Gln Thr Ile Phe Ile Pro Ala Leu Asn Ile Thr
210                215                220
Thr Cys His Asp Val Leu Pro Glu Gln Leu Leu Val Gly Asp Met Phe
225                230                235                240
Asn Tyr Phe Leu Ser Leu Ala Ile Gly Val Phe Leu Phe Pro Ala Phe
                245                250                255
Leu Thr Ala Ser Ala Tyr Val Leu Met Ile Arg Met Leu Arg Ser Ser
                260                265                270
Ala Met Asp Glu Asn Ser Glu Lys Lys Arg Lys Arg Ala Ile Lys Leu
                275                280                285
Ile Val Thr Val Leu Ala Met Tyr Leu Ile Cys Phe Thr Pro Ser Asn
290                295                300
Leu Leu Leu Val Val His Tyr Phe Leu Ile Lys Ser Gln Gly Gln Ser
305                310                315                320
His Val Tyr Ala Leu Tyr Ile Val Ala Leu Cys Leu Ser Thr Leu Asn
                325                330                335
Ser Cys Ile Asp Pro Phe Val Tyr Tyr Phe Val Ser His Asp Phe Arg
                340                345                350
Asp His Ala Lys Asn Ala Leu Leu Cys Arg Ser Val Arg Thr Val Lys
                355                360                365
Gln Met Gln Val Ser Leu Thr Ser Lys Lys His Ser Arg Lys Ser Ser
                370                375                380
Ser Tyr Ser Ser Ser Ser Thr Thr Val Lys Thr Ser Tyr
385                390                395

```

```

<210> 112
<211> 153
<212> PRT
<213> Homo sapiens

```

```

<400> 112
Ala Pro Val Arg Ser Leu Asn Cys Thr Leu Arg Asp Ser Gln Gln Lys
1          5          10          15
Ser Leu Val Met Ser Gly Pro Tyr Glu Leu Lys Ala Leu His Leu Gln
20        25        30
Gly Gln Asp Met Glu Gln Gln Val Val Phe Ser Met Ser Phe Val Gln
35        40        45
Gly Glu Glu Ser Asn Asp Lys Ile Pro Val Ala Leu Gly Leu Lys Glu
50        55        60
Lys Asn Leu Tyr Leu Ser Cys Val Leu Lys Asp Lys Pro Thr Leu
65        70        75        80
Gln Leu Glu Ser Val Asp Pro Lys Asn Tyr Pro Lys Lys Lys Met Glu
85        90        95
Lys Arg Phe Val Phe Asn Lys Ile Glu Ile Asn Asn Lys Leu Glu Phe
100       105       110
Glu Ser Ala Gln Phe Pro Asn Trp Tyr Ile Ser Thr Ser Gln Ala Glu
115       120       125
Asn Met Pro Val Phe Leu Gly Gly Thr Lys Gly Gly Gln Asp Ile Thr
130       135       140
Asp Phe Thr Met Gln Phe Val Ser Ser
145       150

```

```

<210> 113
<211> 385
<212> PRT
<213> Homo sapiens

```

```

<400> 113
Met Trp Gly Arg Leu Leu Leu Trp Pro Leu Val Leu Gly Phe Ser Leu
1          5          10          15

```

Ser Gly Gly Thr Gln Thr Pro Ser Val Tyr Asp Glu Ser Gly Ser Thr  
 20 25 30  
 Gly Gly Gly Asp Asp Ser Thr Pro Ser Ile Leu Pro Ala Pro Arg Gly  
 35 40 45  
 Tyr Pro Gly Gln Val Cys Ala Asn Asp Ser Asp Thr Leu Glu Leu Pro  
 50 55 60  
 Asp Ser Ser Arg Ala Leu Leu Gly Trp Val Pro Thr Arg Leu Val  
 65 70 75 80  
 Pro Ala Leu Tyr Gly Leu Val Leu Val Val Gly Leu Pro Ala Asn Gly  
 85 90 95  
 Leu Ala Leu Trp Val Leu Ala Thr Gln Ala Pro Arg Leu Pro Ser Thr  
 100 105 110  
 Met Leu Leu Met Asn Leu Ala Thr Ala Asp Leu Leu Leu Ala Leu Ala  
 115 120 125  
 Leu Pro Pro Arg Ile Ala Tyr His Leu Arg Gly Gln Arg Trp Pro Phe  
 130 135 140  
 Gly Glu Ala Ala Cys Arg Leu Ala Thr Ala Ala Leu Tyr Gly His Met  
 145 150 155 160  
 Tyr Gly Ser Val Leu Leu Ala Ala Val Ser Leu Asp Arg Tyr Leu  
 165 170 175  
 Ala Leu Val His Pro Leu Arg Ala Arg Ala Leu Arg Gly Arg Arg Leu  
 180 185 190  
 Ala Leu Gly Leu Cys Met Ala Ala Trp Leu Met Ala Ala Leu Ala  
 195 200 205  
 Leu Pro Leu Thr Leu Gln Arg Gln Thr Phe Arg Leu Ala Arg Ser Asp  
 210 215 220  
 Arg Val Leu Cys His Asp Ala Leu Pro Leu Asp Ala Gln Ala Ser His  
 225 230 235 240  
 Trp Gln Pro Ala Phe Thr Cys Leu Ala Leu Leu Gly Cys Phe Leu Pro  
 245 250 255  
 Leu Leu Ala Met Leu Leu Cys Tyr Gly Ala Thr Leu His Thr Leu Ala  
 260 265 270  
 Ala Ser Gly Arg Arg Tyr Gly His Ala Leu Arg Leu Thr Ala Val Val  
 275 280 285  
 Leu Ala Ser Ala Val Ala Phe Phe Val Pro Ser Asn Leu Leu Leu Leu  
 290 295 300  
 Leu His Tyr Ser Asp Pro Ser Pro Ser Ala Trp Gly Asn Leu Tyr Gly  
 305 310 315 320  
 Ala Tyr Val Pro Ser Leu Ala Leu Ser Thr Leu Asn Ser Cys Val Asp  
 325 330 335  
 Pro Phe Ile Tyr Tyr Tyr Val Ser Ala Glu Phe Arg Asp Lys Val Arg  
 340 345 350  
 Ala Gly Leu Phe Gln Arg Ser Pro Gly Asp Thr Val Ala Ser Lys Ala  
 355 360 365  
 Ser Ala Glu Gly Gly Ser Arg Gly Met Gly Thr His Ser Ser Leu Leu  
 370 375 380  
 Gln  
 385

<210> 114  
 <211> 1338  
 <212> PRT  
 <213> Homo sapiens

<400> 114  
 Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser  
 1 5 10 15  
 Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Lys Leu Lys Asp Pro  
 20 25 30  
 Glu Leu Ser Leu Lys Gly Thr Gln His Ile Met Gln Ala Gly Gln Thr  
 35 40 45  
 Leu His Leu Gln Cys Arg Gly Glu Ala Ala His Lys Trp Ser Leu Pro

	50					55					60					
Glu 65	Met	Val	Ser	Lys	Glu 70	Ser	Glu	Arg	Leu	Ser 75	Ile	Thr	Lys	Ser	Ala 80	
Cys	Gly	Arg	Asn	Gly 85	Lys	Gln	Phe	Cys	Ser 90	Thr	Leu	Thr	Leu	Asn 95	Thr	
Ala	Gln	Ala	Asn 100	His	Thr	Gly	Phe	Tyr 105	Ser	Cys	Lys	Tyr	Leu	Ala	Val	
Pro	Thr	Ser 115	Lys	Lys	Lys	Glu	Thr 120	Glu	Ser	Ala	Ile	Tyr	Ile	Phe	Ile	
Ser	Asp 130	Thr	Gly	Arg	Pro	Phe	Val 135	Glu	Met	Tyr	Ser	Glu	Ile	Pro	Glu	
Ile 145	Ile	His	Met	Thr	Glu 150	Gly	Arg	Glu	Leu	Val 155	Ile	Pro	Cys	Arg	Val	
Thr	Ser	Pro	Asn 165	Ile	Thr	Val	Thr	Leu	Lys 170	Lys	Phe	Pro	Leu	Asp	Thr	
Leu	Ile	Pro	Asp 180	Gly	Lys	Arg	Ile	Ile	Trp 185	Asp	Ser	Arg	Lys	Gly	Phe	
Ile	Ile	Ser 195	Asn	Ala	Thr	Tyr	Lys 200	Glu	Ile	Gly	Leu	Leu	Thr	Cys	Glu	
Ala	Thr 210	Val	Asn	Gly	His	Leu	Tyr 215	Lys	Thr	Asn	Tyr	Leu	Thr	His	Arg	
Gln 225	Thr	Asn	Thr	Ile	Ile	Asp	Val 230	Gln	Ile	Ser	Thr	Pro	Arg	Pro	Val	
Lys	Leu	Leu	Arg 245	Gly	His	Thr	Leu	Val	Leu 250	Asn	Cys	Thr	Ala	Thr	Thr	
Pro	Leu	Asn 260	Thr	Arg	Val	Gln	Met	Thr 265	Trp	Ser	Tyr	Pro	Asp	Glu	Lys	
Asn	Lys	Arg 275	Ala	Ser	Val	Arg	Arg	Arg 280	Ile	Asp	Gln	Ser	Asn	Ser	His	
Ala	Asn 290	Ile	Phe	Tyr	Ser	Val	Leu	Thr 295	Ile	Asp	Lys	Met	Gln	Asn	Lys	
Asp 305	Lys	Gly	Leu	Tyr	Thr	Cys	Arg	Val	Arg	Ser 315	Gly	Pro	Ser	Phe	Lys	
Ser	Val	Asn 325	Thr	Ser	Val	His	Ile	Tyr	Asp 330	Lys	Ala	Phe	Ile	Thr	Val	
Lys	His	Arg 340	Lys	Gln	Gln	Val	Leu	Glu	Thr 345	Val	Ala	Gly	Lys	Arg	Ser	
Tyr	Arg	Leu 355	Ser	Met	Lys	Val	Lys	Ala	Phe 360	Pro	Ser	Pro	Glu	Val	Val	
Trp	Leu	Lys 370	Asp	Gly	Leu	Pro	Ala	Thr	Glu 375	Lys	Ser	Ala	Arg	Tyr	Leu	
Thr 385	Arg	Gly	Tyr	Ser	Leu	Ile	Ile	Lys	Asp 390	Val	Thr	Glu	Glu	Asp	Ala	
Gly	Asn	Tyr 405	Thr	Ile	Leu	Leu	Ser	Ile	Lys 410	Gln	Ser	Asn	Val	Phe	Lys	
Asn	Leu	Thr 420	Ala	Thr	Leu	Ile	Val	Asn	Val 425	Lys	Pro	Gln	Ile	Tyr	Glu	
Lys	Ala	Val 435	Ser	Ser	Phe	Pro	Asp	Pro	Ala 440	Leu	Tyr	Pro	Leu	Gly	Ser	
Arg	Gln	Ile 450	Leu	Thr	Cys	Thr	Ala	Tyr	Gly 455	Ile	Pro	Gln	Pro	Thr	Ile	
Lys 465	Trp	Phe	Trp	His	Pro	Cys	Asn	His	Asn 470	His	Ser	Glu	Ala	Arg	Cys	
Asp	Phe	Cys 485	Ser	Asn	Asn	Glu	Glu	Ser	Phe 490	Ile	Leu	Asp	Ala	Asp	Ser	
Asn	Met	Gly 500	Asn	Arg	Ile	Glu	Ser	Ile	Thr 505	Gln	Arg	Met	Ala	Ile	Ile	
Glu	Gly	Lys 515	Asn	Lys	Met	Ala	Ser	Thr	Leu 520	Val	Val	Ala	Asp	Ser	Arg	
Ile	Ser	Gly 530	Ile	Tyr	Ile	Cys	Ile	Ala	Ser 535	Asn	Lys	Val	Gly	Thr	Val	
Gly 545	Arg	Asn	Ile	Ser	Phe	Tyr	Ile	Thr	Asp 550	Val	Pro	Asn	Gly	Phe	His	



Val Asn Leu Glu Lys Met Pro Thr Glu Gly Glu Asp Leu Lys Leu Ser  
 565 570 575  
 Cys Thr Val Asn Lys Phe Leu Tyr Arg Asp Val Thr Trp Ile Leu Leu  
 580 585 590  
 Arg Thr Val Asn Asn Arg Thr Met His Tyr Ser Ile Ser Lys Gln Lys  
 595 600 605  
 Met Ala Ile Thr Lys Glu His Ser Ile Thr Leu Asn Leu Thr Ile Met  
 610 615 620  
 Asn Val Ser Leu Gln Asp Ser Gly Thr Tyr Ala Cys Arg Ala Arg Asn  
 625 630 635 640  
 Val Tyr Thr Gly Glu Glu Ile Leu Gln Lys Lys Glu Ile Thr Ile Arg  
 645 650 655  
 Asp Gln Glu Ala Pro Tyr Leu Leu Arg Asn Leu Ser Asp His Thr Val  
 660 665 670  
 Ala Ile Ser Ser Ser Thr Thr Leu Asp Cys His Ala Asn Gly Val Pro  
 675 680 685  
 Glu Pro Gln Ile Thr Trp Phe Lys Asn Asn His Lys Ile Gln Gln Glu  
 690 695 700  
 Pro Gly Ile Ile Leu Gly Pro Gly Ser Ser Thr Leu Phe Ile Glu Arg  
 705 710 715 720  
 Val Thr Glu Glu Asp Glu Gly Val Tyr His Cys Lys Ala Thr Asn Gln  
 725 730 735  
 Lys Gly Ser Val Glu Ser Ser Ala Tyr Leu Thr Val Gln Gly Thr Ser  
 740 745 750  
 Asp Lys Ser Asn Leu Glu Leu Ile Thr Leu Thr Cys Thr Cys Val Ala  
 755 760 765  
 Ala Thr Leu Phe Trp Leu Leu Leu Thr Leu Leu Ile Arg Lys Met Lys  
 770 775 780  
 Arg Ser Ser Ser Glu Ile Lys Thr Asp Tyr Leu Ser Ile Ile Met Asp  
 785 790 795 800  
 Pro Asp Glu Val Pro Leu Asp Glu Gln Cys Glu Arg Leu Pro Tyr Asp  
 805 810 815  
 Ala Ser Lys Trp Glu Phe Ala Arg Glu Arg Leu Lys Leu Gly Lys Ser  
 820 825 830  
 Leu Gly Arg Gly Ala Phe Gly Lys Val Val Gln Ala Ser Ala Phe Gly  
 835 840 845  
 Ile Lys Lys Ser Pro Thr Cys Arg Thr Val Ala Val Lys Met Leu Lys  
 850 855 860  
 Glu Gly Ala Thr Ala Ser Glu Tyr Lys Ala Leu Met Thr Glu Leu Lys  
 865 870 875 880  
 Ile Leu Thr His Ile Gly His His Leu Asn Val Val Asn Leu Leu Gly  
 885 890 895  
 Ala Cys Thr Lys Gln Gly Gly Pro Leu Met Val Ile Val Glu Tyr Cys  
 900 905 910  
 Lys Tyr Gly Asn Leu Ser Asn Tyr Leu Lys Ser Lys Arg Asp Leu Phe  
 915 920 925  
 Phe Leu Asn Lys Asp Ala Ala Leu His Met Glu Pro Lys Lys Glu Lys  
 930 935 940  
 Met Glu Pro Gly Leu Glu Gln Gly Lys Lys Pro Arg Leu Asp Ser Val  
 945 950 955 960  
 Thr Ser Ser Glu Ser Phe Ala Ser Ser Gly Phe Gln Glu Asp Lys Ser  
 965 970 975  
 Leu Ser Asp Val Glu Glu Glu Glu Asp Ser Asp Gly Phe Tyr Lys Glu  
 980 985 990  
 Pro Ile Thr Met Glu Asp Leu Ile Ser Tyr Ser Phe Gln Val Ala Arg  
 995 1000 1005  
 Gly Met Glu Phe Leu Ser Ser Arg Lys Cys Ile His Arg Asp Leu  
 1010 1015 1020  
 Ala Ala Arg Asn Ile Leu Leu Ser Glu Asn Asn Val Val Lys Ile  
 1025 1030 1035  
 Cys Asp Phe Gly Leu Ala Arg Asp Ile Tyr Lys Asn Pro Asp Tyr  
 1040 1045 1050  
 Val Arg Lys Gly Asp Thr Arg Leu Pro Leu Lys Trp Met Ala Pro

1055	1060	1065
Glu Ser Ile Phe Asp Lys	Ile Tyr Ser Thr Lys	Ser Asp Val Trp
1070	1075	1080
Ser Tyr Gly Val Leu Leu	Trp Glu Ile Phe Ser	Leu Gly Gly Ser
1085	1090	1095
Pro Tyr Pro Gly Val Gln	Met Asp Glu Asp Phe	Cys Ser Arg Leu
1100	1105	1110
Arg Glu Gly Met Arg Met	Arg Ala Pro Glu Tyr	Ser Thr Pro Glu
1115	1120	1125
Ile Tyr Gln Ile Met Leu	Asp Cys Trp His Arg	Asp Pro Lys Glu
1130	1135	1140
Arg Pro Arg Phe Ala Glu	Leu Val Glu Lys Leu	Gly Asp Leu Leu
1145	1150	1155
Gln Ala Asn Val Gln Gln	Asp Gly Lys Asp Tyr	Ile Pro Ile Asn
1160	1165	1170
Ala Ile Leu Thr Gly Asn	Ser Gly Phe Thr Tyr	Ser Thr Pro Ala
1175	1180	1185
Phe Ser Glu Asp Phe Phe	Lys Glu Ser Ile Ser	Ala Pro Lys Phe
1190	1195	1200
Asn Ser Gly Ser Ser Asp	Asp Val Arg Tyr Val	Asn Ala Phe Lys
1205	1210	1215
Phe Met Ser Leu Glu Arg	Ile Lys Thr Phe Glu	Glu Leu Leu Pro
1220	1225	1230
Asn Ala Thr Ser Met Phe	Asp Asp Tyr Gln Gly	Asp Ser Ser Thr
1235	1240	1245
Leu Leu Ala Ser Pro Met	Leu Lys Arg Phe Thr	Trp Thr Asp Ser
1250	1255	1260
Lys Pro Lys Ala Ser Leu	Lys Ile Asp Leu Arg	Val Thr Ser Lys
1265	1270	1275
Ser Lys Glu Ser Gly Leu	Ser Asp Val Ser Arg	Pro Ser Phe Cys
1280	1285	1290
His Ser Ser Cys Gly His	Val Ser Glu Gly Lys	Arg Arg Phe Thr
1295	1300	1305
Tyr Asp His Ala Glu Leu	Glu Arg Lys Ile Ala	Cys Cys Ser Pro
1310	1315	1320
Pro Pro Asp Tyr Asn Ser	Val Val Leu Tyr Ser	Thr Pro Pro Ile
1325	1330	1335

<210> 115  
 <211> 1356  
 <212> PRT  
 <213> Homo sapiens

<400> 115  
 Met Gln Ser Lys Val Leu Leu Ala Val Ala Leu Trp Leu Cys Val Glu  
 1 5 10 15  
 Thr Arg Ala Ala Ser Val Gly Leu Pro Ser Val Ser Leu Asp Leu Pro  
 20 25 30  
 Arg Leu Ser Ile Gln Lys Asp Ile Leu Thr Ile Lys Ala Asn Thr Thr  
 35 40 45  
 Leu Gln Ile Thr Cys Arg Gly Gln Arg Asp Leu Asp Trp Leu Trp Pro  
 50 55 60  
 Asn Asn Gln Ser Gly Ser Glu Gln Arg Val Glu Val Thr Glu Cys Ser  
 65 70 75 80  
 Asp Gly Leu Phe Cys Lys Thr Leu Thr Ile Pro Lys Val Ile Gly Asn  
 85 90 95  
 Asp Thr Gly Ala Tyr Lys Cys Phe Tyr Arg Glu Thr Asp Leu Ala Ser  
 100 105 110  
 Val Ile Tyr Val Tyr Val Gln Asp Tyr Arg Ser Pro Phe Ile Ala Ser  
 115 120 125  
 Val Ser Asp Gln His Gly Val Val Tyr Ile Thr Glu Asn Lys Asn Lys  
 130 135 140

Thr	Val	Val	Ile	Pro	Cys	Leu	Gly	Ser	Ile	Ser	Asn	Leu	Asn	Val	Ser
145					150					155					160
Leu	Cys	Ala	Arg	Tyr	Pro	Glu	Lys	Arg	Phe	Val	Pro	Asp	Gly	Asn	Arg
				165					170					175	
Ile	Ser	Trp	Asp	Ser	Lys	Lys	Gly	Phe	Thr	Ile	Pro	Ser	Tyr	Met	Ile
			180					185					190		
Ser	Tyr	Ala	Gly	Met	Val	Phe	Cys	Glu	Ala	Lys	Ile	Asn	Asp	Glu	Ser
		195					200					205			
Tyr	Gln	Ser	Ile	Met	Tyr	Ile	Val	Val	Val	Val	Gly	Tyr	Arg	Ile	Tyr
	210					215					220				
Asp	Val	Val	Leu	Ser	Pro	Ser	His	Gly	Ile	Glu	Leu	Ser	Val	Gly	Glu
225					230					235					240
Lys	Leu	Val	Leu	Asn	Cys	Thr	Ala	Arg	Thr	Glu	Leu	Asn	Val	Gly	Ile
				245					250					255	
Asp	Phe	Asn	Trp	Glu	Tyr	Pro	Ser	Ser	Lys	His	Gln	His	Lys	Lys	Leu
			260					265					270		
Val	Asn	Arg	Asp	Leu	Lys	Thr	Gln	Ser	Gly	Ser	Glu	Met	Lys	Lys	Phe
		275					280					285			
Leu	Ser	Thr	Leu	Thr	Ile	Asp	Gly	Val	Thr	Arg	Ser	Asp	Gln	Gly	Leu
	290					295					300				
Tyr	Thr	Cys	Ala	Ala	Ser	Ser	Gly	Leu	Met	Thr	Lys	Lys	Asn	Ser	Thr
305					310					315					320
Phe	Val	Arg	Val	His	Glu	Lys	Pro	Phe	Val	Ala	Phe	Gly	Ser	Gly	Met
				325					330					335	
Glu	Ser	Leu	Val	Glu	Ala	Thr	Val	Gly	Glu	Arg	Val	Arg	Ile	Pro	Ala
			340					345					350		
Lys	Tyr	Leu	Gly	Tyr	Pro	Pro	Pro	Glu	Ile	Lys	Trp	Tyr	Lys	Asn	Gly
		355					360					365			
Ile	Pro	Leu	Glu	Ser	Asn	His	Thr	Ile	Lys	Ala	Gly	His	Val	Leu	Thr
	370					375					380				
Ile	Met	Glu	Val	Ser	Glu	Arg	Asp	Thr	Gly	Asn	Tyr	Thr	Val	Ile	Leu
385					390					395					400
Thr	Asn	Pro	Ile	Ser	Lys	Glu	Lys	Gln	Ser	His	Val	Val	Ser	Leu	Val
				405					410					415	
Val	Tyr	Val	Pro	Pro	Gln	Ile	Gly	Glu	Lys	Ser	Leu	Ile	Ser	Pro	Val
			420					425					430		
Asp	Ser	Tyr	Gln	Tyr	Gly	Thr	Thr	Gln	Thr	Leu	Thr	Cys	Thr	Val	Tyr
		435					440					445			
Ala	Ile	Pro	Pro	Pro	His	His	Ile	His	Trp	Tyr	Trp	Gln	Leu	Glu	Glu
	450					455					460				
Glu	Cys	Ala	Asn	Glu	Pro	Ser	Gln	Ala	Val	Ser	Val	Thr	Asn	Pro	Tyr
465					470					475					480
Pro	Cys	Glu	Glu	Trp	Arg	Ser	Val	Glu	Asp	Phe	Gln	Gly	Gly	Asn	Lys
				485					490					495	
Ile	Glu	Val	Asn	Lys	Asn	Gln	Phe	Ala	Leu	Ile	Glu	Gly	Lys	Asn	Lys
			500					505					510		
Thr	Val	Ser	Thr	Leu	Val	Ile	Gln	Ala	Ala	Asn	Val	Ser	Ala	Leu	Tyr
		515					520					525			
Lys	Cys	Glu	Ala	Val	Asn	Lys	Val	Gly	Arg	Gly	Glu	Arg	Val	Ile	Ser
		530				535					540				
Phe	His	Val	Thr	Arg	Gly	Pro	Glu	Ile	Thr	Leu	Gln	Pro	Asp	Met	Gln
545					550					555					560
Pro	Thr	Glu	Gln	Glu	Ser	Val	Ser	Leu	Trp	Cys	Thr	Ala	Asp	Arg	Ser
				565					570					575	
Thr	Phe	Glu	Asn	Leu	Thr	Trp	Tyr	Lys	Leu	Gly	Pro	Gln	Pro	Leu	Pro
			580					585					590		
Ile	His	Val	Gly	Glu	Leu	Pro	Thr	Pro	Val	Cys	Lys	Asn	Leu	Asp	Thr
		595					600						605		
Leu	Trp	Lys	Leu	Asn	Ala	Thr	Met	Phe	Ser	Asn	Ser	Thr	Asn	Asp	Ile
	610					615					620				
Leu	Ile	Met	Glu	Leu	Lys	Asn	Ala	Ser	Leu	Gln	Asp	Gln	Gly	Asp	Tyr
625					630					635					640
Val	Cys	Leu	Ala	Gln	Asp	Arg	Lys	Thr	Lys	Lys	Arg	His	Cys	Val	Val

				645					650					655		
Arg	Gln	Leu	Thr	Val	Leu	Glu	Arg	Val	Ala	Pro	Thr	Ile	Thr	Gly	Asn	
			660					665					670			
Leu	Glu	Asn	Gln	Thr	Thr	Ser	Ile	Gly	Glu	Ser	Ile	Glu	Val	Ser	Cys	
		675					680					685				
Thr	Ala	Ser	Gly	Asn	Pro	Pro	Pro	Gln	Ile	Met	Trp	Phe	Lys	Asp	Asn	
	690					695				700						
Glu	Thr	Leu	Val	Glu	Asp	Ser	Gly	Ile	Val	Leu	Lys	Asp	Gly	Asn	Arg	
705				710					715					720		
Asn	Leu	Thr	Ile	Arg	Arg	Val	Arg	Lys	Glu	Asp	Glu	Gly	Leu	Tyr	Thr	
			725					730					735			
Cys	Gln	Ala	Cys	Ser	Val	Leu	Gly	Cys	Ala	Lys	Val	Glu	Ala	Phe	Phe	
		740					745					750				
Ile	Ile	Glu	Gly	Ala	Gln	Glu	Lys	Thr	Asn	Leu	Glu	Ile	Ile	Ile	Leu	
	755					760					765					
Val	Gly	Thr	Ala	Val	Ile	Ala	Met	Phe	Phe	Trp	Leu	Leu	Leu	Val	Ile	
	770					775					780					
Ile	Leu	Arg	Thr	Val	Lys	Arg	Ala	Asn	Gly	Gly	Glu	Leu	Lys	Thr	Gly	
785				790					795					800		
Tyr	Leu	Ser	Ile	Val	Met	Asp	Pro	Asp	Glu	Leu	Pro	Leu	Asp	Glu	His	
			805					810					815			
Cys	Glu	Arg	Leu	Pro	Tyr	Asp	Ala	Ser	Lys	Trp	Glu	Phe	Pro	Arg	Asp	
		820					825					830				
Arg	Leu	Lys	Leu	Gly	Lys	Pro	Leu	Gly	Arg	Gly	Ala	Phe	Gly	Gln	Val	
	835					840					845					
Ile	Glu	Ala	Asp	Ala	Phe	Gly	Ile	Asp	Lys	Thr	Ala	Thr	Cys	Arg	Thr	
	850					855				860						
Val	Ala	Val	Lys	Met	Leu	Lys	Glu	Gly	Ala	Thr	His	Ser	Glu	His	Arg	
865				870					875					880		
Ala	Leu	Met	Ser	Glu	Leu	Lys	Ile	Leu	Ile	His	Ile	Gly	His	His	Leu	
			885					890					895			
Asn	Val	Val	Asn	Leu	Leu	Gly	Ala	Cys	Thr	Lys	Pro	Gly	Gly	Pro	Leu	
		900					905					910				
Met	Val	Ile	Val	Glu	Phe	Cys	Lys	Phe	Gly	Asn	Leu	Ser	Thr	Tyr	Leu	
	915					920					925					
Arg	Ser	Lys	Arg	Asn	Glu	Phe	Val	Pro	Tyr	Lys	Thr	Lys	Gly	Ala	Arg	
	930					935				940						
Phe	Arg	Gln	Gly	Lys	Asp	Tyr	Val	Gly	Ala	Ile	Pro	Val	Asp	Leu	Lys	
945				950					955					960		
Arg	Arg	Leu	Asp	Ser	Ile	Thr	Ser	Ser	Gln	Ser	Ser	Ala	Ser	Ser	Gly	
			965					970					975			
Phe	Val	Glu	Glu	Lys	Ser	Leu	Ser	Asp	Val	Glu	Glu	Glu	Glu	Ala	Pro	
	980						985									

His Gly Glu Pro Ser Gln Arg Pro Thr Phe Ser Glu Leu Val Glu  
 1145 1150 1155  
 His Leu Gly Asn Leu Leu Gln Ala Asn Ala Gln Gln Asp Gly Lys  
 1160 1165 1170  
 Asp Tyr Ile Val Leu Pro Ile Ser Glu Thr Leu Ser Met Glu Glu  
 1175 1180 1185  
 Asp Ser Gly Leu Ser Leu Pro Thr Ser Pro Val Ser Cys Met Glu  
 1190 1195 1200  
 Glu Glu Glu Val Cys Asp Pro Lys Phe His Tyr Asp Asn Thr Ala  
 1205 1210 1215  
 Gly Ile Ser Gln Tyr Leu Gln Asn Ser Lys Arg Lys Ser Arg Pro  
 1220 1225 1230  
 Val Ser Val Lys Thr Phe Glu Asp Ile Pro Leu Glu Glu Pro Glu  
 1235 1240 1245  
 Val Lys Val Ile Pro Asp Asp Asn Gln Thr Asp Ser Gly Met Val  
 1250 1255 1260  
 Leu Ala Ser Glu Glu Leu Lys Thr Leu Glu Asp Arg Thr Lys Leu  
 1265 1270 1275  
 Ser Pro Ser Phe Gly Gly Met Val Pro Ser Lys Ser Arg Glu Ser  
 1280 1285 1290  
 Val Ala Ser Glu Gly Ser Asn Gln Thr Ser Gly Tyr Gln Ser Gly  
 1295 1300 1305  
 Tyr His Ser Asp Asp Thr Asp Thr Thr Val Tyr Ser Ser Glu Glu  
 1310 1315 1320  
 Ala Glu Leu Leu Lys Leu Ile Glu Ile Gly Val Gln Thr Gly Ser  
 1325 1330 1335  
 Thr Ala Gln Ile Leu Gln Pro Asp Ser Gly Thr Thr Leu Ser Ser  
 1340 1345 1350  
 Pro Pro Val  
 1355

<210> 116  
 <211> 1186  
 <212> PRT  
 <213> Homo sapiens

<400> 116  
 Leu Glu Glu Lys Lys Val Cys Gln Gly Thr Ser Asn Lys Leu Thr Gln  
 1 5 10 15  
 Leu Gly Thr Phe Glu Asp His Phe Leu Ser Leu Gln Arg Met Phe Asn  
 20 25 30  
 Asn Cys Glu Val Val Leu Gly Asn Leu Glu Ile Thr Tyr Val Gln Arg  
 35 40 45  
 Asn Tyr Asp Leu Ser Phe Leu Lys Thr Ile Gln Glu Val Ala Gly Tyr  
 50 55 60  
 Val Leu Ile Ala Leu Asn Thr Val Glu Arg Ile Pro Leu Glu Asn Leu  
 65 70 75 80  
 Gln Ile Ile Arg Gly Asn Met Tyr Tyr Glu Asn Ser Tyr Ala Leu Ala  
 85 90 95  
 Val Leu Ser Asn Tyr Asp Ala Asn Lys Thr Gly Leu Lys Glu Leu Pro  
 100 105 110  
 Met Arg Asn Leu Gln Glu Ile Leu His Gly Ala Val Arg Phe Ser Asn  
 115 120 125  
 Asn Pro Ala Leu Cys Asn Val Glu Ser Ile Gln Trp Arg Asp Ile Val  
 130 135 140  
 Ser Ser Asp Phe Leu Ser Asn Met Ser Met Asp Phe Gln Asn His Leu  
 145 150 155 160  
 Gly Ser Cys Gln Lys Cys Asp Pro Ser Cys Pro Asn Gly Ser Cys Trp  
 165 170 175  
 Gly Ala Gly Glu Glu Asn Cys Gln Lys Leu Thr Lys Ile Ile Cys Ala  
 180 185 190  
 Gln Gln Cys Ser Gly Arg Cys Arg Gly Lys Ser Pro Ser Asp Cys Cys

	195					200					205				
His	Asn	Gln	Cys	Ala	Ala	Gly	Cys	Thr	Gly	Pro	Arg	Glu	Ser	Asp	Cys
	210					215					220				
Leu	Val	Cys	Arg	Lys	Phe	Arg	Asp	Glu	Ala	Thr	Cys	Lys	Asp	Thr	Cys
225					230					235					240
Pro	Pro	Leu	Met	Leu	Tyr	Asn	Pro	Thr	Thr	Tyr	Gln	Met	Asp	Val	Asn
				245					250					255	
Pro	Glu	Gly	Lys	Tyr	Ser	Phe	Gly	Ala	Thr	Cys	Val	Lys	Lys	Cys	Pro
			260				265						270		
Arg	Asn	Tyr	Val	Val	Thr	Asp	His	Gly	Ser	Cys	Val	Arg	Ala	Cys	Gly
		275					280					285			
Ala	Asp	Ser	Tyr	Glu	Met	Glu	Glu	Asp	Gly	Val	Arg	Lys	Cys	Lys	Lys
	290					295					300				
Cys	Glu	Gly	Pro	Cys	Arg	Lys	Val	Cys	Asn	Gly	Ile	Gly	Ile	Gly	Glu
305					310					315					320
Phe	Lys	Asp	Ser	Leu	Ser	Ile	Asn	Ala	Thr	Asn	Ile	Lys	His	Phe	Lys
				325					330					335	
Asn	Cys	Thr	Ser	Ile	Ser	Gly	Asp	Leu	His	Ile	Leu	Pro	Val	Ala	Phe
			340				345						350		
Arg	Gly	Asp	Ser	Phe	Thr	His	Thr	Pro	Pro	Leu	Asp	Pro	Gln	Glu	Leu
		355					360					365			
Asp	Ile	Leu	Lys	Thr	Val	Lys	Glu	Ile	Thr	Gly	Phe	Leu	Leu	Ile	Gln
	370					375					380				
Ala	Trp	Pro	Glu	Asn	Arg	Thr	Asp	Leu	His	Ala	Phe	Glu	Asn	Leu	Glu
385					390					395					400
Ile	Ile	Arg	Gly	Arg	Thr	Lys	Gln	His	Gly	Gln	Phe	Ser	Leu	Ala	Val
				405					410					415	
Val	Ser	Leu	Asn	Ile	Thr	Ser	Leu	Gly	Leu	Arg	Ser	Leu	Lys	Glu	Ile
			420					425					430		
Ser	Asp	Gly	Asp	Val	Ile	Ile	Ser	Gly	Asn	Lys	Asn	Leu	Cys	Tyr	Ala
		435					440				445				
Asn	Thr	Ile	Asn	Trp	Lys	Lys	Leu	Phe	Gly	Thr	Ser	Gly	Gln	Lys	Thr
					455						460				
Lys	Ile	Ile	Ser	Asn	Arg	Gly	Glu	Asn	Ser	Cys	Lys	Ala	Thr	Gly	Gln
465					470					475					480
Val	Cys	His	Ala	Leu	Cys	Ser	Pro	Glu	Gly	Cys	Trp	Gly	Pro	Glu	Pro
				485					490					495	
Arg	Asp	Cys	Val	Ser	Cys	Arg	Asn	Val	Ser	Arg	Gly	Arg	Glu	Cys	Val
			500					505					510		
Asp	Lys	Cys	Asn	Leu	Leu	Glu	Gly	Glu	Pro	Arg	Glu	Phe	Val	Glu	Asn
		515					520				525				
Ser	Glu	Cys	Ile	Gln	Cys	His	Pro	Glu	Cys	Leu	Pro	Gln	Ala	Met	Asn
	530					535					540				
Ile	Thr	Cys	Thr	Gly	Arg	Gly	Pro	Asp	Asn	Cys	Ile	Gln	Cys	Ala	His
545					550					555					56

Gly	Leu	Trp	Ile	Pro	Glu	Gly	Glu	Lys	Val	Lys	Ile	Pro	Val	Ala	Ile		
705					710					715					720		
Lys	Glu	Leu	Arg	Glu	Ala	Thr	Ser	Pro	Lys	Ala	Asn	Lys	Glu	Ile	Leu		
				725					730						735		
Asp	Glu	Ala	Tyr	Val	Met	Ala	Ser	Val	Asp	Asn	Pro	His	Val	Cys	Arg		
			740					745					750				
Leu	Leu	Gly	Ile	Cys	Leu	Thr	Ser	Thr	Val	Gln	Leu	Ile	Thr	Gln	Leu		
		755						760				765					
Met	Pro	Phe	Gly	Cys	Leu	Leu	Asp	Tyr	Val	Arg	Glu	His	Lys	Asp	Asn		
		770					775				780						
Ile	Gly	Ser	Gln	Tyr	Leu	Leu	Asn	Trp	Cys	Val	Gln	Ile	Ala	Lys	Gly		
785					790					795					800		
Met	Asn	Tyr	Leu	Glu	Asp	Arg	Arg	Leu	Val	His	Arg	Asp	Leu	Ala	Ala		
				805					810						815		
Arg	Asn	Val	Leu	Val	Lys	Thr	Pro	Gln	His	Val	Lys	Ile	Thr	Asp	Phe		
			820					825					830				
Gly	Leu	Ala	Lys	Leu	Leu	Gly	Ala	Glu	Glu	Lys	Glu	Tyr	His	Ala	Glu		
		835					840					845					
Gly	Gly	Lys	Val	Pro	Ile	Lys	Trp	Met	Ala	Leu	Glu	Ser	Ile	Leu	His		
	850					855					860						
Arg	Ile	Tyr	Thr	His	Gln	Ser	Asp	Val	Trp	Ser	Tyr	Gly	Val	Thr	Val		
865					870					875					880		
Trp	Glu	Leu	Met	Thr	Phe	Gly	Ser	Lys	Pro	Tyr	Asp	Gly	Ile	Pro	Ala		
				885					890						895		
Ser	Glu	Ile	Ser	Ser	Ile	Leu	Glu	Lys	Gly	Glu	Arg	Leu	Pro	Gln	Pro		
			900					905					910				
Pro	Ile	Cys	Thr	Ile	Asp	Val	Tyr	Met	Ile	Met	Val	Lys	Cys	Trp	Met		
		915					920					925					
Ile	Asp	Ala	Asp	Ser	Arg	Pro	Lys	Phe	Arg	Glu	Leu	Ile	Ile	Glu	Phe		
	930					935					940						
Ser	Lys	Met	Ala	Arg	Asp	Pro	Gln	Arg	Tyr	Leu	Val	Ile	Gln	Gly	Asp		
945					950					955					960		
Glu	Arg	Met	His	Leu	Pro	Ser	Pro	Thr	Asp	Ser	Asn	Phe	Tyr	Arg	Ala		
				965					970						975		
Leu	Met	Asp	Glu	Glu	Asp	Met	Asp	Asp	Val	Val	Asp	Ala	Asp	Glu	Tyr		
		980						985					990				
Leu	Ile	Pro	Gln	Gln	Gly	Phe	Phe	Ser	Ser	Pro	Ser	Thr	Ser	Arg	Thr		
		995					1000						1005				
Pro	Leu	Leu	Ser	Ser	Leu	Ser	Ala	Thr	Ser	Asn	Asn	Ser	Thr	Val			
	1010					1015						1020					
Ala	Cys	Ile	Asp	Arg	Asn	Gly	Leu	Gln	Ser	Cys	Pro	Ile	Lys	Glu			
	1025					1030						1035					
Asp	Ser	Phe	Leu	Gln	Arg	Tyr	Ser	Ser	Asp	Pro	Thr	Gly	Ala	Leu			
	1040					1045						1050					
Thr	Glu	Asp	Ser	Ile	Asp	Asp	Thr	Phe	Leu	Pro	Val	Pro	Glu	Tyr			
	1055					1060						1065					
Ile	Asn	Gln	Ser	Val	Pro	Lys	Arg	Pro	Ala	Gly	Ser	Val	Gln	Asn			
	1070					1075						1080					
Pro	Val	Tyr	His	Asn	Gln	Pro	Leu	Asn	Pro	Ala	Pro	Ser	Arg	Asp			
	1085					1090						1095					
Pro	His	Tyr	Gln	Asp	Pro	His	Ser	Thr	Ala	Val	Gly	Asn	Pro	Glu			
	1100					1105						1110					
Tyr	Leu	Asn	Thr	Val	Gln	Pro	Thr	Cys	Val	Asn	Ser	Thr	Phe	Asp			
	1115					1120						1125					
Ser	Pro	Ala	His	Trp	Ala	Gln	Lys	Gly	Ser	His	Gln	Ile	Ser	Leu			
	1130					1135						1140					
Asp	Asn	Pro	Asp	Tyr	Gln	Gln	Asp	Phe	Phe	Pro	Lys	Glu	Ala	Lys			
	1145					1150						1155					
Pro	Asn	Gly	Ile	Phe	Lys	Gly	Ser	Thr	Ala	Glu	Asn	Ala	Glu	Tyr			
	1160					1165						1170					
Leu	Arg	Val	Ala	Pro	Gln	Ser	Ser	Glu	Phe	Ile	Gly	Ala					
	1175					1180						1185					

<400>	117														
Met	Asp	Val	Leu	Ser	Pro	Gly	Gln	Gly	Asn	Asn	Thr	Thr	Ser	Pro	Pro
1				5					10					15	
Ala	Pro	Phe	Glu	Thr	Gly	Gly	Asn	Thr	Thr	Gly	Ile	Ser	Asp	Val	Thr
			20					25					30		
Val	Ser	Tyr	Gln	Val	Ile	Thr	Ser	Leu	Leu	Leu	Gly	Thr	Leu	Ile	Phe
		35					40					45			
Cys	Ala	Val	Leu	Gly	Asn	Ala	Cys	Val	Val	Ala	Ala	Ile	Ala	Leu	Glu
	50					55				60					
Arg	Ser	Leu	Gln	Asn	Val	Ala	Asn	Tyr	Leu	Ile	Gly	Ser	Leu	Ala	Val
65					70					75					80
Thr	Asp	Leu	Met	Val	Ser	Val	Leu	Val	Leu	Pro	Met	Ala	Ala	Leu	Tyr
				85					90					95	
Gln	Val	Leu	Asn	Lys	Trp	Thr	Leu	Gly	Gln	Val	Thr	Cys	Asp	Leu	Phe
			100					105					110		
Ile	Ala	Leu	Asp	Val	Leu	Cys	Cys	Thr	Ser	Ser	Ile	Leu	His	Leu	Cys
		115					120					125			
Ala	Ile	Ala	Leu	Asp	Arg	Tyr	Trp	Ala	Ile	Thr	Asp	Pro	Ile	Asp	Tyr
		130				135					140				
Val	Asn	Lys	Arg	Thr	Pro	Arg	Arg	Ala	Ala	Ala	Leu	Ile	Ser	Leu	Thr
145					150					155					160
Trp	Leu	Ile	Gly	Phe	Leu	Ile	Ser	Ile	Pro	Pro	Met	Leu	Gly	Trp	Arg
				165					170					175	
Thr	Pro	Glu	Asp	Arg	Ser	Asp	Pro	Asp	Ala	Cys	Thr	Ile	Ser	Lys	Asp
			180					185					190		
His	Gly	Tyr	Thr	Ile	Tyr	Ser	Thr	Phe	Gly	Ala	Phe	Tyr	Ile	Pro	Leu
		195					200					205			
Leu	Leu	Met	Leu	Val	Leu	Tyr	Gly	Arg	Ile	Phe	Arg	Ala	Ala	Arg	Phe
		210				215					220				
Arg	Ile	Arg	Lys	Thr	Val	Lys	Lys	Val	Glu	Lys	Thr	Gly	Ala	Asp	Thr
225					230					235					240
Arg	His	Gly	Ala	Ser	Pro	Ala	Pro	Gln	Pro	Lys	Lys	Ser	Val	Asn	Gly
				245					250					255	
Glu	Ser	Gly	Ser	Arg	Asn	Trp	Arg	Leu	Gly	Val	Glu	Ser	Lys	Ala	Gly
			260					265					270		
Gly	Ala	Leu	Cys	Ala	Asn	Gly	Ala	Val	Arg	Gln	Gly	Asp	Asp	Gly	Ala
		275					280					285			
Ala	Leu	Glu	Val	Ile	Glu	Val	His	Arg	Val	Gly	Asn	Ser	Lys	Glu	His
		290				295					300				
Leu	Pro	Leu	Pro	Ser	Glu	Ala	Gly	Pro	Thr	Pro	Cys	Ala	Pro	Ala	Ser
305					310					315					320
Phe	Glu	Arg	Lys	Asn	Glu	Arg	Asn	Ala	Glu	Ala	Lys	Arg	Lys	Met	Ala
				325					330					335	
Leu	Ala	Arg	Glu	Arg	Lys	Thr	Val	Lys	Thr	Leu	Gly	Ile	Ile	Met	Gly
			340												



<210> 118  
 <211> 129  
 <212> PRT  
 <213> Homo sapiens

<400> 118  
 His Lys Cys Asp Ile Thr Leu Gln Glu Ile Ile Lys Thr Leu Asn Ser  
 1 5 10 15  
 Leu Thr Glu Gln Lys Thr Leu Cys Thr Glu Leu Thr Val Thr Asp Ile  
 20 25 30  
 Phe Ala Ala Ser Lys Asn Thr Thr Glu Lys Glu Thr Phe Cys Arg Ala  
 35 40 45  
 Ala Thr Val Leu Arg Gln Phe Tyr Ser His His Glu Lys Asp Thr Arg  
 50 55 60  
 Cys Leu Gly Ala Thr Ala Gln Gln Phe His Arg His Lys Gln Leu Ile  
 65 70 75 80  
 Arg Phe Leu Lys Arg Leu Asp Arg Asn Leu Trp Gly Leu Ala Gly Leu  
 85 90 95  
 Asn Ser Cys Pro Val Lys Glu Ala Asn Gln Ser Thr Leu Glu Asn Phe  
 100 105 110  
 Leu Glu Arg Leu Lys Thr Ile Met Arg Glu Lys Tyr Ser Lys Cys Ser  
 115 120 125  
 Ser

<210> 119  
 <211> 113  
 <212> PRT  
 <213> Homo sapiens

<400> 119  
 Met Gly Pro Val Pro Pro Ser Thr Ala Leu Arg Glu Leu Ile Glu Glu  
 1 5 10 15  
 Leu Val Asn Ile Thr Gln Asn Gln Lys Ala Pro Leu Cys Asn Gly Ser  
 20 25 30  
 Met Val Trp Ser Ile Asn Leu Thr Ala Gly Met Tyr Cys Ala Ala Leu  
 35 40 45  
 Glu Ser Leu Ile Asn Val Ser Gly Cys Ser Ala Ile Glu Lys Thr Gln  
 50 55 60  
 Arg Met Leu Ser Gly Phe Cys Pro His Lys Val Ser Ala Gly Gln Phe  
 65 70 75 80  
 Ser Ser Leu His Val Arg Asp Thr Lys Ile Glu Val Ala Gln Phe Val  
 85 90 95  
 Lys Asp Leu Leu Leu His Leu Lys Lys Leu Phe Arg Glu Gly Arg Phe  
 100 105 110  
 Asn

<210> 120  
 <211> 726  
 <212> PRT  
 <213> Homo sapiens

<400> 120  
 Val Thr Lys Leu Leu Pro Ala Leu Leu Leu Gln His Val Leu Leu His  
 1 5 10 15  
 Leu Leu Leu Leu Pro Ile Ala Ile Pro Tyr Ala Glu Gly Gln Arg Lys  
 20 25 30  
 Arg Arg Asn Thr Ile His Glu Phe Lys Lys Ser Ala Lys Thr Thr Leu  
 35 40 45  
 Ile Lys Ile Asp Pro Ala Leu Lys Ile Lys Thr Lys Lys Val Asn Thr  
 50 55 60  
 Ala Asp Gln Cys Ala Asn Arg Cys Thr Arg Asn Lys Gly Leu Pro Phe

65					70					75					80
Thr	Cys	Lys	Ala	Phe	Val	Phe	Asp	Lys	Ala	Arg	Lys	Gln	Cys	Leu	Trp
			85						90					95	
Phe	Pro	Phe	Asn	Ser	Met	Ser	Ser	Gly	Val	Lys	Lys	Glu	Phe	Gly	His
			100					105					110		
Glu	Phe	Asp	Leu	Tyr	Glu	Asn	Lys	Asp	Tyr	Ile	Arg	Asn	Cys	Ile	Ile
		115					120					125			
Gly	Lys	Gly	Arg	Ser	Tyr	Lys	Gly	Thr	Val	Ser	Ile	Thr	Lys	Ser	Gly
	130					135					140				
Ile	Lys	Cys	Gln	Pro	Trp	Ser	Ser	Met	Ile	Pro	His	Glu	His	Ser	Phe
145					150					155					160
Leu	Pro	Ser	Ser	Tyr	Arg	Gly	Lys	Asp	Leu	Gln	Glu	Asn	Tyr	Cys	Arg
				165					170					175	
Asn	Pro	Arg	Gly	Glu	Glu	Gly	Gly	Pro	Trp	Cys	Phe	Thr	Ser	Asn	Pro
			180					185					190		
Glu	Val	Arg	Tyr	Glu	Val	Cys	Asp	Ile	Pro	Gln	Cys	Ser	Glu	Val	Glu
		195					200					205			
Cys	Met	Thr	Cys	Asn	Gly	Glu	Ser	Tyr	Arg	Gly	Leu	Met	Asp	His	Thr
	210				215						220				
Glu	Ser	Gly	Lys	Ile	Cys	Gln	Arg	Trp	Asp	His	Gln	Thr	Pro	His	Arg
225				230						235					240
His	Lys	Phe	Leu	Pro	Glu	Arg	Tyr	Pro	Asp	Lys	Gly	Phe	Asp	Asp	Asn
				245					250					255	
Tyr	Cys	Arg	Asn	Pro	Asp	Gly	Gln	Pro	Arg	Pro	Trp	Cys	Tyr	Thr	Leu
			260					265					270		
Asp	Pro	His	Thr	Arg	Trp	Glu	Tyr	Cys	Ala	Ile	Lys	Thr	Cys	Ala	Asp
		275					280					285			
Asn	Thr	Met	Asn	Asp	Thr	Asp	Val	Pro	Leu	Glu	Thr	Thr	Glu	Cys	Ile
	290					295					300				
Gln	Gly	Gln	Gly	Glu	Gly	Tyr	Arg	Gly	Thr	Val	Asn	Thr	Ile	Trp	Asn
305				310						315					320
Gly	Ile	Pro	Cys	Gln	Arg	Trp	Asp	Ser	Gln	Tyr	Pro	His	Glu	His	Asp
				325					330					335	
Met	Thr	Pro	Glu	Asn	Phe	Lys	Cys	Lys	Asp	Leu	Arg	Glu	Asn	Tyr	Cys
			340					345					350		
Arg	Asn	Pro	Asp	Gly	Ser	Glu	Ser	Pro	Trp	Cys	Phe	Thr	Thr	Asp	Pro
		355					360					365			
Asn	Ile	Arg	Val	Gly	Tyr	Cys	Ser	Gln	Ile	Pro	Asn	Cys	Asp	Met	Ser
	370					375					380				
His	Gly	Gln	Asp	Cys	Tyr	Arg	Gly	Asn	Gly	Lys	Asn	Tyr	Met	Gly	Asn
385				390						395					400
Leu	Ser	Gln	Thr	Arg	Ser	Gly	Leu	Thr	Cys	Ser	Met	Trp	Asp	Lys	Asn
				405					410					415	
Met	Glu	Asp	Leu	His	Arg	His	Ile	Phe	Trp	Glu	Pro	Asp	Ala	Ser	Lys
			420					425					430		

Leu Val Leu Met Lys Leu Ala Arg Pro Ala Val Leu Asp Asp Phe Val  
                   580                                  585                                  590  
 Ser Thr Ile Asp Leu Pro Asn Tyr Gly Cys Thr Ile Pro Glu Lys Thr  
                   595                                  600                                  605  
 Ser Cys Ser Val Tyr Gly Trp Gly Tyr Thr Gly Leu Ile Asn Tyr Asp  
                   610                                  615                                  620  
 Gly Leu Leu Arg Val Ala His Leu Tyr Ile Met Gly Asn Glu Lys Cys  
                   625                                  630                                  635                                  640  
 Ser Gln His His Arg Gly Lys Val Thr Leu Asn Glu Ser Glu Ile Cys  
                                   645                                  650                                  655  
 Ala Gly Ala Glu Lys Ile Gly Ser Gly Pro Cys Glu Gly Asp Tyr Gly  
                                   660                                  665                                  670  
 Gly Pro Leu Val Cys Glu Gln His Lys Met Arg Met Val Leu Gly Val  
                                   675                                  680                                  685  
 Ile Val Pro Gly Arg Gly Cys Ala Ile Pro Asn Arg Pro Gly Ile Phe  
                                   690                                  695                                  700  
 Val Arg Val Ala Tyr Tyr Ala Lys Trp Ile His Lys Ile Ile Leu Thr  
                                   705                                  710                                  715                                  720  
 Tyr Lys Val Pro Gln Ser  
                                   725

<210> 121  
 <211> 191  
 <212> PRT  
 <213> Homo sapiens

<400> 121  
 Phe Pro Thr Ile Pro Leu Ser Arg Leu Phe Asp Asn Ala Met Leu Arg  
 1                  5                                  10                                  15  
 Ala His Arg Leu His Gln Leu Ala Phe Asp Thr Tyr Gln Glu Phe Glu  
                   20                                  25                                  30  
 Glu Ala Tyr Ile Pro Lys Glu Gln Lys Tyr Ser Phe Leu Gln Asn Pro  
                   35                                  40                                  45  
 Gln Thr Ser Leu Cys Phe Ser Glu Ser Ile Pro Thr Pro Ser Asn Arg  
                   50                                  55                                  60  
 Glu Glu Thr Gln Gln Lys Ser Asn Leu Glu Leu Leu Arg Ile Ser Leu  
 65                                  70                                  75                                  80  
 Leu Leu Ile Gln Ser Trp Leu Glu Pro Val Gln Phe Leu Arg Ser Val  
                                   85                                  90                                  95  
 Phe Ala Asn Ser Leu Val Tyr Gly Ala Ser Asp Ser Asn Val Tyr Asp  
                   100                                  105                                  110  
  
 Leu Leu Lys Asp Leu Glu Glu Gly Ile Gln Thr Leu Met Gly Arg Leu  
                   115                                  120                                  125  
 Glu Asp Gly Ser Pro Arg Thr Gly Gln Ile Phe Lys Gln Thr Tyr Ser  
                   130                                  135                                  140  
 Lys Phe Asp Thr Asn Ser His Asn Asp Asp Ala Leu Leu Lys Asn Tyr  
 145                                  150                                  155                                  160  
 Gly Leu Leu Tyr Cys Phe Arg Lys Asp Met Asp Lys Val Glu Thr Phe  
                                   165                                  170                                  175  
 Leu Arg Ile Val Gln Cys Arg Ser Val Glu Gly Ser Cys Gly Phe  
                                   180                                  185                                  190

<210> 122  
 <211> 156  
 <212> PRT  
 <213> Homo sapiens

<400> 122  
 Ala Tyr Arg Pro Ser Glu Thr Leu Cys Gly Gly Glu Leu Val Asp Thr  
 1                  5                                  10                                  15  
 Leu Gln Phe Val Cys Gly Asp Arg Gly Phe Tyr Phe Ser Arg Pro Ala

```

      20      25      30
Ser Arg Val Ser Arg Arg Ser Arg Gly Ile Val Glu Glu Cys Cys Phe
      35      40      45
Arg Ser Cys Asp Leu Ala Leu Leu Glu Thr Tyr Cys Ala Thr Pro Ala
      50      55      60
Lys Ser Glu Arg Asp Val Ser Thr Pro Pro Thr Val Leu Pro Asp Asn
      65      70      75      80
Phe Pro Arg Tyr Pro Val Gly Lys Phe Phe Gln Tyr Asp Thr Trp Lys
      85      90      95
Gln Ser Thr Gln Arg Leu Arg Arg Gly Leu Pro Ala Leu Leu Arg Ala
      100      105      110
Arg Arg Gly His Val Leu Ala Lys Glu Leu Glu Ala Phe Arg Glu Ala
      115      120      125
Lys Arg His Arg Pro Leu Ile Ala Leu Pro Thr Gln Asp Pro Ala His
      130      135      140
Gly Gly Ala Pro Pro Glu Met Ala Ser Asn Arg Lys
      145      150      155

```

```

<210> 123
<211> 735
<212> PRT
<213> Homo sapiens

```

```

<400> 123
Glu Val Lys Gln Glu Asn Arg Leu Leu Asn Glu Ser Glu Ser Ser Ser
1      5      10      15
Gln Gly Leu Leu Gly Tyr Tyr Phe Ser Asp Leu Asn Phe Gln Ala Pro
      20      25      30
Met Val Val Thr Ser Ser Thr Thr Gly Asp Leu Ser Ile Pro Ser Ser
      35      40      45
Glu Leu Glu Asn Ile Pro Ser Glu Asn Gln Tyr Phe Gln Ser Ala Ile
      50      55      60
Trp Ser Gly Phe Ile Lys Val Lys Lys Ser Asp Glu Tyr Thr Phe Ala
      65      70      75      80
Thr Ser Ala Asp Asn His Val Thr Met Trp Val Asp Asp Gln Glu Val
      85      90      95
Ile Asn Lys Ala Ser Asn Ser Asn Lys Ile Arg Leu Glu Lys Gly Arg
      100      105      110
Leu Tyr Gln Ile Lys Ile Gln Tyr Gln Arg Glu Asn Pro Thr Glu Lys
      115      120      125
Gly Leu Asp Phe Lys Leu Tyr Trp Thr Asp Ser Gln Asn Lys Lys Glu
      130      135      140
Val Ile Ser Ser Asp Asn Leu Gln Leu Pro Glu Leu Lys Gln Lys Ser
      145      150      155      160
Ser Asn Ser Arg Lys Lys Arg Ser Thr Ser Ala Gly Pro Thr Val Pro
      165      170      175
Asp Arg Asp Asn Asp Gly Ile Pro Asp Ser Leu Glu Val Glu Gly Tyr
      180      185      190
Thr Val Asp Val Lys Asn Lys Arg Thr Phe Leu Ser Pro Trp Ile Ser
      195      200      205
Asn Ile His Glu Lys Lys Gly Leu Thr Lys Tyr Lys Ser Ser Pro Glu
      210      215      220
Lys Trp Ser Thr Ala Ser Asp Pro Tyr Ser Asp Phe Glu Lys Val Thr
      225      230      235      240
Gly Arg Ile Asp Lys Asn Val Ser Pro Glu Ala Arg His Pro Leu Val
      245      250      255
Ala Ala Tyr Pro Ile Val His Val Asp Met Glu Asn Ile Ile Leu Ser
      260      265      270
Lys Asn Glu Asp Gln Ser Thr Gln Asn Thr Asp Ser Gln Thr Arg Thr
      275      280      285
Ile Ser Lys Asn Thr Ser Thr Ser Arg Thr His Thr Ser Glu Val His
      290      295      300

```

Gly Asn Ala Glu Val His Ala Ser Phe Phe Asp Ile Gly Gly Ser Val  
 305 310 315 320  
 Ser Ala Gly Phe Ser Asn Ser Asn Ser Ser Thr Val Ala Ile Asp His  
 325 330 335  
 Ser Leu Ser Leu Ala Gly Glu Arg Thr Trp Ala Glu Thr Met Gly Leu  
 340 345 350  
 Asn Thr Ala Asp Thr Ala Arg Leu Asn Ala Asn Ile Arg Tyr Val Asn  
 355 360 365  
 Thr Gly Thr Ala Pro Ile Tyr Asn Val Leu Pro Thr Thr Ser Leu Val  
 370 375 380  
 Leu Gly Lys Asn Gln Thr Leu Ala Thr Ile Lys Ala Lys Glu Asn Gln  
 385 390 395 400  
 Leu Ser Gln Ile Leu Ala Pro Asn Asn Tyr Tyr Pro Ser Lys Asn Leu  
 405 410 415  
 Ala Pro Ile Ala Leu Asn Ala Gln Asp Asp Phe Ser Ser Thr Pro Ile  
 420 425 430  
 Thr Met Asn Tyr Asn Gln Phe Leu Glu Leu Glu Lys Thr Lys Gln Leu  
 435 440 445  
 Arg Leu Asp Thr Asp Gln Val Tyr Gly Asn Ile Ala Thr Tyr Asn Phe  
 450 455 460  
 Glu Asn Gly Arg Val Arg Val Asp Thr Gly Ser Asn Trp Ser Glu Val  
 465 470 475 480  
 Leu Pro Gln Ile Gln Glu Thr Thr Ala Arg Ile Ile Phe Asn Gly Lys  
 485 490 495  
 Asp Leu Asn Leu Val Glu Arg Arg Ile Ala Val Asn Pro Ser Asp  
 500 505 510  
 Pro Leu Glu Thr Thr Lys Pro Asp Met Thr Leu Lys Glu Ala Leu Lys  
 515 520 525  
 Ile Ala Phe Gly Phe Asn Glu Pro Asn Gly Asn Leu Gln Tyr Gln Gly  
 530 535 540  
 Lys Asp Ile Thr Glu Phe Asp Phe Asn Phe Asp Gln Gln Thr Ser Gln  
 545 550 555 560  
 Asn Ile Lys Asn Gln Leu Ala Glu Leu Asn Ala Thr Asn Ile Tyr Thr  
 565 570 575  
 Val Leu Asp Lys Ile Lys Leu Asn Ala Lys Met Asn Ile Leu Ile Arg  
 580 585 590  
 Asp Lys Arg Phe His Tyr Asp Arg Asn Asn Ile Ala Val Gly Ala Asp  
 595 600 605  
 Glu Ser Val Val Lys Glu Ala His Arg Glu Val Ile Asn Ser Ser Thr  
 610 615 620  
 Glu Gly Leu Leu Leu Asn Ile Asp Lys Asp Ile Arg Lys Ile Leu Ser  
 625 630 635 640  
 Gly Tyr Ile Val Glu Ile Glu Asp Thr Glu Gly Leu Lys Glu Val Ile  
 645 650 655  
 Asn Asp Arg Tyr Asp Met Leu Asn Ile Ser Ser Leu Arg Gln Asp Gly  
 660 665 670  
 Lys Thr Phe Ile Asp Phe Lys Lys Tyr Asn Asp Lys Leu Pro Leu Tyr  
 675 680 685  
 Ile Ser Asn Pro Asn Tyr Lys Val Asn Val Tyr Ala Val Thr Lys Glu  
 690 695 700  
 Asn Thr Ile Ile Asn Pro Ser Glu Asn Gly Asp Thr Ser Thr Asn Gly  
 705 710 715 720  
 Ile Lys Lys Ile Leu Ile Phe Ser Lys Lys Gly Tyr Glu Ile Gly  
 725 730 735

<210> 124  
 <211> 509  
 <212> PRT  
 <213> Homo sapiens

<400> 124  
 Met Lys Val Lys Gly Thr Arg Arg Asn Tyr Gln His Leu Trp Arg Trp

1				5						10					15
Gly	Thr	Leu	Leu	Leu	Gly	Met	Leu	Met	Ile	Cys	Ser	Ala	Thr	Glu	Lys
			20					25					30		
Leu	Trp	Val	Thr	Val	Tyr	Tyr	Gly	Val	Pro	Val	Trp	Lys	Glu	Ala	Thr
		35					40					45			
Thr	Thr	Leu	Phe	Cys	Ala	Ser	Asp	Ala	Arg	Ala	Tyr	Asp	Thr	Glu	Val
		50					55					60			
His	Asn	Val	Trp	Ala	Thr	His	Ala	Cys	Val	Pro	Thr	Asp	Pro	Asn	Pro
65					70					75					80
Gln	Glu	Val	Val	Leu	Gly	Asn	Val	Thr	Glu	Asn	Phe	Asn	Met	Trp	Lys
				85					90					95	
Asn	Asn	Met	Val	Glu	Gln	Met	Gln	Glu	Asp	Ile	Ile	Ser	Leu	Trp	Asp
			100					105					110		
Gln	Ser	Leu	Lys	Pro	Cys	Val	Lys	Leu	Thr	Pro	Leu	Cys	Val	Thr	Leu
		115					120					125			
Asn	Cys	Thr	Asp	Leu	Gly	Lys	Ala	Thr	Asn	Thr	Asn	Ser	Ser	Asn	Trp
		130				135						140			
Lys	Glu	Glu	Ile	Lys	Gly	Glu	Ile	Lys	Asn	Cys	Ser	Phe	Asn	Ile	Thr
145					150					155					160
Thr	Ser	Ile	Arg	Asp	Lys	Ile	Gln	Lys	Glu	Asn	Ala	Leu	Phe	Arg	Asn
				165					170					175	
Leu	Asp	Val	Val	Pro	Ile	Asp	Asn	Ala	Ser	Thr	Thr	Thr	Asn	Tyr	Thr
			180					185					190		
Asn	Tyr	Arg	Leu	Ile	His	Cys	Asn	Arg	Ser	Val	Ile	Thr	Gln	Ala	Cys
		195					200						205		
Pro	Lys	Val	Ser	Phe	Glu	Pro	Ile	Pro	Ile	His	Tyr	Cys	Thr	Pro	Ala
		210				215					220				
Gly	Phe	Ala	Ile	Leu	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Lys	Gly
225					230					235					240
Pro	Cys	Thr	Asn	Val	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro
				245						250				255	
Ile	Val	Ser	Thr	Gln	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Glu
			260					265					270		
Val	Val	Ile	Arg	Ser	Asp	Asn	Phe	Thr	Asn	Asn	Ala	Lys	Thr	Ile	Ile
		275					280					285			
Val	Gln	Leu	Asn	Glu	Ser	Val	Ala	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn
		290				295					300				
Asn	Thr	Arg	Lys	Ser	Ile	Tyr	Ile	Gly	Pro	Gly	Arg	Ala	Phe	His	Thr
305					310					315					320
Thr	Gly	Arg	Ile	Ile	Gly	Asp	Ile	Arg	Lys	Ala	His	Cys	Asn	Ile	Ser
				325					330					335	
Arg	Ala	Gln	Trp	Asn	Asn	Thr	Leu	Glu	Gln	Ile	Val	Lys	Lys	Leu	Arg
			340					345					350		
Glu	Gln	Phe	Gly	Asn	Asn	Lys	Thr	Ile	Val	Phe	Asn	Gln	Ser	Ser	Gly
		355					360				365				
Gly	Asp	Pro	Glu	Ile	Val	Met	His	Ser	Phe	Asn	Cys	Arg	Gly	Glu	Phe
		370				375				380					
Phe	Tyr	Cys	Asn	Thr	Thr	Gln	Leu	Phe	Asn	Asn	Thr	Trp	Arg	Leu	Asn
385					390					395					400
His	Thr	Glu	Gly	Thr	Lys	Gly	Asn	Asp	Thr	Ile	Ile	Leu	Pro	Cys	Arg
				405					410					415	
Ile	Lys	Gln	Ile	Ile	Asn	Met	Trp	Gln	Glu	Val	Gly	Lys	Ala	Met	Tyr
			420					425					430		
Ala	Pro	Pro	Ile	Gly	Gly	Gln	Ile	Ser	Cys	Ser	Ser	Asn	Ile	Thr	Gly
		435					440					445			
Leu	Leu	Leu	Thr	Arg	Asp	Gly	Gly	Thr	Asn	Val	Thr	Asn	Asp	Thr	Glu
		450				455					460				
Val	Phe	Arg	Pro	Gly	Gly	Gly	Asp	Met	Arg	Asp	Asn	Trp	Arg	Ser	Glu
465					470					475					480
Leu	Tyr	Lys	Tyr	Lys	Val	Ile	Lys	Ile	Glu	Pro	Leu	Gly	Ile	Ala	Pro
				485					490					495	
Thr	Lys	Ala	Lys	Arg	Arg	Val	Val	Gln	Arg	Glu	Lys	Arg			
			500					505							

<210> 125  
 <211> 101  
 <212> PRT  
 <213> Homo sapiens

<400> 125  
 Ser Trp Val Ile Pro Pro Ile Ser Cys Pro Glu Asn Glu Lys Gly Pro  
 1 5 10 15  
 Phe Pro Lys Asn Leu Val Gln Ile Lys Ser Asn Lys Asp Lys Glu Gly  
 20 25 30  
 Lys Val Phe Tyr Ser Ile Thr Gly Gln Gly Ala Asp Thr Pro Pro Val  
 35 40 45  
 Gly Val Phe Ile Ile Glu Arg Glu Thr Gly Trp Leu Lys Val Thr Glu  
 50 55 60  
 Pro Leu Asp Arg Glu Arg Ile Ala Thr Tyr Thr Leu Phe Ser His Ala  
 65 70 75 80  
 Val Ser Ser Asn Gly Asn Ala Val Glu Asp Pro Met Glu Ile Leu Ile  
 85 90 95  
 Thr Val Thr Asp Gln  
 100

<210> 126  
 <211> 459  
 <212> PRT  
 <213> Homo sapiens

<400> 126  
 Glu Ile Cys Gly Pro Gly Ile Asp Ile Arg Asn Asp Tyr Gln Gln Leu  
 1 5 10 15  
 Lys Arg Leu Glu Asn Cys Thr Val Ile Glu Gly Tyr Leu His Ile Leu  
 20 25 30  
 Leu Ile Ser Lys Ala Glu Asp Tyr Arg Ser Tyr Arg Phe Pro Lys Leu  
 35 40 45  
 Thr Val Ile Thr Glu Tyr Leu Leu Leu Phe Arg Val Ala Gly Leu Glu  
 50 55 60  
 Ser Leu Gly Asp Leu Phe Pro Asn Leu Thr Val Ile Arg Gly Trp Lys  
 65 70 75 80  
 Leu Phe Tyr Asn Tyr Ala Leu Val Ile Phe Glu Met Thr Asn Leu Lys  
 85 90 95  
 Asp Ile Gly Leu Tyr Asn Leu Arg Asn Ile Thr Arg Gly Ala Ile Arg  
 100 105 110  
 Ile Glu Lys Asn Ala Asp Leu Cys Tyr Leu Ser Thr Val Asp Trp Ser  
 115 120 125  
 Leu Ile Leu Asp Ala Val Ser Asn Asn Tyr Ile Val Gly Asn Lys Pro  
 130 135 140  
 Pro Lys Glu Cys Gly Asp Leu Cys Pro Gly Thr Met Glu Glu Lys Pro  
 145 150 155 160  
 Met Cys Glu Lys Thr Thr Ile Asn Asn Glu Tyr Asn Tyr Arg Cys Trp  
 165 170 175  
 Thr Thr Asn Arg Cys Gln Lys Met Cys Pro Ser Thr Cys Gly Lys Arg  
 180 185 190  
 Ala Cys Thr Glu Asn Asn Glu Cys Cys His Pro Glu Cys Leu Gly Ser  
 195 200 205  
 Cys Ser Ala Pro Asp Asn Asp Thr Ala Cys Val Ala Cys Arg His Tyr  
 210 215 220  
 Tyr Tyr Ala Gly Val Cys Val Pro Ala Cys Pro Pro Asn Thr Tyr Arg  
 225 230 235 240  
 Phe Glu Gly Trp Arg Cys Val Asp Arg Asp Phe Cys Ala Asn Ile Leu  
 245 250 255  
 Ser Ala Glu Ser Ser Asp Ser Glu Gly Phe Val Ile His Asp Gly Glu

```

      260      265      270
Cys Met Gln Glu Cys Pro Ser Gly Phe Ile Arg Asn Gly Ser Gln Ser
      275      280      285
Met Tyr Cys Ile Pro Cys Glu Gly Pro Cys Pro Lys Val Cys Glu Glu
      290      295      300
Glu Lys Lys Thr Lys Thr Ile Asp Ser Val Thr Ser Ala Gln Met Leu
305      310      315
Gln Gly Cys Thr Ile Phe Lys Gly Asn Leu Leu Ile Asn Ile Arg Arg
      325      330      335
Gly Asn Asn Ile Ala Ser Glu Leu Glu Asn Phe Met Gly Leu Ile Glu
      340      345      350
Val Val Thr Gly Tyr Val Lys Ile Arg His Ser His Ala Leu Val Ser
      355      360      365
Leu Ser Phe Leu Lys Asn Leu Arg Leu Ile Leu Gly Glu Glu Gln Leu
      370      375      380
Glu Gly Asn Tyr Ser Phe Tyr Val Leu Asp Asn Gln Asn Leu Gln Gln
385      390      395
Leu Trp Asp Trp Asp His Arg Asn Leu Thr Ile Lys Ala Gly Lys Met
      405      410      415
Tyr Phe Ala Phe Asn Pro Lys Leu Cys Val Ser Glu Ile Tyr Arg Met
      420      425      430
Glu Glu Val Thr Gly Thr Lys Gly Arg Gln Ser Lys Gly Asp Ile Asn
      435      440      445
Thr Arg Asn Asn Gly Glu Arg Ala Ser Cys Glu
      450      455

```

```

<210> 127
<211> 146
<212> PRT
<213> Homo sapiens

```

```

<400> 127
Val Pro Ile Gln Lys Val Gln Asp Asp Thr Lys Thr Leu Ile Lys Thr
1      5      10      15
Ile Val Thr Arg Ile Asn Asp Ile Ser His Thr Gln Ser Val Ser Ser
      20      25      30
Lys Gln Lys Val Thr Gly Leu Asp Phe Ile Pro Gly Leu His Pro Ile
      35      40      45
Leu Thr Leu Ser Lys Met Asp Gln Thr Leu Ala Val Tyr Gln Gln Ile
      50      55      60
Leu Thr Ser Met Pro Ser Arg Asn Val Ile Gln Ile Ser Asn Asp Leu
65      70      75      80
Glu Asn Leu Arg Asp Leu Leu His Val Leu Ala Phe Ser Lys Ser Cys
      85      90      95
His Leu Pro Trp Ala Ser Gly Leu Glu Thr Leu Asp Ser Leu Gly Gly
      100      105      110
Val Leu Glu Ala Ser Gly Tyr Ser Thr Glu Val Val Ala Leu Ser Arg
      115      120      125
Leu Gln Gly Ser Leu Gln Asp Met Leu Trp Gln Leu Asp Leu Ser Pro
      130      135      140
Gly Cys
145

```

```

<210> 128
<211> 327
<212> PRT
<213> Homo sapiens

```

```

<400> 128
Lys Glu Ile Thr Asn Ala Leu Glu Thr Trp Gly Ala Leu Gly Gln Asp
1      5      10      15

```



```

Ile Asn Leu Asp Ile Pro Ser Phe Gln Met Ser Asp Asp Ile Asp Asp
      20      25      30
Ile Lys Trp Glu Lys Thr Ser Asp Lys Lys Lys Ile Ala Gln Phe Arg
      35      40      45
Lys Glu Lys Glu Thr Phe Lys Glu Lys Asp Thr Tyr Lys Leu Phe Lys
      50      55      60
Asn Gly Thr Leu Lys Ile Lys His Leu Lys Thr Asp Asp Gln Asp Ile
      65      70      75      80
Tyr Lys Val Ser Ile Tyr Asp Thr Lys Gly Lys Asn Val Leu Glu Lys
      85      90      95
Ile Phe Asp Leu Lys Ile Gln Glu Arg Val Ser Lys Pro Lys Ile Ser
      100      105      110
Trp Thr Cys Ile Asn Thr Thr Leu Thr Cys Glu Val Met Asn Gly Thr
      115      120      125
Asp Pro Glu Leu Asn Leu Tyr Gln Asp Gly Lys His Leu Lys Leu Ser
      130      135      140
Gln Arg Val Ile Thr His Lys Trp Thr Thr Ser Leu Ser Ala Lys Phe
      145      150      155      160
Lys Cys Thr Ala Gly Asn Lys Val Ser Lys Glu Ser Ser Val Glu Pro
      165      170      175
Val Ser Cys Pro Glu Lys Gly Leu Asp Ile Tyr Leu Ile Ile Gly Ile
      180      185      190
Cys Gly Gly Gly Ser Leu Leu Met Val Phe Val Ala Leu Leu Val Phe
      195      200      205
Tyr Ile Thr Lys Arg Lys Lys Gln Arg Ser Arg Arg Asn Asp Glu Glu
      210      215      220
Leu Glu Thr Arg Ala His Arg Val Ala Thr Glu Glu Arg Gly Arg Lys
      225      230      235      240
Pro Gln Gln Ile Pro Ala Ser Thr Pro Gln Asn Pro Ala Thr Ser Gln
      245      250      255
His Pro Pro Pro Pro Pro Gly His Arg Ser Gln Ala Pro Ser His Arg
      260      265      270
Pro Pro Pro Pro Gly His Arg Val Gln His Gln Pro Gln Lys Arg Pro
      275      280      285
Pro Ala Pro Ser Gly Thr Gln Val His Gln Gln Lys Gly Pro Pro Leu
      290      295      300
Pro Arg Pro Arg Val Gln Pro Lys Pro Pro His Gly Ala Ala Glu Asn
      305      310      315      320
Ser Leu Ser Pro Ser Ser Asn
      325

```

```

<210> 129
<211> 433
<212> PRT
<213> Homo sapiens

```

```

<400> 129
Lys Lys Val Val Leu Gly Lys Lys Gly Asp Thr Val Glu Leu Thr Cys
1      5      10      15
Thr Ala Ser Gln Lys Lys Ser Ile Gln Phe His Trp Lys Asn Ser Asn
      20      25      30
Gln Ile Lys Ile Leu Gly Asn Gln Gly Ser Phe Leu Thr Lys Gly Pro
      35      40      45
Ser Lys Leu Asn Asp Arg Ala Asp Ser Arg Arg Ser Leu Trp Asp Gln
      50      55      60
Gly Asn Phe Pro Leu Ile Ile Lys Asn Leu Lys Ile Glu Asp Ser Asp
      65      70      75      80
Thr Tyr Ile Cys Glu Val Glu Asp Gln Lys Glu Glu Val Gln Leu Leu
      85      90      95
Val Phe Gly Leu Thr Ala Asn Ser Asp Thr His Leu Leu Gln Gly Gln
      100      105      110
Ser Leu Thr Leu Thr Leu Glu Ser Pro Pro Gly Ser Ser Pro Ser Val

```

```

      115      120      125
Gln Cys Arg Ser Pro Arg Gly Lys Asn Ile Gln Gly Gly Lys Thr Leu
  130      135      140
Ser Val Ser Gln Leu Glu Leu Gln Asp Ser Gly Thr Trp Thr Cys Thr
  145      150      155      160
Val Leu Gln Asn Gln Lys Lys Val Glu Phe Lys Ile Asp Ile Val Val
      165      170      175
Leu Ala Phe Gln Lys Ala Ser Ser Ile Val Tyr Lys Lys Glu Gly Glu
      180      185      190
Gln Val Glu Phe Ser Phe Pro Leu Ala Phe Thr Val Glu Lys Leu Thr
      195      200      205
Gly Ser Gly Glu Leu Trp Trp Gln Ala Glu Arg Ala Ser Ser Ser Lys
      210      215      220
Ser Trp Ile Thr Phe Asp Leu Lys Asn Lys Glu Val Ser Val Lys Arg
  225      230      235      240
Val Thr Gln Asp Pro Lys Leu Gln Met Gly Lys Lys Leu Pro Leu His
      245      250      255
Leu Thr Leu Pro Gln Ala Leu Pro Gln Tyr Ala Gly Ser Gly Asn Leu
      260      265      270
Thr Leu Ala Leu Glu Ala Lys Thr Gly Lys Leu His Gln Glu Val Asn
      275      280      285
Leu Val Val Met Arg Ala Thr Gln Leu Gln Lys Asn Leu Thr Cys Glu
      290      295      300
Val Trp Gly Pro Thr Ser Pro Lys Leu Met Leu Ser Leu Lys Leu Glu
  305      310      315      320
Asn Lys Glu Ala Lys Val Ser Lys Arg Glu Lys Ala Val Trp Val Leu
      325      330      335
Asn Pro Glu Ala Gly Met Trp Gln Cys Leu Leu Ser Asp Ser Gly Gln
      340      345      350
Val Leu Leu Glu Ser Asn Ile Lys Val Leu Pro Thr Trp Ser Thr Pro
      355      360      365
Val Gln Pro Met Ala Leu Ile Val Leu Gly Gly Val Ala Gly Leu Leu
      370      375      380
Leu Phe Ile Gly Leu Gly Ile Phe Phe Cys Val Arg Cys Arg His Arg
  385      390      395      400
Arg Arg Gln Ala Glu Arg Met Ser Gln Ile Lys Arg Leu Leu Ser Glu
      405      410      415
Lys Lys Thr Cys Gln Cys Pro His Arg Phe Gln Lys Thr Cys Ser Pro
      420      425      430
Ile

```

```

<210> 130
<211> 1145
<212> PRT
<213> Homo sapiens

```

```

<400> 130
Tyr Asn Leu Asp Val Arg Gly Ala Arg Ser Phe Ser Pro Pro Arg Ala
  1      5      10      15
Gly Arg His Phe Gly Tyr Arg Val Leu Gln Val Gly Asn Gly Val Ile
      20      25      30
Val Gly Ala Pro Gly Glu Gly Asn Ser Thr Gly Ser Leu Tyr Gln Cys
      35      40      45
Gln Ser Gly Thr Gly His Cys Leu Pro Val Thr Leu Arg Gly Ser Asn
      50      55      60
Tyr Thr Ser Lys Tyr Leu Gly Met Thr Leu Ala Thr Asp Pro Thr Asp
  65      70      75      80
Gly Ser Ile Leu Ala Cys Asp Pro Gly Leu Ser Arg Thr Cys Asp Gln
      85      90      95
Asn Thr Tyr Leu Ser Gly Leu Cys Tyr Leu Phe Arg Gln Asn Leu Gln
      100      105      110
Gly Pro Met Leu Gln Gly Arg Pro Gly Phe Gln Glu Cys Ile Lys Gly

```



Ile Thr Ile Cys Phe Gln Ile Lys Ser Leu Tyr Pro Gln Phe Gln Gly  
 625 630 635 640  
 Arg Leu Val Ala Asn Leu Thr Tyr Thr Leu Gln Leu Asp Gly His Arg  
 645 650 655  
 Thr Arg Arg Arg Gly Leu Phe Pro Gly Gly Arg His Glu Leu Arg Arg  
 660 665 670  
 Asn Ile Ala Val Thr Thr Ser Met Ser Cys Thr Asp Phe Ser Phe His  
 675 680 685  
 Phe Pro Val Cys Val Gln Asp Leu Ile Ser Pro Ile Asn Val Ser Leu  
 690 695 700  
 Asn Phe Ser Leu Trp Glu Glu Glu Gly Thr Pro Arg Asp Gln Arg Ala  
 705 710 715 720  
 Gln Gly Lys Asp Ile Pro Pro Ile Leu Arg Pro Ser Leu His Ser Glu  
 725 730 735  
 Thr Trp Glu Ile Pro Phe Glu Lys Asn Cys Gly Glu Asp Lys Lys Cys  
 740 745 750  
 Glu Ala Asn Leu Arg Val Ser Phe Ser Pro Ala Arg Ser Arg Ala Leu  
 755 760 765  
 Arg Leu Thr Ala Phe Ala Ser Leu Ser Val Glu Leu Ser Leu Ser Asn  
 770 775 780  
 Leu Glu Glu Asp Ala Tyr Trp Val Gln Leu Asp Leu His Phe Pro Pro  
 785 790 795 800  
 Gly Leu Ser Phe Arg Lys Val Glu Met Leu Lys Pro His Ser Gln Ile  
 805 810 815  
 Pro Val Ser Cys Glu Glu Leu Pro Glu Glu Ser Arg Leu Leu Ser Arg  
 820 825 830  
 Ala Leu Ser Cys Asn Val Ser Ser Pro Ile Phe Lys Ala Gly His Ser  
 835 840 845  
 Val Ala Leu Gln Met Met Phe Asn Thr Leu Val Asn Ser Ser Trp Gly  
 850 855 860  
 Asp Ser Val Glu Leu His Ala Asn Val Thr Cys Asn Asn Glu Asp Ser  
 865 870 875 880  
 Asp Leu Leu Glu Asp Asn Ser Ala Thr Thr Ile Ile Pro Ile Leu Tyr  
 885 890 895  
 Pro Ile Asn Ile Leu Ile Gln Asp Gln Glu Asp Ser Thr Leu Tyr Val  
 900 905 910  
 Ser Phe Thr Pro Lys Gly Pro Lys Ile His Gln Val Lys His Met Tyr  
 915 920 925  
 Gln Val Arg Ile Gln Pro Ser Ile His Asp His Asn Ile Pro Thr Leu  
 930 935 940  
 Glu Ala Val Val Gly Val Pro Gln Pro Pro Ser Glu Gly Pro Ile Thr  
 945 950 955 960  
 His Gln Trp Ser Val Gln Met Glu Pro Pro Val Pro Cys His Tyr Glu  
 965 970 975  
 Asp Leu Glu Arg Leu Pro Asp Ala Ala Glu Pro Cys Leu Pro Gly Ala  
 980 985 990  
 Leu Phe Arg Cys Pro Val Val Phe Arg Gln Glu Ile Leu Val Gln Val  
 995 1000 1005  
 Ile Gly Thr Leu Glu Leu Val Gly Glu Ile Glu Ala Ser Ser Met  
 1010 1015 1020  
 Phe Ser Leu Cys Ser Ser Leu Ser Ile Ser Phe Asn Ser Ser Lys  
 1025 1030 1035  
 His Phe His Leu Tyr Gly Ser Asn Ala Ser Leu Ala Gln Val Val  
 1040 1045 1050  
 Met Lys Val Asp Val Val Tyr Glu Lys Gln Met Leu Tyr Leu Tyr  
 1055 1060 1065  
 Val Leu Ser Gly Ile Gly Gly Leu Leu Leu Leu Leu Leu Ile Phe  
 1070 1075 1080  
 Ile Val Leu Tyr Lys Val Gly Phe Phe Lys Arg Asn Leu Lys Glu  
 1085 1090 1095  
 Lys Met Glu Ala Gly Arg Gly Val Pro Asn Gly Ile Pro Ala Glu  
 1100 1105 1110  
 Asp Ser Glu Gln Leu Ala Ser Gly Gln Glu Ala Gly Asp Pro Gly

1115	1120	1125
Cys Leu Lys Pro Leu His	Glu Lys Asp Ser Glu Ser	Gly Gly Gly
1130	1135	1140
Lys Asp		
1145		

<210> 131  
 <211> 660  
 <212> PRT  
 <213> Homo sapiens

<400> 131

Met Glu Ala Leu Met Ala Arg Gly Ala Leu Thr Gly Pro Leu Arg Ala	
1 5 10 15	
Leu Cys Leu Leu Gly Cys Leu Leu Ser His Ala Ala Ala Ala Pro Ser	
20 25 30	
Pro Ile Ile Lys Phe Pro Gly Asp Val Ala Pro Lys Thr Asp Lys Glu	
35 40 45	
Leu Ala Val Gln Tyr Leu Asn Thr Phe Tyr Gly Cys Pro Lys Glu Ser	
50 55 60	
Cys Asn Leu Phe Val Leu Lys Asp Thr Leu Lys Lys Met Gln Lys Phe	
65 70 75 80	
Phe Gly Leu Pro Gln Thr Gly Asp Leu Asp Gln Asn Thr Ile Glu Thr	
85 90 95	
Met Arg Lys Pro Arg Cys Gly Asn Pro Asp Val Ala Asn Tyr Asn Phe	
100 105 110	
Phe Pro Arg Lys Pro Lys Trp Asp Lys Asn Gln Ile Thr Tyr Arg Ile	
115 120 125	
Ile Gly Tyr Thr Pro Asp Leu Asp Pro Glu Thr Val Asp Asp Ala Phe	
130 135 140	
Ala Arg Ala Phe Gln Val Trp Ser Asp Val Thr Pro Leu Arg Phe Ser	
145 150 155 160	
Arg Ile His Asp Gly Glu Ala Asp Ile Met Ile Asn Phe Gly Arg Trp	
165 170 175	
Glu His Gly Asp Gly Tyr Pro Phe Asp Gly Lys Asp Gly Leu Leu Ala	
180 185 190	
His Ala Phe Ala Pro Gly Thr Gly Val Gly Gly Asp Ser His Phe Asp	
195 200 205	
Asp Asp Glu Leu Trp Thr Leu Gly Glu Gly Gln Val Val Arg Val Lys	
210 215 220	
Tyr Gly Asn Ala Asp Gly Glu Tyr Cys Lys Phe Pro Phe Leu Phe Asn	
225 230 235 240	
Gly Lys Glu Tyr Asn Ser Cys Thr Asp Thr Gly Arg Ser Asp Gly Phe	
245 250 255	
Leu Trp Cys Ser Thr Thr Tyr Asn Phe Glu Lys Asp Gly Lys Tyr Gly	
260 265 270	
Phe Cys Pro His Glu Ala Leu Phe Thr Met Gly Gly Asn Ala Glu Gly	
275 280 285	
Gln Pro Cys Lys Phe Pro Phe Arg Phe Gln Gly Thr Ser Tyr Asp Ser	
290 295 300	
Cys Thr Thr Glu Gly Arg Thr Asp Gly Tyr Arg Trp Cys Gly Thr Thr	
305 310 315 320	
Glu Asp Tyr Asp Arg Asp Lys Lys Tyr Gly Phe Cys Pro Glu Thr Ala	
325 330 335	
Met Ser Thr Val Gly Gly Asn Ser Glu Gly Ala Pro Cys Val Phe Pro	
340 345 350	
Phe Thr Phe Leu Gly Asn Lys Tyr Glu Ser Cys Thr Ser Ala Gly Arg	
355 360 365	
Ser Asp Gly Lys Met Trp Cys Ala Thr Thr Ala Asn Tyr Asp Asp Asp	
370 375 380	
Arg Lys Trp Gly Phe Cys Pro Asp Gln Gly Tyr Ser Leu Phe Leu Val	
385 390 395 400	

Ala Ala His Glu Phe Gly His Ala Met Gly Leu Glu His Ser Gln Asp  
 405 410 415  
 Pro Gly Ala Leu Met Ala Pro Ile Tyr Thr Tyr Thr Lys Asn Phe Arg  
 420 425 430  
 Leu Ser Gln Asp Asp Ile Lys Gly Ile Gln Glu Leu Tyr Gly Ala Ser  
 435 440 445  
 Pro Asp Ile Asp Leu Gly Thr Gly Pro Thr Pro Thr Leu Gly Pro Val  
 450 455 460  
 Thr Pro Glu Ile Cys Lys Gln Asp Ile Val Phe Asp Gly Ile Ala Gln  
 465 470 475 480  
 Ile Arg Gly Glu Ile Phe Phe Phe Lys Asp Arg Phe Ile Trp Arg Thr  
 485 490 495  
 Val Thr Pro Arg Asp Lys Pro Met Gly Pro Leu Leu Val Ala Thr Phe  
 500 505 510  
 Trp Pro Glu Leu Pro Glu Lys Ile Asp Ala Val Tyr Glu Ala Pro Gln  
 515 520 525  
 Glu Glu Lys Ala Val Phe Phe Ala Gly Asn Glu Tyr Trp Ile Tyr Ser  
 530 535 540  
 Ala Ser Thr Leu Glu Arg Gly Tyr Pro Lys Pro Leu Thr Ser Leu Gly  
 545 550 555 560  
 Leu Pro Pro Asp Val Gln Arg Val Asp Ala Ala Phe Asn Trp Ser Lys  
 565 570 575  
 Asn Lys Lys Thr Tyr Ile Phe Ala Gly Asp Lys Phe Trp Arg Tyr Asn  
 580 585 590  
 Glu Val Lys Lys Lys Met Asp Pro Gly Phe Pro Lys Leu Ile Ala Asp  
 595 600 605  
 Ala Trp Asn Ala Ile Pro Asp Asn Leu Asp Ala Val Val Asp Leu Gln  
 610 615 620  
 Gly Gly Gly His Ser Tyr Phe Phe Lys Gly Ala Tyr Tyr Leu Lys Leu  
 625 630 635 640  
 Glu Asn Gln Ser Leu Lys Ser Val Lys Phe Gly Ser Ile Lys Ser Asp  
 645 650 655  
 Trp Leu Gly Cys  
 660

<210> 132  
 <211> 707  
 <212> PRT  
 <213> Homo sapiens

<400> 132  
 Met Ser Leu Trp Gln Pro Leu Val Leu Val Leu Leu Val Leu Gly Cys  
 1 5 10 15  
 Cys Phe Ala Ala Pro Arg Gln Arg Gln Ser Thr Leu Val Leu Phe Pro  
 20 25 30  
 Gly Asp Leu Arg Thr Asn Leu Thr Asp Arg Gln Leu Ala Glu Glu Tyr  
 35 40 45  
 Leu Tyr Arg Tyr Gly Tyr Thr Arg Val Ala Glu Met Arg Gly Glu Ser  
 50 55 60  
 Lys Ser Leu Gly Pro Ala Leu Leu Leu Leu Gln Lys Gln Leu Ser Leu  
 65 70 75 80  
 Pro Glu Thr Gly Glu Leu Asp Ser Ala Thr Leu Lys Ala Met Arg Thr  
 85 90 95  
 Pro Arg Cys Gly Val Pro Asp Leu Gly Arg Phe Gln Thr Phe Glu Gly  
 100 105 110  
 Asp Leu Lys Trp His His His Asn Ile Thr Tyr Trp Ile Gln Asn Tyr  
 115 120 125  
 Ser Glu Asp Leu Pro Arg Ala Val Ile Asp Asp Ala Phe Ala Arg Ala  
 130 135 140  
 Phe Ala Leu Trp Ser Ala Val Thr Pro Leu Thr Phe Thr Arg Val Tyr  
 145 150 155 160  
 Ser Arg Asp Ala Asp Ile Val Ile Gln Phe Gly Val Ala Glu His Gly



Phe Cys Gln Asp Arg Phe Tyr Trp Arg Val Ser Ser Arg Ser Glu Leu  
                   675                                  680                  685  
 Asn Gln Val Asp Gln Val Gly Tyr Val Thr Tyr Asp Ile Leu Gln Cys  
                   690                                  695                  700  
 Pro Glu Asp  
 705

<210> 133  
 <211> 115  
 <212> PRT  
 <213> Homo sapiens

<400> 133  
 Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu Thr Leu Ala  
 1                  5                                  10                  15  
 Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu Thr Leu Arg  
                   20                                  25                  30  
 Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr Glu Glu Ile  
                   35                                  40                  45  
 Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln Gly Gly Thr  
                   50                                  55                  60  
 Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys Tyr Ile Asp  
 65                  70                                  75                  80  
 Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val Asn Gln Phe  
                   85                                  90                  95  
 Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr Glu Trp Ile  
                   100                                  105                  110  
 Ile Glu Ser  
                   115

<210> 134  
 <211> 185  
 <212> PRT  
 <213> Homo sapiens

<400> 134  
 Ala Pro Val Pro Pro Gly Glu Asp Ser Lys Asp Val Ala Ala Pro His  
 1                  5                                  10                  15  
 Arg Gln Pro Leu Thr Ser Ser Glu Arg Ile Asp Lys Gln Ile Arg Tyr  
                   20                                  25                  30  
 Ile Leu Asp Gly Ile Ser Ala Leu Arg Lys Glu Thr Cys Asn Lys Ser  
                   35                                  40                  45  
 Asn Met Cys Glu Ser Ser Lys Glu Ala Leu Ala Glu Asn Asn Leu Asn  
                   50                                  55                  60  
 Leu Pro Lys Met Ala Glu Lys Asp Gly Cys Phe Gln Ser Gly Phe Asn  
 65                  70                                  75                  80  
 Glu Glu Thr Cys Leu Val Lys Ile Ile Thr Gly Leu Leu Glu Phe Glu  
                   85                                  90                  95  
 Val Tyr Leu Glu Tyr Leu Gln Asn Arg Phe Glu Ser Ser Glu Glu Gln  
                   100                                  105                  110  
 Ala Arg Ala Val Gln Met Ser Thr Lys Val Leu Ile Gln Phe Leu Gln  
                   115                                  120                  125  
 Lys Lys Ala Lys Asn Leu Asp Ala Ile Thr Thr Pro Asp Pro Thr Thr  
                   130                                  135                  140  
 Asn Ala Ser Leu Leu Thr Lys Leu Gln Ala Gln Asn Gln Trp Leu Gln  
 145                  150                                  155                  160  
 Asp Met Thr Thr His Leu Ile Leu Arg Ser Phe Lys Glu Phe Leu Gln  
                   165                                  170                  175  
 Ser Ser Leu Arg Ala Leu Arg Gln Met  
                   180                                  185



<210> 135  
 <211> 160  
 <212> PRT  
 <213> Homo sapiens

<400> 135  
 Ser Pro Gly Gln Gly Thr Gln Ser Glu Asn Ser Cys Thr His Phe Pro  
 1 5 10 15  
 Gly Asn Leu Pro Asn Met Leu Arg Asp Leu Arg Asp Ala Phe Ser Arg  
 20 25 30  
 Val Lys Thr Phe Phe Gln Met Lys Asp Gln Leu Asp Asn Leu Leu Leu  
 35 40 45  
 Lys Glu Ser Leu Leu Glu Asp Phe Lys Gly Tyr Leu Gly Cys Gln Ala  
 50 55 60  
 Leu Ser Glu Met Ile Gln Phe Tyr Leu Glu Glu Val Met Pro Gln Ala  
 65 70 75 80  
 Glu Asn Gln Asp Pro Asp Ile Lys Ala His Val Asn Ser Leu Gly Glu  
 85 90 95  
 Asn Leu Lys Thr Leu Arg Leu Arg Leu Arg Arg Cys His Arg Phe Leu  
 100 105 110  
 Pro Cys Glu Asn Lys Ser Lys Ala Val Glu Gln Val Lys Asn Ala Phe  
 115 120 125  
 Asn Lys Leu Gln Glu Lys Gly Ile Tyr Lys Ala Met Ser Glu Phe Asp  
 130 135 140  
 Ile Phe Ile Asn Tyr Ile Glu Ala Tyr Met Thr Met Lys Ile Arg Asn  
 145 150 155 160

<210> 136  
 <211> 472  
 <212> PRT  
 <213> Homo sapiens

<400> 136  
 Glu Met Gly Thr Ala Asp Leu Gly Pro Ser Ser Val Pro Thr Pro Thr  
 1 5 10 15  
 Asn Val Thr Ile Glu Ser Tyr Asn Met Asn Pro Ile Val Tyr Trp Glu  
 20 25 30  
 Tyr Gln Ile Met Pro Gln Val Pro Val Phe Thr Val Glu Val Lys Asn  
 35 40 45  
 Tyr Gly Val Lys Asn Ser Glu Trp Ile Asp Ala Cys Ile Asn Ile Ser  
 50 55 60  
 His His Tyr Cys Asn Ile Ser Asp His Val Gly Asp Pro Ser Asn Ser  
 65 70 75 80  
 Leu Trp Val Arg Val Lys Ala Arg Val Gly Gln Lys Glu Ser Ala Tyr  
 85 90 95  
 Ala Lys Ser Glu Glu Phe Ala Val Cys Arg Asp Gly Lys Ile Gly Pro  
 100 105 110  
 Pro Lys Leu Asp Ile Arg Lys Glu Glu Lys Gln Ile Met Ile Asp Ile  
 115 120 125  
 Phe His Pro Ser Val Phe Val Asn Gly Asp Glu Gln Glu Val Asp Tyr  
 130 135 140  
 Asp Pro Glu Thr Thr Cys Tyr Ile Arg Val Tyr Asn Val Tyr Val Arg  
 145 150 155 160  
 Met Asn Gly Ser Glu Ile Gln Tyr Lys Ile Leu Thr Gln Lys Glu Asp  
 165 170 175  
 Asp Cys Asp Glu Ile Gln Cys Gln Leu Ala Ile Pro Val Ser Ser Leu  
 180 185 190  
 Asn Ser Gln Tyr Cys Val Ser Ala Glu Gly Val Leu His Val Trp Gly  
 195 200 205  
 Val Thr Thr Glu Lys Ser Lys Glu Val Cys Ile Thr Ile Phe Asn Ser  
 210 215 220  
 Ser Ile Lys Gly Ser Leu Trp Ile Pro Val Val Ala Ala Leu Leu Leu

```

225          230          235          240
Phe Leu Val Leu Ser Leu Val Phe Ile Cys Phe Tyr Ile Lys Lys Ile
          245          250          255
Asn Pro Leu Lys Glu Lys Ser Ile Ile Leu Pro Lys Ser Leu Ile Ser
          260          265          270
Val Val Arg Ser Ala Thr Leu Glu Thr Lys Pro Glu Ser Lys Tyr Val
          275          280          285
Ser Leu Ile Thr Ser Tyr Gln Pro Phe Ser Leu Glu Lys Glu Val Val
          290          295          300
Cys Glu Glu Pro Leu Ser Pro Ala Thr Val Pro Gly Met His Thr Glu
305          310          315          320
Asp Asn Pro Gly Lys Val Glu His Thr Glu Glu Leu Ser Ser Ile Thr
          325          330          335
Glu Val Val Thr Thr Glu Glu Asn Ile Pro Asp Val Val Pro Gly Ser
          340          345          350
His Leu Thr Pro Ile Glu Arg Glu Ser Ser Ser Pro Leu Ser Ser Asn
          355          360          365
Gln Ser Glu Pro Gly Ser Ile Ala Leu Asn Ser Tyr His Ser Arg Asn
370          375          380
Cys Ser Glu Ser Asp His Ser Arg Asn Gly Phe Asp Thr Asp Ser Ser
385          390          395          400
Cys Leu Glu Ser His Ser Ser Leu Ser Asp Ser Glu Phe Pro Pro Asn
          405          410          415
Asn Lys Gly Glu Ile Lys Thr Glu Gly Gln Glu Leu Ile Thr Val Ile
          420          425          430
Lys Ala Pro Thr Ser Phe Gly Tyr Asp Lys Pro His Val Leu Val Asp
          435          440          445
Leu Leu Val Asp Asp Ser Gly Lys Glu Ser Leu Ile Gly Tyr Arg Pro
          450          455          460
Thr Glu Asp Ser Lys Glu Phe Ser
465          470

```

```

<210> 137
<211> 143
<212> PRT
<213> Homo sapiens

```

```

<400> 137
Gln Asp Pro Tyr Val Lys Glu Ala Glu Asn Leu Lys Lys Tyr Phe Asn
1          5          10          15
Ala Gly His Ser Asp Val Ala Asp Asn Gly Thr Leu Phe Leu Gly Ile
          20          25          30
Leu Lys Asn Trp Lys Glu Glu Ser Asp Arg Lys Ile Met Gln Ser Gln
          35          40          45
Ile Val Ser Phe Tyr Phe Lys Leu Phe Lys Asn Phe Lys Asp Asp Gln
          50          55          60
Ser Ile Gln Lys Ser Val Glu Thr Ile Lys Glu Asp Met Asn Val Lys
65          70          75          80
Phe Phe Asn Ser Asn Lys Lys Lys Arg Asp Asp Phe Glu Lys Leu Thr
          85          90          95
Asn Tyr Ser Val Thr Asp Leu Asn Val Gln Arg Lys Ala Ile His Glu
          100          105          110
Leu Ile Gln Val Met Ala Glu Leu Ser Pro Ala Ala Lys Thr Gly Lys
          115          120          125
Arg Lys Arg Ser Gln Met Leu Phe Arg Gly Arg Arg Ala Ser Gln
130          135          140

```

```

<210> 138
<211> 143
<212> PRT

```

&lt;213&gt; Homo sapiens

&lt;400&gt; 138

```

Met Glu Ser Pro Ser Ala Pro Pro His Arg Trp Cys Ile Pro Trp Gln
1          5          10          15
Arg Leu Leu Leu Thr Ala Ser Leu Leu Thr Phe Trp Asn Pro Pro Thr
          20          25          30
Thr Ala Lys Leu Thr Ile Glu Ser Thr Pro Phe Asn Val Ala Glu Gly
          35          40          45
Lys Glu Val Leu Leu Leu Val His Asn Leu Pro Gln His Leu Phe Gly
          50          55          60
Tyr Ser Trp Tyr Lys Gly Glu Arg Val Asp Gly Asn Arg Gln Ile Ile
65          70          75          80
Gly Tyr Val Ile Gly Thr Gln Gln Ala Thr Pro Gly Pro Ala Tyr Ser
          85          90          95
Gly Arg Glu Ile Ile Tyr Pro Asn Ala Ser Leu Leu Ile Gln Asn Ile
          100          105          110
Ile Gln Asn Asp Thr Gly Phe Tyr Thr Leu His Val Ile Lys Ser Asp
          115          120          125
Leu Val Asn Glu Glu Ala Thr Gly Gln Phe Arg Val Tyr Arg Glu
          130          135          140

```

&lt;210&gt; 139

&lt;211&gt; 440

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 139

```

Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val
1          5          10          15
Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp
          20          25          30
Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu
          35          40          45
His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg
          50          55          60
Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu
65          70          75          80
Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg
          85          90          95
Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly
          100          105          110
Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg
          115          120          125
Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr
          130          135          140
His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro
145          150          155          160
Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile
          165          170          175
Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro
          180          185          190
Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile
          195          200          205
Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys
          210          215          220
Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val
225          230          235          240
Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys
          245          250          255
Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala
          260          265          270

```

Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met  
 275 280 285  
 Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln  
 290 295 300  
 Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr  
 305 310 315 320  
 Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val  
 325 330 335  
 Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile  
 340 345 350  
 Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala  
 355 360 365  
 Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr  
 370 375 380  
 Asn Ile Pro Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val  
 385 390 395 400  
 Met Ala Ala Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val  
 405 410 415  
 Cys Gln Trp Arg Cys Leu Arg Cys Leu Arg Gln Gln His Asp Asp Phe  
 420 425 430  
 Ala Asp Asp Ile Ser Leu Leu Lys  
 435 440

<210> 140  
 <211> 810  
 <212> PRT  
 <213> Homo sapiens

<400> 140  
 Met Glu His Lys Glu Val Val Leu Leu Leu Leu Phe Leu Lys Ser  
 1 5 10 15  
 Gly Gln Gly Glu Pro Leu Asp Asp Tyr Val Asn Thr Gln Gly Ala Ser  
 20 25 30  
 Leu Phe Ser Val Thr Lys Lys Gln Leu Gly Ala Gly Ser Ile Glu Glu  
 35 40 45  
 Cys Ala Ala Lys Cys Glu Glu Asp Glu Glu Phe Thr Cys Arg Ala Phe  
 50 55 60  
 Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Arg  
 65 70 75 80  
 Lys Ser Ser Ile Ile Ile Arg Met Arg Asp Val Val Leu Phe Glu Lys  
 85 90 95  
 Lys Val Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg  
 100 105 110  
 Gly Thr Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser  
 115 120 125  
 Ser Thr Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser  
 130 135 140  
 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln  
 145 150 155 160  
 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys  
 165 170 175  
 Asp Ile Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn  
 180 185 190  
 Tyr Asp Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala  
 195 200 205  
 Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe  
 210 215 220  
 Pro Asn Lys Asn Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu  
 225 230 235 240  
 Leu Arg Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu  
 245 250 255  
 Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Ser Gly Pro Thr  
 260 265 270

Tyr Gln Cys Leu Lys Gly Thr Gly Glu Asn Tyr Arg Gly Asn Val Ala  
 275 280 285  
 Val Thr Val Ser Gly His Thr Cys Gln His Trp Ser Ala Gln Thr Pro  
 290 295 300  
 His Thr His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Asp  
 305 310 315 320  
 Glu Asn Tyr Cys Arg Asn Pro Asp Gly Lys Arg Ala Pro Trp Cys His  
 325 330 335  
 Thr Thr Asn Ser Gln Val Arg Trp Glu Tyr Cys Lys Ile Pro Ser Cys  
 340 345 350  
 Asp Ser Ser Pro Val Ser Thr Glu Gln Leu Ala Pro Thr Ala Pro Pro  
 355 360 365  
 Glu Leu Thr Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Gln Ser  
 370 375 380  
 Tyr Arg Gly Thr Ser Ser Thr Thr Thr Thr Gly Lys Lys Cys Gln Ser  
 385 390 395 400  
 Trp Ser Ser Met Thr Pro His Arg His Gln Lys Thr Pro Glu Asn Tyr  
 405 410 415  
 Pro Asn Ala Gly Leu Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp  
 420 425 430  
 Lys Gly Pro Trp Cys Phe Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr  
 435 440 445  
 Cys Asn Leu Lys Lys Cys Ser Gly Thr Glu Ala Ser Val Val Ala Pro  
 450 455 460  
 Pro Pro Val Val Leu Leu Pro Asp Val Glu Thr Pro Ser Glu Glu Asp  
 465 470 475 480  
 Cys Met Phe Gly Asn Gly Lys Gly Tyr Arg Gly Lys Arg Ala Thr Thr  
 485 490 495  
 Val Thr Gly Thr Pro Cys Gln Asp Trp Ala Ala Gln Glu Pro His Arg  
 500 505 510  
 His Ser Ile Phe Thr Pro Glu Thr Asn Pro Arg Ala Gly Leu Glu Lys  
 515 520 525  
 Asn Tyr Cys Arg Asn Pro Asp Gly Asp Val Gly Gly Pro Trp Cys Tyr  
 530 535 540  
 Thr Thr Asn Pro Arg Lys Leu Tyr Asp Tyr Cys Asp Val Pro Gln Cys  
 545 550 555 560  
 Ala Ala Pro Ser Phe Asp Cys Gly Lys Pro Gln Val Glu Pro Lys Lys  
 565 570 575  
 Cys Pro Gly Arg Val Val Gly Gly Cys Val Ala His Pro His Ser Trp  
 580 585 590  
 Pro Trp Gln Val Ser Leu Arg Thr Arg Phe Gly Met His Phe Cys Gly  
 595 600 605  
 Gly Thr Leu Ile Ser Pro Glu Trp Val Leu Thr Ala Ala His Cys Leu  
 610 615 620  
 Glu Lys Ser Pro Arg Pro Ser Ser Tyr Lys Val Ile Leu Gly Ala His  
 625 630 635 640  
 Gln Glu Val Asn Leu Glu Pro His Val Gln Glu Ile Glu Val Ser Arg  
 645 650 655  
 Leu Phe Leu Glu Pro Thr Arg Lys Asp Ile Ala Leu Leu Lys Leu Ser  
 660 665 670  
 Ser Pro Ala Val Ile Thr Asp Lys Val Ile Pro Ala Cys Leu Pro Ser  
 675 680 685  
 Pro Asn Tyr Val Val Ala Asp Arg Thr Glu Cys Phe Ile Thr Gly Trp  
 690 695 700  
 Gly Glu Thr Gln Gly Thr Phe Gly Ala Gly Leu Leu Lys Glu Ala Gln  
 705 710 715 720  
 Leu Pro Val Ile Glu Asn Lys Val Cys Asn Arg Tyr Glu Phe Leu Asn  
 725 730 735  
 Gly Arg Val Gln Ser Thr Glu Leu Cys Ala Gly His Leu Ala Gly Gly  
 740 745 750  
 Thr Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys Phe Glu  
 755 760 765  
 Lys Asp Lys Tyr Ile Leu Gln Gly Val Thr Ser Trp Gly Leu Gly Cys

770		775		780
Ala Arg Pro Asn Lys Pro Gly Val Tyr Val Arg Val Ser Arg Phe Val				
785		790		795
Thr Trp Ile Glu Gly Val Met Arg Asn Asn				800
	805		810	

<210> 141  
 <211> 762  
 <212> PRT  
 <213> Homo sapiens

<400> 141

Gly Pro Asn Ile Cys Thr Thr Arg Gly Val Ser Ser Cys Gln Gln Cys				
1	5	10	15	
Leu Ala Val Ser Pro Met Cys Ala Trp Cys Ser Asp Glu Ala Leu Pro				
	20	25	30	
Leu Gly Ser Pro Arg Cys Asp Leu Lys Glu Asn Leu Leu Lys Asp Asn				
	35	40	45	
Cys Ala Pro Glu Ser Ile Glu Phe Pro Val Ser Glu Ala Arg Val Leu				
	50	55	60	
Glu Asp Arg Pro Leu Ser Asp Lys Gly Ser Gly Asp Ser Ser Gln Val				
65	70	75	80	
Thr Gln Val Ser Pro Gln Arg Ile Ala Leu Arg Leu Arg Pro Asp Asp				
	85	90	95	
Ser Lys Asn Phe Ser Ile Gln Val Arg Gln Val Glu Asp Tyr Pro Val				
	100	105	110	
Asp Ile Tyr Tyr Leu Met Asp Leu Ser Tyr Ser Met Lys Asp Asp Leu				
	115	120	125	
Trp Ser Ile Gln Asn Leu Gly Thr Lys Leu Ala Thr Gln Met Arg Lys				
	130	135	140	
Leu Thr Ser Asn Leu Arg Ile Gly Phe Gly Ala Phe Val Asp Lys Pro				
145	150	155	160	
Val Ser Pro Tyr Met Tyr Ile Ser Pro Pro Glu Ala Leu Glu Asn Pro				
	165	170	175	
Cys Tyr Asp Met Lys Thr Thr Cys Leu Pro Met Phe Gly Tyr Lys His				
	180	185	190	
Val Leu Thr Leu Thr Asp Gln Val Thr Arg Phe Asn Glu Glu Val Lys				
	195	200	205	
Lys Gln Ser Val Ser Arg Asn Arg Asp Ala Pro Glu Gly Gly Phe Asp				
	210	215	220	
Ala Ile Met Gln Ala Thr Val Cys Asp Glu Lys Ile Gly Trp Arg Asn				
225	230	235	240	
Asp Ala Ser His Leu Leu Val Phe Thr Thr Asp Ala Lys Thr His Ile				
	245	250	255	
Ala Leu Asp Gly Arg Leu Ala Gly Ile Val Gln Pro Asn Asp Gly Gln				
	260	265	270	
Cys His Val Gly Ser Asp Asn His Tyr Ser Ala Ser Thr Thr Met Asp				
	275	280	285	
Tyr Pro Ser Leu Gly Leu Met Thr Glu Lys Leu Ser Gln Lys Asn Ile				
	290	295	300	
Asn Leu Ile Phe Ala Val Thr Glu Asn Val Val Asn Leu Tyr Gln Asn				
305	310	315	320	
Tyr Ser Glu Leu Ile Pro Gly Thr Thr Val Gly Val Leu Ser Met Asp				
	325	330	335	
Ser Ser Asn Val Leu Gln Leu Ile Val Asp Ala Tyr Gly Lys Ile Arg				
	340	345	350	
Ser Lys Val Glu Leu Glu Val Arg Asp Leu Pro Glu Glu Leu Ser Leu				
	355	360	365	
Ser Phe Asn Ala Thr Cys Leu Asn Asn Glu Val Ile Pro Gly Leu Lys				
	370	375	380	
Ser Cys Met Gly Leu Lys Ile Gly Asp Thr Val Ser Phe Ser Ile Glu				
385	390	395	400	

Ala	Lys	Val	Arg	Gly	Cys	Pro	Gln	Glu	Lys	Glu	Lys	Ser	Phe	Thr	Ile
				405					410					415	
Lys	Pro	Val	Gly	Phe	Lys	Asp	Ser	Leu	Ile	Val	Gln	Val	Thr	Phe	Asp
			420					425					430		
Cys	Asp	Cys	Ala	Cys	Gln	Ala	Gln	Ala	Glu	Pro	Asn	Ser	His	Arg	Cys
			435				440					445			
Asn	Asn	Gly	Asn	Gly	Thr	Phe	Glu	Cys	Gly	Val	Cys	Arg	Cys	Gly	Pro
			450			455				460					
Gly	Trp	Leu	Gly	Ser	Gln	Cys	Glu	Cys	Ser	Glu	Glu	Asp	Tyr	Arg	Pro
465				470					475					480	
Ser	Gln	Gln	Asp	Glu	Cys	Ser	Pro	Arg	Glu	Gly	Gln	Pro	Val	Cys	Ser
			485						490					495	
Gln	Arg	Gly	Glu	Cys	Leu	Cys	Gly	Gln	Cys	Val	Cys	His	Ser	Ser	Asp
			500				505					510			
Phe	Gly	Lys	Ile	Thr	Gly	Lys	Tyr	Cys	Glu	Cys	Asp	Asp	Phe	Ser	Cys
			515				520				525				
Val	Arg	Tyr	Lys	Gly	Glu	Met	Cys	Ser	Gly	His	Gly	Gln	Cys	Ser	Cys
			530			535				540					
Gly	Asp	Cys	Leu	Cys	Asp	Ser	Asp	Trp	Thr	Gly	Tyr	Tyr	Cys	Asn	Cys
545				550					555					560	
Thr	Thr	Arg	Thr	Asp	Thr	Cys	Met	Ser	Ser	Asn	Gly	Leu	Leu	Cys	Ser
			565					570						575	
Gly	Arg	Gly	Lys	Cys	Glu	Cys	Gly	Ser	Cys	Val	Cys	Ile	Gln	Pro	Gly
			580				585					590			
Ser	Tyr	Gly	Asp	Thr	Cys	Glu	Lys	Cys	Pro	Thr	Cys	Pro	Asp	Ala	Cys
			595			600				605					
Thr	Phe	Lys	Lys	Glu	Cys	Val	Glu	Cys	Lys	Lys	Phe	Asp	Arg	Glu	Pro
			610			615				620					
Tyr	Met	Thr	Glu	Asn	Thr	Cys	Asn	Arg	Tyr	Cys	Arg	Asp	Glu	Ile	Glu
625				630					635					640	
Ser	Val	Lys	Glu	Leu	Lys	Asp	Thr	Gly	Lys	Asp	Ala	Val	Asn	Cys	Thr
			645					650					655		
Tyr	Lys	Asn	Glu	Asp	Asp	Cys	Val	Val	Arg	Phe	Gln	Tyr	Tyr	Glu	Asp
			660				665					670			
Ser	Ser	Gly	Lys	Ser	Ile	Leu	Tyr	Val	Val	Glu	Glu	Pro	Glu	Cys	Pro
			675			680				685					
Lys	Gly	Pro	Asp	Ile	Leu	Val	Val	Leu	Leu	Ser	Val	Met	Gly	Ala	Ile
			690			695				700					
Leu	Leu	Ile	Gly	Leu	Ala	Ala	Leu	Leu	Ile	Trp	Lys	Leu	Leu	Ile	Thr
705				710					715					720	
Ile	His	Asp	Arg	Lys	Glu	Phe	Ala	Lys	Phe	Glu	Glu	Glu	Arg	Ala	Arg
			725					730					735		
Ala	Lys	Trp	Asp	Thr	Ala	Asn	Asn	Pro							

```
<210> 142
<211> 505
<212> PRT
<213> Homo sapiens
```

<400> 142																
Gln	Thr	Ser	Val	Ser	Pro	Ser	Lys	Val	Ile	Leu	Pro	Arg	Gly	Gly	Ser	
1				5					10					15		
Val	Leu	Val	Thr	Cys	Ser	Thr	Ser	Cys	Asp	Gln	Pro	Lys	Leu	Leu	Gly	
			20					25					30			
Ile	Glu	Thr	Pro	Leu	Pro	Lys	Lys	Glu	Leu	Leu	Leu	Pro	Gly	Asn	Asn	
		35					40					45				
Arg	Lys	Val	Tyr	Glu	Leu	Ser	Asn	Val	Gln	Glu	Asp	Ser	Gln	Pro	Met	
	50					55					60					
Cys	Tyr	Ser	Asn	Cys	Pro	Asp	Gly	Gln	Ser	Thr	Ala	Lys	Thr	Phe	Leu	

65					70					75				80	
Thr	Val	Tyr	Trp	Thr	Pro	Glu	Arg	Val	Glu	Leu	Ala	Pro	Leu	Pro	Ser
				85					90					95	
Trp	Gln	Pro	Val	Gly	Lys	Asn	Leu	Thr	Leu	Arg	Cys	Gln	Val	Glu	Gly
			100					105					110		
Gly	Ala	Pro	Arg	Ala	Asn	Leu	Thr	Val	Val	Leu	Leu	Arg	Gly	Glu	Lys
		115					120					125			
Glu	Leu	Lys	Arg	Glu	Pro	Ala	Val	Gly	Glu	Pro	Ala	Glu	Val	Thr	Thr
	130					135					140				
Thr	Val	Leu	Val	Arg	Arg	Asp	His	His	Gly	Ala	Asn	Phe	Ser	Cys	Arg
145					150					155				160	
Thr	Glu	Leu	Asp	Leu	Arg	Pro	Gln	Gly	Leu	Glu	Leu	Phe	Glu	Asn	Thr
			165					170					175		
Ser	Ala	Pro	Tyr	Gln	Leu	Gln	Thr	Phe	Val	Leu	Pro	Ala	Thr	Pro	Pro
			180				185						190		
Gln	Leu	Val	Ser	Pro	Arg	Val	Leu	Glu	Val	Asp	Thr	Gln	Gly	Thr	Val
	195					200						205			
Val	Cys	Ser	Leu	Asp	Gly	Leu	Phe	Pro	Val	Ser	Glu	Ala	Gln	Val	His
	210					215					220				
Leu	Ala	Leu	Gly	Asp	Gln	Arg	Leu	Asn	Pro	Thr	Val	Thr	Tyr	Gly	Asn
225					230					235				240	
Asp	Ser	Phe	Ser	Ala	Lys	Ala	Ser	Val	Ser	Val	Thr	Ala	Glu	Asp	Glu
			245					250					255		
Gly	Thr	Gln	Arg	Leu	Thr	Cys	Ala	Val	Ile	Leu	Gly	Asn	Gln	Ser	Gln
		260					265						270		
Glu	Thr	Leu	Gln	Thr	Val	Thr	Ile	Tyr	Ser	Phe	Pro	Ala	Pro	Asn	Val
	275					280						285			
Ile	Leu	Thr	Lys	Pro	Glu	Val	Ser	Glu	Gly	Thr	Glu	Val	Thr	Val	Lys
	290				295						300				
Cys	Glu	Ala	His	Pro	Arg	Ala	Lys	Val	Thr	Leu	Asn	Gly	Val	Pro	Ala
305					310					315				320	
Gln	Pro	Leu	Gly	Pro	Arg	Ala	Gln	Leu	Leu	Lys	Ala	Thr	Pro	Glu	
			325					330					335		
Asp	Asn	Gly	Arg	Ser	Phe	Ser	Cys	Ser	Ala	Thr	Leu	Glu	Val	Ala	Gly
		340					345					350			
Gln	Leu	Ile	His	Lys	Asn	Gln	Thr	Arg	Glu	Leu	Arg	Val	Leu	Tyr	Gly
	355					360						365			
Pro	Arg	Leu	Asp	Glu	Arg	Asp	Cys	Pro	Gly	Asn	Trp	Thr	Trp	Pro	Glu
	370					375					380				
Asn	Ser	Gln	Gln	Thr	Pro	Met	Cys	Gln	Ala	Trp	Gly	Asn	Pro	Leu	Pro
385					390					395				400	
Glu	Leu	Lys	Cys	Leu	Lys	Asp	Gly	Thr	Phe	Pro	Leu	Pro	Ile	Gly	Glu
			405					410					415		
Ser	Val	Thr	Val	Thr	Arg	Asp	Leu	Glu	Gly	Thr	Tyr	Leu	Cys	Arg	Ala
		420					425					430			
Arg	Ser	Thr	Gln	Gly	Glu	Val	Thr	Arg	Glu	Val	Thr	Val	Asn	Val	Leu
	435					440					445				
Ser	Pro	Arg	Tyr	Glu	Ile	Val	Ile	Ile	Thr	Val	Val	Ala	Ala	Ala	Val
	450					455					460				
Ile	Met	Gly	Thr	Ala	Gly	Leu	Ser	Thr	Tyr	Leu	Tyr	Asn	Arg	Gln	Arg
465					470					475				480	
Lys	Ile	Lys	Lys	Tyr	Arg	Leu	Gln	Gln	Ala	Gln	Lys	Gly	Thr	Pro	Met
			485					490					495		
Lys	Pro	Asn	Thr	Gln	Ala	Thr	Pro	Pro							
		500					505								

<210> 143  
 <211> 261  
 <212> PRT  
 <213> Homo sapiens

<400> 143



```

Met Ile Glu Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly
1      5      10      15
Leu Pro Ile Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu
20      25      30
Ile Thr Gln Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg
35      40      45
Arg Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val
50      55      60
Phe Met Lys Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser
65      70      75      80
Leu Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys
85      90      95
Asp Ile Met Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu
100      105      110
Met Gln Lys Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser
115      120      125
Glu Ala Ser Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly
130      135      140
Tyr Tyr Thr Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln
145      150      155      160
Leu Thr Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr
165      170      175
Phe Cys Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser
180      185      190
Leu Cys Leu Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala
195      200      205
Ala Asn Thr His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His
210      215      220
Leu Gly Gly Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn
225      230      235      240
Val Thr Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe
245      250      255
Gly Leu Leu Lys Leu
260

```

```

<210> 144
<211> 187
<212> PRT
<213> Homo sapiens

```

```

<400> 144
Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly
1      5      10      15
Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu
20      25      30
Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val
35      40      45
Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp
50      55      60
Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile
65      70      75      80
Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu
85      90      95
Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln
100      105      110
Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Phe Leu Leu
115      120      125
Trp Ile Leu Ala Ala Val Ser Ser Gly Leu Phe Phe Tyr Ser Phe Leu
130      135      140
Leu Thr Ala Val Ser Leu Ser Lys Met Leu Lys Lys Arg Ser Pro Leu
145      150      155      160
Thr Thr Gly Val Tyr Val Lys Met Pro Pro Thr Glu Pro Glu Cys Glu

```

165 170 175  
 Lys Gln Phe Gln Pro Tyr Phe Ile Pro Ile Asn  
 180 185  
  
 <210> 145  
 <211> 544  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 145  
 Ile Pro Pro His Val Gln Lys Ser Val Asn Asn Asp Met Ile Val Thr  
 1 5 10 15  
 Asp Asn Asn Gly Ala Val Lys Phe Pro Gln Leu Cys Lys Phe Cys Asp  
 20 25 30  
 Val Arg Phe Ser Thr Cys Asp Asn Gln Lys Ser Cys Met Ser Asn Cys  
 35 40 45  
 Ser Ile Thr Ser Ile Cys Glu Lys Pro Gln Glu Val Cys Val Ala Val  
 50 55 60  
 Trp Arg Lys Asn Asp Glu Asn Ile Thr Leu Glu Thr Val Cys His Asp  
 65 70 75 80  
 Pro Lys Leu Pro Tyr His Asp Phe Ile Leu Glu Asp Ala Ala Ser Pro  
 85 90 95  
 Lys Cys Ile Met Lys Glu Lys Lys Lys Pro Gly Glu Thr Phe Phe Met  
 100 105 110  
 Cys Ser Cys Ser Ser Asp Glu Cys Asn Asp Asn Ile Ile Phe Ser Glu  
 115 120 125  
 Glu Tyr Asn Thr Ser Asn Pro Asp Leu Leu Leu Val Ile Phe Gln Val  
 130 135 140  
 Thr Gly Ile Ser Leu Leu Pro Pro Leu Gly Val Ala Ile Ser Val Ile  
 145 150 155 160  
 Ile Ile Phe Tyr Cys Tyr Arg Val Asn Arg Gln Gln Lys Leu Ser Ser  
 165 170 175  
 Thr Trp Glu Thr Gly Lys Thr Arg Lys Leu Met Glu Phe Ser Glu His  
 180 185 190  
 Cys Ala Ile Ile Leu Glu Asp Asp Arg Ser Asp Ile Ser Ser Thr Cys  
 195 200 205  
 Ala Asn Asn Ile Asn His Asn Thr Glu Leu Leu Pro Ile Glu Leu Asp  
 210 215 220  
 Thr Leu Val Gly Lys Gly Arg Phe Ala Glu Val Tyr Lys Ala Lys Leu  
 225 230 235 240  
 Lys Gln Asn Thr Ser Glu Gln Phe Glu Thr Val Ala Val Lys Ile Phe  
 245 250 255  
 Pro Tyr Glu Glu Tyr Ala Ser Trp Lys Thr Glu Lys Asp Ile Phe Ser  
 260 265 270  
 Asp Ile Asn Leu Lys His Glu Asn Ile Leu Gln Phe Leu Thr Ala Glu  
 275 280 285  
 Glu Arg Lys Thr Glu Leu Gly Lys Gln Tyr Trp Leu Ile Thr Ala Phe  
 290 295 300  
 His Ala Lys Gly Asn Leu Gln Glu Tyr Leu Thr Arg His Val Ile Ser  
 305 310 315 320  
 Trp Glu Asp Leu Arg Lys Leu Gly Ser Ser Leu Ala Arg Gly Ile Ala  
 325 330 335  
 His Leu His Ser Asp His Thr Pro Cys Gly Arg Pro Lys Met Pro Ile  
 340 345 350  
 Val His Arg Asp Leu Lys Ser Ser Asn Ile Leu Val Lys Asn Asp Leu  
 355 360 365  
 Thr Cys Cys Leu Cys Asp Phe Gly Leu Ser Leu Arg Leu Asp Pro Thr  
 370 375 380  
 Leu Ser Val Asp Asp Leu Ala Asn Ser Gly Gln Val Gly Thr Ala Arg  
 385 390 395 400  
 Tyr Met Ala Pro Glu Val Leu Glu Ser Arg Met Asn Leu Glu Asn Ala  
 405 410 415

Glu Ser Phe Lys Gln Thr Asp Val Tyr Ser Met Ala Leu Val Leu Trp  
                   420                  425                  430  
 Glu Met Thr Ser Arg Cys Asn Ala Val Gly Glu Val Lys Asp Tyr Glu  
                   435                  440                  445  
 Pro Pro Phe Gly Ser Lys Val Arg Glu His Pro Cys Val Glu Ser Met  
                   450                  455                  460  
 Lys Asp Asn Val Leu Arg Asp Arg Gly Arg Pro Glu Ile Pro Ser Phe  
 465                  470                  475                  480  
 Trp Leu Asn His Gln Gly Ile Gln Met Val Cys Glu Thr Leu Thr Glu  
                   485                  490                  495  
 Cys Trp Asp His Asp Pro Glu Ala Arg Leu Thr Ala Gln Cys Val Ala  
                   500                  505                  510  
 Glu Arg Phe Ser Glu Leu Glu His Leu Asp Arg Leu Ser Gly Arg Ser  
                   515                  520                  525  
 Cys Ser Glu Glu Lys Ile Pro Glu Asp Gly Ser Leu Asn Thr Thr Lys  
                   530                  535                  540

<210> 146  
 <211> 358  
 <212> PRT  
 <213> Homo sapiens

<400> 146  
 Cys Glu Glu Pro Pro Thr Phe Glu Ala Met Glu Leu Ile Gly Lys Pro  
 1                  5                  10                  15  
 Lys Pro Tyr Tyr Glu Ile Gly Glu Arg Val Asp Tyr Lys Cys Lys Lys  
                   20                  25                  30  
 Gly Tyr Phe Tyr Ile Pro Pro Leu Ala Thr His Thr Ile Cys Asp Arg  
                   35                  40                  45  
 Asn His Thr Trp Leu Pro Val Ser Asp Asp Ala Cys Tyr Arg Glu Thr  
                   50                  55                  60  
 Cys Pro Tyr Ile Arg Asp Pro Leu Asn Gly Gln Ala Val Pro Ala Asn  
 65                  70                  75                  80  
 Gly Thr Tyr Glu Phe Gly Tyr Gln Met His Phe Ile Cys Asn Glu Gly  
                   85                  90                  95  
 Tyr Tyr Leu Ile Gly Glu Glu Ile Leu Tyr Cys Glu Leu Lys Gly Ser  
                   100                  105                  110  
 Val Ala Ile Trp Ser Gly Lys Pro Pro Ile Cys Glu Lys Val Leu Cys  
                   115                  120                  125  
 Thr Pro Pro Pro Lys Ile Lys Asn Gly Lys His Thr Phe Ser Glu Val  
                   130                  135                  140  
 Glu Val Phe Glu Tyr Leu Asp Ala Val Thr Tyr Ser Cys Asp Pro Ala  
 145                  150                  155                  160  
 Pro Gly Pro Asp Pro Phe Ser Leu Ile Gly Glu Ser Thr Ile Tyr Cys  
                   165                  170                  175  
 Gly Asp Asn Ser Val Trp Ser Arg Ala Ala Pro Glu Cys Lys Val Val  
                   180                  185                  190  
 Lys Cys Arg Phe Pro Val Val Glu Asn Gly Lys Gln Ile Ser Gly Phe  
                   195                  200                  205  
 Gly Lys Lys Phe Tyr Tyr Lys Ala Thr Val Met Phe Glu Cys Asp Lys  
                   210                  215                  220  
 Gly Phe Tyr Leu Asp Gly Ser Asp Thr Ile Val Cys Asp Ser Asn Ser  
 225                  230                  235                  240  
 Thr Trp Asp Pro Pro Val Pro Lys Cys Leu Lys Val Leu Pro Pro Ser  
                   245                  250                  255  
 Ser Thr Lys Pro Pro Ala Leu Ser His Ser Val Ser Thr Ser Ser Thr  
                   260                  265                  270  
 Thr Lys Ser Pro Ala Ser Ser Ala Ser Gly Pro Arg Pro Thr Tyr Lys  
                   275                  280                  285  
 Pro Pro Val Ser Asn Tyr Pro Gly Tyr Pro Lys Pro Glu Glu Gly Ile  
                   290                  295                  300  
 Leu Asp Ser Leu Asp Val Trp Val Ile Ala Val Ile Val Ile Ala Ile

[illegible]

```
<210> 147
<211> 1148
<212> PRT
<213> Homo sapiens
```

<div>&lt;400&gt;</div>	<div>147</div>														
Leu	Pro	Glu	Ala	Lys	Ile	Phe	Ser	Gly	Pro	Ser	Ser	Glu	Gln	Phe	Gly
1				5					10					15	
Tyr	Ala	Val	Gln	Gln	Phe	Ile	Asn	Pro	Lys	Gly	Asn	Trp	Leu	Leu	Val
			20						25				30		
Gly	Ser	Pro	Trp	Ser	Gly	Phe	Pro	Glu	Asn	Arg	Met	Gly	Asp	Val	Tyr
		35					40					45			
Lys	Cys	Pro	Val	Asp	Leu	Ser	Thr	Ala	Thr	Cys	Glu	Lys	Leu	Asn	Leu
	50					55					60				
Gln	Thr	Ser	Thr	Ser	Ile	Pro	Asn	Val	Thr	Glu	Met	Lys	Thr	Asn	Met
65					70					75					80
Ser	Leu	Gly	Leu	Ile	Leu	Thr	Arg	Asn	Met	Gly	Thr	Gly	Gly	Phe	Leu
				85					90					95	
Thr	Cys	Gly	Pro	Leu	Trp	Ala	Gln	Gln	Cys	Gly	Asn	Gln	Tyr	Tyr	Thr
			100					105					110		
Thr	Gly	Val	Cys	Ser	Asp	Ile	Ser	Pro	Asp	Phe	Gln	Leu	Ser	Ala	Ser
		115					120					125			
Phe	Ser	Pro	Ala	Thr	Gln	Pro	Cys	Pro	Ser	Leu	Ile	Asp	Val	Val	Val
		130				135					140				
Val	Cys	Asp	Glu	Ser	Asn	Ser	Ile	Tyr	Pro	Trp	Asp	Ala	Val	Lys	Asn
145					150					155					160
Phe	Leu	Glu	Lys	Phe	Val	Gln	Gly	Leu	Asp	Ile	Gly	Pro	Thr	Lys	Thr
				165					170					175	
Gln	Val	Gly	Leu	Ile	Gln	Tyr	Ala	Asn	Asn	Pro	Arg	Val	Val	Phe	Asn
			180					185					190		
Leu	Asn	Thr	Tyr	Lys	Thr	Lys	Glu	Glu	Met	Ile	Val	Ala	Thr	Ser	Gln
		195					200					205			
Thr	Ser	Gln	Tyr	Gly	Gly	Asp	Leu	Thr	Asn	Thr	Phe	Gly	Ala	Ile	Gln
		210				215					220				
Tyr	Ala	Arg	Lys	Tyr	Ala	Tyr	Ser	Ala	Ala	Ser	Gly	Gly	Arg	Arg	Ser
225					230					235					240
Ala	Thr	Lys	Val	Met	Val	Val	Val	Thr	Asp	Gly	Glu	Ser	His	Asp	Gly
				245					250					255	
Ser	Met	Leu	Lys	Ala	Val	Ile	Asp	Gln	Cys	Asn	His	Asp	Asn	Ile	Leu
			260				265						270		
Arg	Phe	Gly	Ile	Ala	Val	Leu	Gly	Tyr	Leu	Asn	Arg	Asn	Ala	Leu	Asp
		275					280					285			
Thr	Lys	Asn	Leu	Ile	Lys	Glu	Ile	Lys	Ala	Ile	Ala	Ser	Ile	Pro	Thr
		290				295					300				
Glu	Arg	Tyr	Phe	Phe	Asn	Val	Ser	Asp	Glu	Ala	Ala	Leu	Leu	Glu	Lys
305					310					315					320
Ala	Gly	Thr	Leu	Gly	Glu	Gln	Ile	Phe	Ser	Ile	Glu	Gly	Thr	Val	Gln
				325					330					335	
Gly	Gly	Asp	Asn	Phe	Gln	Met	Glu	Met	Ser	Gln	Val	Gly	Phe	Ser	

385	390										395					400		
Ser	Ser	Tyr	Leu	Gly	Tyr	Ser	Val	Ala	Ala	Ile	Ser	Thr	Gly	Glu	Ser			
				405					410					415				
Thr	His	Phe	Val	Ala	Gly	Ala	Pro	Arg	Ala	Asn	Tyr	Thr	Gly	Gln	Ile			
				420				425					430					
Val	Leu	Tyr	Ser	Val	Asn	Glu	Asn	Gly	Asn	Ile	Thr	Val	Ile	Gln	Ala			
				435			440					445						
His	Arg	Gly	Asp	Gln	Ile	Gly	Ser	Tyr	Phe	Gly	Ser	Val	Leu	Cys	Ser			
	450					455					460							
Val	Asp	Val	Asp	Lys	Asp	Thr	Ile	Thr	Asp	Val	Leu	Leu	Val	Gly	Ala			
465					470					475					480			
Pro	Met	Tyr	Met	Ser	Asp	Leu	Lys	Lys	Glu	Glu	Gly	Arg	Val	Tyr	Leu			
				485					490					495				
Phe	Thr	Ile	Lys	Lys	Gly	Ile	Leu	Gly	Gln	His	Gln	Phe	Leu	Glu	Gly			
				500				505					510					
Pro	Glu	Gly	Ile	Glu	Asn	Thr	Arg	Phe	Gly	Ser	Ala	Ile	Ala	Ala	Leu			
				515			520					525						
Ser	Asp	Ile	Asn	Met	Asp	Gly	Phe	Asn	Asp	Val	Ile	Val	Gly	Ser	Pro			
						535					540							
Leu	Glu	Asn	Gln	Asn	Ser	Gly	Ala	Val	Tyr	Ile	Tyr	Asn	Gly	His	Gln			
545					550					555					560			
Gly	Thr	Ile	Arg	Thr	Lys	Tyr	Ser	Gln	Lys	Ile	Leu	Gly	Ser	Asp	Gly			
				565					570					575				
Ala	Phe	Arg	Ser	His	Leu	Gln	Tyr	Phe	Gly	Arg	Ser	Leu	Asp	Gly	Tyr			
				580				585					590					
Gly	Asp	Leu	Asn	Gly	Asp	Ser	Ile	Thr	Asp	Val	Ser	Ile	Gly	Ala	Phe			
				595			600					605						
Gly	Gln	Val	Val	Gln	Leu	Trp	Ser	Gln	Ser	Ile	Ala	Asp	Val	Ala	Ile			
						615					620							
Glu	Ala	Ser	Phe	Thr	Pro	Glu	Lys	Ile	Thr	Leu	Val	Asn	Lys	Asn	Ala			
625					630					635					640			
Gln	Ile	Ile	Leu	Lys	Leu	Cys	Phe	Ser	Ala	Lys	Phe	Arg	Pro	Thr	Lys			
				645					650					655				
Gln	Asn	Asn	Gln	Val	Ala	Ile	Val	Tyr	Asn	Ile	Thr	Leu	Asp	Ala	Asp			
				660				665					670					
Gly	Phe	Ser	Ser	Arg	Val	Thr	Ser	Arg	Gly	Leu	Phe	Lys	Glu	Asn	Asn			
				675			680					685						
Glu	Arg	Cys	Leu	Gln	Lys	Asn	Met	Val	Val	Asn	Gln	Ala	Gln	Ser	Cys			
						695					700							
Pro	Glu	His	Ile	Ile	Tyr	Ile	Gln	Glu	Pro	Ser	Asp	Val	Val	Asn	Ser			
705					710					715					720			
Leu	Asp	Leu	Arg	Val	Asp	Ile	Ser	Leu	Glu	Asn	Pro	Gly	Thr	Ser	Pro			
				725					730					735				
Ala	Leu	Glu	Ala	Tyr	Ser	Glu	Thr	Ala	Lys	Val	Phe	Ser	Ile	Pro	Phe			
				740				745					750					

Ile Pro Leu Leu Tyr Asp Ala Glu Ile His Leu Thr Arg Ser Thr Asn  
                   900                  905                  910  
 Ile Asn Phe Tyr Glu Ile Ser Ser Asp Gly Asn Val Pro Ser Ile Val  
                   915                  920                  925  
 His Ser Phe Glu Asp Val Gly Pro Lys Phe Ile Phe Ser Leu Lys Val  
                   930                  935                  940  
 Thr Thr Gly Ser Val Pro Val Ser Met Ala Thr Val Ile Ile His Ile  
                   945                  950                  955                  960  
 Pro Gln Tyr Thr Lys Glu Lys Asn Pro Leu Met Tyr Leu Thr Gly Val  
                   965                  970                  975  
 Gln Thr Asp Lys Ala Gly Asp Ile Ser Cys Asn Ala Asp Ile Asn Pro  
                   980                  985                  990  
 Leu Lys Ile Gly Gln Thr Ser Ser Ser Val Ser Phe Lys Ser Glu Asn  
                   995                  1000                  1005  
 Phe Arg His Thr Lys Glu Leu Asn Cys Arg Thr Ala Ser Cys Ser  
                   1010                  1015                  1020  
 Asn Val Thr Cys Trp Leu Lys Asp Val His Met Lys Gly Glu Tyr  
                   1025                  1030                  1035  
 Phe Val Asn Val Thr Thr Arg Ile Trp Asn Gly Thr Phe Ala Ser  
                   1040                  1045                  1050  
 Ser Thr Phe Gln Thr Val Gln Leu Thr Ala Ala Ala Glu Ile Asn  
                   1055                  1060                  1065  
 Thr Tyr Asn Pro Glu Ile Tyr Val Ile Glu Asp Asn Thr Val Thr  
                   1070                  1075                  1080  
 Ile Pro Leu Met Ile Met Lys Pro Asp Glu Lys Ala Glu Val Pro  
                   1085                  1090                  1095  
 Thr Gly Val Ile Ile Gly Ser Ile Ile Ala Gly Ile Leu Leu Leu  
                   1100                  1105                  1110  
 Leu Ala Leu Val Ala Ile Leu Trp Lys Leu Gly Phe Phe Lys Arg  
                   1115                  1120                  1125  
 Lys Tyr Glu Lys Met Thr Lys Asn Pro Asp Glu Ile Asp Glu Thr  
                   1130                  1135                  1140  
 Thr Glu Leu Ser Ser  
                   1145

<210> 148  
 <211> 133  
 <212> PRT  
 <213> Homo sapiens

<400> 148  
 Ala Pro Met Thr Gln Thr Thr Pro Leu Lys Thr Ser Trp Val Asn Cys  
 1                  5                  10                  15  
 Ser Asn Met Ile Asp Glu Ile Ile Thr His Leu Lys Gln Pro Pro Leu  
                   20                  25                  30  
 Pro Leu Leu Asp Phe Asn Asn Leu Asn Gly Glu Asp Gln Asp Ile Leu  
                   35                  40                  45  
 Met Glu Asn Asn Leu Arg Arg Pro Asn Leu Glu Ala Phe Asn Arg Ala  
                   50                  55                  60  
 Val Lys Ser Leu Gln Asn Ala Ser Ala Ile Glu Ser Ile Leu Lys Asn  
                   65                  70                  75                  80  
 Leu Leu Pro Cys Leu Pro Leu Ala Thr Ala Ala Pro Thr Arg His Pro  
                   85                  90                  95  
 Ile His Ile Lys Asp Gly Asp Trp Asn Glu Phe Arg Arg Lys Leu Thr  
                   100                  105                  110  
 Phe Tyr Leu Lys Thr Leu Glu Asn Ala Gln Ala Gln Gln Thr Thr Leu  
                   115                  120                  125  
 Ser Leu Ala Ile Phe  
                   130

<210> 149  
 <211> 622  
 <212> PRT  
 <213> Homo sapiens

<400> 149  
 Met Ala His Val Arg Gly Leu Gln Leu Pro Gly Cys Leu Ala Leu Ala  
 1 5 10 15  
 Ala Leu Cys Ser Leu Val His Ser Gln His Val Phe Leu Ala Pro Gln  
 20 25 30  
 Gln Ala Arg Ser Leu Leu Gln Arg Val Arg Arg Ala Asn Thr Phe Leu  
 35 40 45  
 Glu Glu Val Arg Lys Gly Asn Leu Glu Arg Glu Cys Val Glu Glu Thr  
 50 55 60  
 Cys Ser Tyr Glu Glu Ala Phe Glu Ala Leu Glu Ser Ser Thr Ala Thr  
 65 70 75 80  
 Asp Val Phe Trp Ala Lys Tyr Thr Ala Cys Glu Thr Ala Arg Thr Pro  
 85 90 95  
 Arg Asp Lys Leu Ala Ala Cys Leu Glu Gly Asn Cys Ala Glu Gly Leu  
 100 105 110  
 Gly Thr Asn Tyr Arg Gly His Val Asn Ile Thr Arg Ser Gly Ile Glu  
 115 120 125  
 Cys Gln Leu Trp Arg Ser Arg Tyr Pro His Lys Pro Glu Ile Asn Ser  
 130 135 140  
 Thr Thr His Pro Gly Ala Asp Leu Gln Glu Asn Phe Cys Arg Asn Pro  
 145 150 155 160  
 Asp Ser Ser Thr Thr Gly Pro Trp Cys Tyr Thr Thr Asp Pro Thr Val  
 165 170 175  
 Arg Arg Gln Glu Cys Ser Ile Pro Val Cys Gly Gln Asp Gln Val Thr  
 180 185 190  
 Val Ala Met Thr Pro Arg Ser Glu Gly Ser Ser Val Asn Leu Ser Pro  
 195 200 205  
 Pro Leu Glu Gln Cys Val Pro Asp Arg Gly Gln Gln Tyr Gln Gly Arg  
 210 215 220  
 Leu Ala Val Thr Thr His Gly Leu Pro Cys Leu Ala Trp Ala Ser Ala  
 225 230 235 240  
 Gln Ala Lys Ala Leu Ser Lys His Gln Asp Phe Asn Ser Ala Val Gln  
 245 250 255  
 Leu Val Glu Asn Phe Cys Arg Asn Pro Asp Gly Asp Glu Glu Gly Val  
 260 265 270  
 Trp Cys Tyr Val Ala Gly Lys Pro Gly Asp Phe Gly Tyr Cys Asp Leu  
 275 280 285  
 Asn Tyr Cys Glu Glu Ala Val Glu Glu Glu Thr Gly Asp Gly Leu Asp  
 290 295 300  
 Glu Asp Ser Asp Arg Ala Ile Glu Gly Arg Thr Ala Thr Ser Glu Tyr  
 305 310 315 320  
 Gln Thr Phe Phe Asn Pro Arg Thr Phe Gly Ser Gly Glu Ala Asp Cys  
 325 330 335  
 Gly Leu Arg Pro Leu Phe Glu Lys Lys Ser Leu Glu Asp Lys Thr Glu  
 340 345 350  
 Arg Glu Leu Leu Glu Ser Tyr Ile Asp Gly Arg Ile Val Glu Gly Ser  
 355 360 365  
 Asp Ala Glu Ile Gly Met Ser Pro Trp Gln Val Met Leu Phe Arg Lys  
 370 375 380  
 Ser Pro Gln Glu Leu Leu Cys Gly Ala Ser Leu Ile Ser Asp Arg Trp  
 385 390 395 400  
 Val Leu Thr Ala Ala His Cys Leu Leu Tyr Pro Pro Trp Asp Lys Asn  
 405 410 415  
 Phe Thr Glu Asn Asp Leu Leu Val Arg Ile Gly Lys His Ser Arg Thr  
 420 425 430  
 Arg Tyr Glu Arg Asn Ile Glu Lys Ile Ser Met Leu Glu Lys Ile Tyr  
 435 440 445  
 Ile His Pro Arg Tyr Asn Trp Arg Glu Asn Leu Asp Arg Asp Ile Ala

450		455		460	
Leu Met Lys Leu Lys Lys	Pro Val Ala Phe Ser	Asp Tyr Ile His Pro			
465		470		475	480
Val Cys Leu Pro Asp Arg	Glu Thr Ala Ala Ser	Leu Leu Gln Ala Gly			
	485		490		495
Tyr Lys Gly Arg Val Thr	Gly Trp Gly Asn Leu	Lys Glu Thr Trp Thr			
	500		505		510
Ala Asn Val Gly Lys Gly	Gln Pro Ser Val Leu	Gln Val Val Asn Leu			
	515		520		525
Pro Ile Val Glu Arg Pro	Val Cys Lys Asp Ser	Thr Arg Ile Arg Ile			
	530		535		540
Thr Asp Asn Met Phe Cys	Ala Gly Tyr Lys Pro	Asp Glu Gly Lys Arg			
545		550		555	560
Gly Asp Ala Cys Glu Gly	Asp Ser Gly Gly Pro	Phe Val Met Lys Ser			
	565		570		575
Pro Phe Asn Asn Arg Trp	Tyr Gln Met Gly Ile	Val Ser Trp Gly Glu			
	580		585		590
Gly Cys Asp Arg Asp Gly	Lys Tyr Gly Phe Tyr	Thr His Val Phe Arg			
	595		600		605
Leu Lys Lys Trp Ile Gln	Lys Val Ile Asp Gln	Phe Gly Glu			
610		615		620	



# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP2004/051173

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/56 C12N9/50 C12N9/64 C12N9/00 A61K38/48

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	WO 00/78332 A (BJARNASON JON BRAGI) 28 December 2000 (2000-12-28) page 8, line 6 - page 11, line 5	1-67, 77-82
Y	AGGARWAL B B ET AL: "HUMAN TUMOR NECROSIS FACTOR PRODUCTION, PURIFICATION, AND CHARACTERIZATION" JOURNAL OF BIOLOGICAL CHEMISTRY, THE AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, INC.,, US, vol. 260, no. 4, 25 February 1985 (1985-02-25), pages 2345-2354, XP000654946 ISSN: 0021-9258 page 2345, left-hand column, line 1 - page 2345, right-hand column, line 4 figure 7 ----- -/-	1-67, 77-82

☒ Further documents are listed in the continuation of box C

☒ Patent family members are listed in annex

### \* Special categories of cited documents

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*8\* document member of the same patent family

Date of the actual completion of the international search:

10 November 2004

Date of mailing of the international search report

08/12/2004

Name and mailing address of the ISA

European Patent Office, P B 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx 31 651 epo nl  
Fax (+31-70) 340-3016

Authorized officer

Seroz, T

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/051173

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passage*	Relevant to claim No
Y	<p>VAN KESSEL K P ET AL: "Inactivation of recombinant human tumor necrosis factor-alpha by proteolytic enzymes released from stimulated human neutrophils." JOURNAL OF IMMUNOLOGY (BALTIMORE, MD. : 1950) 1 DEC 1991, vol. 147, no. 11, 1 December 1991 (1991-12-01), pages 3862-3868, XP002304822 ISSN: 0022-1767 page 3867, left-hand column, paragraph 2; figure 2</p>	1-67, 77-82
Y	<p>XU Y ET AL: "Mutational analysis of the primary substrate specificity pocket of complement factor B. Asp(226) is a major structural determinant for p(1)-Arg binding." THE JOURNAL OF BIOLOGICAL CHEMISTRY. 7 JAN 2000, vol. 275, no. 1, 7 January 2000 (2000-01-07), pages 378-385, XP002304823 ISSN: 0021-9258 page 381, left-hand column, last paragraph - page 382, left-hand column, paragraph 1 page 385, left-hand column</p>	1,2, 68-71, 79-82
Y	<p>FAUTREL B ET AL: "INTERET DES MOLECULES ANTI-TNF-ALPHA DANS LE MALADIES INFLAMMATOIRES ET INFECTIEUSES" REVUE DE MEDECINE INTERNE, CMR, ASNIERES, FR, vol. 21, no. 10, 2000, pages 872-888, XP000965586 ISSN: 0248-8663 page 873, left-hand column, paragraph 1</p>	1-67, 77-82
P,Y	<p>RUGGLES SANDRA WAUGH ET AL: "Characterization of structural determinants of granzyme B reveals potent mediators of extended substrate specificity." THE JOURNAL OF BIOLOGICAL CHEMISTRY. 16 JUL 2004, vol. 279, no. 29, 16 July 2004 (2004-07-16), pages 30751-30759, XP002304824 ISSN: 0021-9258 page 30751, right-hand column, paragraph 1 figure 5</p>	1,2, 68-71, 79-82

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP2004/051173

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication where appropriate, of the relevant passage	Relevant to claim No.
A	<p>ALTAMIRANO M M ET AL: "Directed evolution of new catalytic activity using the alpha/beta-barrel scaffold"  NATURE, MACMILLAN JOURNALS LTD. LONDON, GB,  vol. 403, no. 6770,  10 February 2000 (2000-02-10), pages 617-622, XP002173865  ISSN: 0028-0836  page 618, right-hand column, last paragraph - page 619, left-hand column, paragraph 2; figure 3  -----</p>	1-82

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP2004/051173

## Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.  
because they relate to subject matter not required to be searched by this Authority, namely  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest
- ☐ No protest accompanied the payment of additional search fees

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box II.1

Although claim 79 is directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound.

Although claims 81 and 82 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound.

-----

## Continuation of Box II.2

Present claims 1-5, 7-71, 74-82 relate to a product defined by reference to a desirable characteristic or property, namely, the capability to hydrolyse defined substrates at defined positions.

An attempt is made to define the product by reference to a result to be achieved. This lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the engineered trypsin, subtilisin E, human pepsin and human caspase 7.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP2004/051173

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 0078332	A	28-12-2000	AT	278417 T	15-10-2004
			AU	4947800 A	09-01-2001
			CA	2377357 A1	28-12-2000
			CN	1356907 T	03-07-2002
			EP	1202743 A2	08-05-2002
			WO	0078332 A2	28-12-2000
			JP	2003502071 T	21-01-2003
			NO	20016159 A	17-12-2001
			NZ	516632 A	30-04-2004
			PL	352318 A1	11-08-2003
			US	2002141987 A1	03-10-2002

---